

Table I. Table of nuclear and decay properties**EXPLANATION OF TABLE**

Data are presented in groups ordered according to increasing mass number A .

Nuclide	Nuclidic name: mass number $A = N + Z$ and element symbol (for $Z > 109$ see Section 2). Element indications with suffix ‘ m ’, ‘ n ’, ‘ p ’ or ‘ q ’ indicate assignments to excited isomeric states (defined, see text, as upper states with half-lives larger than 100 ns). Suffixes ‘ p ’ and ‘ q ’ indicate also non-isomeric levels, of use in the AME2003. Suffix ‘ r ’ indicates a state from a proton resonance occurring in (p, γ) reactions (e.g. $^{28}\text{Si}^r$). Suffix ‘ x ’ applies to mixtures of levels (with relative ratio R , given in the ‘Half-life’ column), e.g. occurring in spallation reactions (indicated ‘spmix’ in the ‘ J^π ’ column) or fission (‘fsmix’).
Mass excess	Mass excess [$M(\text{in u}) - A$], in keV, and its one standard deviation uncertainty as given in the ‘Atomic Mass Evaluation’ (AME2003, second part of this volume). Rounding policy: in cases where the furthest-left significant digit in the error is larger than 3, values and errors are rounded off, but not to more than tens of keV. (Examples: $2345.67 \pm 2.78 \rightarrow 2345.7 \pm 2.8$, $2345.67 \pm 4.68 \rightarrow 2346 \pm 5$, but $2346.7 \pm 468.2 \rightarrow 2350 \pm 470$). # in place of decimal point: value and uncertainty derived not from purely experimental data, but at least partly from systematic trends (cf. AME2003).
Excitation energy	For excited isomers only: energy difference, in keV, between levels adopted as higher level isomer and ground state isomer, and its one standard deviation uncertainty, as given in AME2003 when derived from the AME, otherwise as given by ENSDF. The rounding policy is the same as for the mass excess (see above). # in place of decimal point: value and uncertainty derived from systematic trends. The excitation energy is followed by its origin code when derived from a method other than γ -ray spectrometry: MD Mass doublet RQ Reaction energy difference AD α energy difference BD β energy difference p proton decay XL L X-rays Nm estimated value derived with help of Nilsson model When the existence of an isomer is questionable the following codes are used: EU existence of isomer is under discussion (e.g. $^{141}\text{Tb}^m$). If existence is strongly doubted, no excitation energy and no mass are given. They are replaced by the mention “non-existent” (e.g. $^{138}\text{Pm}^n$). RN isomer is proved not to exist (e.g. $^{184}\text{Lu}^m$). Excitation energy and mass are replaced by the mention “non-existent”. Remark: codes EU and RN are also used when the discovery of a nuclide (e.g. ^{260}Fm) is questioned. In this case however we always give an estimate, derived from systematic trends, for the ground state mass. Isomeric assignment: * In case the uncertainty σ on the excitation energy E is larger than half that energy ($\sigma > E/2$), these quantities are followed by an asterisk (e.g. ^{130}In and $^{130}\text{In}^m$). & In case the ordering of the ground- and isomeric-states are reversed compared to ENSDF, an ampersand sign is added (e.g. ^{90}Tc and $^{90}\text{Tc}^m$).

- Half-life s = seconds; m = minutes; h = hours; d = days; y = years;
 1 y = 31 556 926 s or 365.2422 d
 adopted values for NUBASE (see text)
 STABLE = stable nuclide or nuclide for which no finite value for half-life
 has been found.
 # value estimated from systematic trends in neighboring nuclides with the same Z
 and N parities.
 subunits:
 ms: 10^{-3} s millisecond ky: 10^3 y kiloyear
 μ s: 10^{-6} s microsecond My: 10^6 y megayear
 ns: 10^{-9} s nanosecond Gy: 10^9 y gigayear
 ps: 10^{-12} s picosecond Ty: 10^{12} y terayear
 fs: 10^{-15} s femtosecond Py: 10^{15} y petayear
 as: 10^{-18} s attosecond Ey: 10^{18} y exayear
 zs: 10^{-21} s zeptosecond Zy: 10^{21} y zettayear
 ys: 10^{-24} s yoctosecond Yy: 10^{24} y yottayear
 For isomeric mixtures: R is the production ratio of excited isomeric state to ground-state.
- J^π Spin and parity:
 () uncertain spin and/or parity.
 # values estimated from systematic trends in neighboring nuclides with the same Z
 and N parities.
 high high spin.
 low low spin.
 am same J^π as α -decay parent;
 For isomeric mixtures: mix (spmix and fsmix if coming from spallation and fission respec-
 tively).
- Ens Year of the archival file of the ENSDF
 (in order to reduce the width of the Table, the two digits for the centuries are omitted).
- Reference Reference keys:
 (in order to reduce the width of the Table, the two digits for the centuries are omitted; at
 the end of this volume however, the full reference key-number is given: 1992Pa05 and not
 92Pa05)
 92Pa05 Updates to ENSDF derived from regular journal. These keys are taken from
 Nuclear Data Sheets. Where not yet available, the style 03Ya.1 is provisionally
 adopted.
 95Am.A Updates to ENSDF derived from abstract, preprint, private communication, con-
 ference, thesis or annual report.
 ABBW Re-interpretation by the present authors.
 The reference key-numbers are followed by one, two or three letter codes which specifies
 the added or modified physical quantities:
 T for half-life
 J for spin and/or parity
 E for the isomer excitation energy
 D for decay mode and/or intensity
 I for identification

Decay modes and intensities Decay modes followed by their intensities (in %), and their one standard deviation uncertainties. The special notation 1.8e-12 stands for 1.8×10^{-12} .
 The uncertainties are given - only in this field - in the ENSDF-style: $\alpha=25.9\ 23$ stands for $\alpha=25.9 \pm 2.3\ %$

The ordering is according to decreasing intensities.

α	α emission
p 2p	proton emission 2-proton emission
n 2n	neutron emission 2-neutron emission
ε	electron capture
e^+	positron emission
β^+	β^+ decay ($\beta^+ = \varepsilon + e^+$)
β^-	β^- decay
$2\beta^-$	double β^- decay
$2\beta^+$	double β^+ decay
β^-n	β^- delayed neutron emission
β^-2n	β^- delayed 2-neutron emission
β^+p	β^+ delayed proton emission
β^+2p	β^+ delayed 2-proton emission
$\beta^- \alpha$	β^- delayed α emission
$\beta^+ \alpha$	β^+ delayed α emission
β^-d	β^- delayed deuteron emission
IT	internal transition
SF	spontaneous fission
β^+SF	β^+ delayed fission
β^-SF	β^- delayed fission
^{24}Ne	heavy cluster emission
...	list is continued in a remark, at the end of the A-group

For long-lived nuclides:

IS Isotopic abundance

* A remark on the corresponding nuclide is given below the block of data corresponding to the same A.

Remarks. For nuclides indicated with an asterix at the end of the line, remarks have been added. They are collected in groups at the end of each block of data corresponding to the same A. They start with a code letter, like the ones following the reference key-number, as given above, indicating to which quantity the remark applies. They give:

- i) Continuation for the list of decays. In this case, the remark starts with three dots.
- ii) Information explaining how a value has been derived.
- iii) Reasons for changing a value or its uncertainty as given by the authors or for rejecting it.
- iv) Complementary references for updated data.
- v) Separate values entering an adopted average.

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^1_0n	8071.3171	0.0005	613.9 s	0.6	1/2 ⁺	00 02PaDG T	β^- =100	
^1_1H	7288.9705	0.0001	STABLE		1/2 ⁺	00 98Ro45 D	IS=99.9885 70	
* ^1_1H	D : all isotopic abundances in NUBASE are from 98Ro45							**
^2_1H	13135.7216	0.0003	STABLE		1 ⁺	99	IS=0.0115 70	
^3_1H	14949.8060	0.0023	12.32 y	0.02	1/2 ⁺	00	β^- =100	
^3_2He	14931.2148	0.0024	STABLE		1/2 ⁺	98	IS=0.000137 3	
^3_3Li	28670#	2000#	RN	p-unstable		98	p ?	
^4_1H	25900	100	139 ys	10	2 ⁻	98 03Me11 T	n=100	
^4_2He	2424.9156	0.0001	STABLE		0 ⁺	98	IS=99.999863 3	
^4_3Li	25320	210	91 ys	9	2 ⁻	98 65Ce02 T	p=100	
* ^4_1H	T : width=3.28(0.23) MeV; also 91Go19=4.7(1.0) outweighed, not used							**
^5_1H	32890	100	> 910 ys		(1/2 ⁺)	02 03Go11 T	2n=100	
^5_2He	11390	50	700 ys	30	3/2 ⁻	02	n=100	
^5_3Li	11680	50	370 ys	30	3/2 ⁻	02	p=100	
^5_4Be	38000#	4000#			1/2 ⁺ #	02	p ?	
* ^5_1H	T : from width < 0.5 MeV; at variance with 01Ko52=280(50)ys, width=1.9(0.4)							**
* ^5_2He	T : (same authors) but with instrumental resolution=1.3 MeV							**
* ^5_3Li	T : others 91Go19=66(25) ys 95Al31=110 ys probably for higher state							**
* ^5_4Be	J : from angular distribution consistent with $l = 0$							**
^6_1H	41860	260	290 ys	70	2 ⁻ #	02	n ?; 3n ?	
^6_2He	17595.1	0.8	806.7 ms	1.5	0 ⁺	02 90Ri01 D	β^- =100; β^- d=0.00028 5	
^6_3Li	14086.793	0.015	STABLE		1 ⁺	02	IS=7.59 4	
^6_4Be	18375	5	5.0 zs	0.3	0 ⁺	02	2p=100	
^6_5B	43600#	700#	p-unstable#		2 ⁻ #		2p ?	
^7_1H	49140#	1010#	23 ys	6	1/2 ⁺ #	03Ko11 T	2n ?	
^7_2He	26101	17	2.9 zs	0.5	(3/2 ⁻)	03 02Me07 T	n=100	
^7_3Li	14908.14	0.08	STABLE		3/2 ⁻	03	IS=92.41 4	
^7_4Be	15770.03	0.11	53.22 d	0.06	3/2 ⁻	03	ϵ =100	
^7_5B	27870	70	350 ys	50	(3/2 ⁻)	03	p=100	
* ^7_1H	T : from estimated width 20(5) MeV in Fig. 5							**
* ^7_2He	T : from 159(28) keV, average 02Me07=150(80) 69St02=160(30)							**
^8_2He	31598	7	119.0 ms	1.5	0 ⁺	99 88Aj01 D	β^- =100; β^- n=16 1; β^- t=0.9 1	
^8_3Li	20946.84	0.09	840.3 ms	0.9	2 ⁺	99 90Sa16 T	β^- =100; β^- α =100	
^8_4Be	4941.67	0.04	67 as	17	0 ⁺	99	α =100	
^8_5B	22921.5	1.0	770 ms	3	2 ⁺	99 88Aj01 D	β^+ =100; β^+ α =100	
^8_6C	35094	23	2.0 zs	0.4	0 ⁺	99	2p=100	
* ^8_2He	D : β^- n intensity is from 88Aj01; β^- t intensity from 86Bo41							**
* ^8_3Li	D : β^- decay to first 2 ⁺ state in ^8Be , which decays 100% in 2 α							**
* ^8_5B	D : β^+ to 2 excited states in ^8Be , then α and γ , but not to ^8Be ground-state							**
^9_2He	40939	29	7 zs	4	1/2 ^(-#)	99 99Bo26 T	n=100	
^9_3Li	24954.3	1.9	178.3 ms	0.4	3/2 ⁻	99 95Re.A D	β^- =100; β^- n=50.8 2	
^9_4Be	11347.6	0.4	STABLE		3/2 ⁻	99	IS=100.	
^9_5B	12415.7	1.0	800 zs	300	3/2 ⁻	99	p=100	
^9_6C	28910.5	2.1	126.5 ms	0.9	(3/2 ⁻)	99 88Aj01 D	β^+ =100; β^+ p=23; β^+ α =17	
* ^9_2He	T : derived from width 100(60) keV J : from 01Ch31							**
* ^9_3Li	D : also 92Te03 β^- n=51(1)% 81La11=49(5) outweighed, not used							**
* ^9_5B	D : β^+ =12% and 11% to 2 excited p-emitting states in ^9B , and 17% to α emitter							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{10}He	48810	70	2.7 zs	1.8	0^+	99 94Os04 T	2n=100	*
^{10}Li	33051	15	2.0 zs	0.5	$(1^-, 2^-)$	99 94Yo01 TJ	n=100	
$^{10}\text{Li}^m$	33250	40	200 40 RQ	3.7	zs 1.5	97Zi04 T	IT=100	*
$^{10}\text{Li}^n$	33530	40	480 40 RQ	1.35	zs 0.24	99 94Yo01 T	IT=100	*
^{10}Be	12606.7	0.4	1.51 My	0.06	0^+	99	β^- =100	
^{10}B	12050.7	0.4	STABLE		3^+	99	IS=19.9 7	
^{10}C	15698.7	0.4	19.290 s	0.012	0^+	99 90Ba02 T	β^+ =100	
^{10}N	38800	400	200 ys	140	(2^-)	99 02Le16 TJ	p ?	
* ^{10}He	D : most probably 2 neutron emitter from $S_{2n} = -1070(70)$ keV							**
* $^{10}\text{Li}^m$	T : average 97Zi04=120(+100-50) 94Yo01=100(70) keV							**
* $^{10}\text{Li}^n$	T : average 94Yo01=358(23) 93Bo03=150(70) keV, Birge ratio $B=2.8$							**
^{11}Li	40797	19	8.75 ms	0.14	$3/2^-$	00 97Mo35 T	β^- =100; β^- n=84.9 8; ...	*
^{11}Be	20174	6	13.81 s	0.08	$1/2^+$	00 81Al03 D	β^- =100; β^- n=2.9 4	
^{11}B	8667.9	0.4	STABLE		$3/2^-$	00	IS=80.1 7	
^{11}C	10650.3	1.0	20.39 m	0.02	$3/2^-$	00	β^+ =100	
^{11}N	24300	50	590 ys	210	$1/2^+$	00 03Gu06 T	p=100	*
$^{11}\text{N}^m$	25040	80	740 60	690	ys 80	1/2-	96Ax01 ETJ	p=100
* ^{11}Li	D : ... ; β^- 2n=4.1 4; β^- 3n=1.9 2; β^- n α =1.00 6; β^- t=0.014 3; β^- d=0.013 5							**
* ^{11}Li	D : β^- n, β^- 2n and β^- 3n intensities are from 89Ha.B's evaluation;							**
* ^{11}Li	D : β^- n α intensity is from 84La27; β^- d intensity from 96Mu19;							**
* ^{11}Li	D : β^- t: average 84La27=0.010(4)% 96Mu19=0.020(5)%							**
* ^{11}Li	T : average 97Mo35=8.99(0.10) 96Mu19=8.2(0.2) 95Re.A=8.4(0.2)							**
* ^{11}Li	T : 81Bj01=8.83(0.12) and 74Ro31=8.5(0.2)							**
* ^{11}N	T : unweighed average 03Gu06=0.24(0.24) 00Ma62=1.44(0.2) MeV 00OI01=0.4(0.1)							**
* ^{11}N	T : and 96Ax01=0.99(0.20) MeV (Birge ratio $B=3.03$)							**
^{12}Li	50100#	1000#	< 10 ns			00 74Bo05 I	n ?	
^{12}Be	25077	15	21.50 ms	0.04	0^+	00 01Be53 T	β^- =100; β^- n=0.50 3	*
^{12}B	13368.9	1.4	20.20 ms	0.02	1^+	00 66Sc23 D	β^- =100; β^- n=1.6 3	
^{12}C	0.0	0.0	STABLE		0^+	00	IS=98.93 8	
^{12}N	17338.1	1.0	11.000 ms	0.016	1^+	00 66Sc23 D	β^+ =100; β^+ n=3.5 5	
^{12}O	32048	18	580 ys	30	0^+	00 95Kr03 T	2p=60 30; β^+ ?	
* ^{12}Be	D : from 99Be53; also 95Re.A=0.52 9% outweighed, not used							**
^{13}Be	33250	70	0.5 ns	0.1	$(1/2^+)$	01Th01 TJ	n ?	
$^{13}\text{Be}^p$	33950	90	700 120 RQ	2.7	zs 1.8	$(1/2^-)$ 00		
$^{13}\text{Be}^q$	35160	50	1910 90 RQ			$(5/2^+)$		
^{13}B	16562.2	1.1	17.33 ms	0.17	$3/2^-$	00	β^- =100; β^- n=0.28 4	
^{13}C	3125.0113	0.0009	STABLE		$1/2^-$	01	IS=1.07 8	
^{13}N	5345.48	0.27	9.965 m	0.004	$1/2^-$	00	β^+ =100	
^{13}O	23112	10	8.58 ms	0.05	$(3/2^-)$	00 70Es03 D	β^+ =100; β^+ p=10.9 20	
^{14}Be	39950	130	4.35 ms	0.17	0^+	01 02Je11 D	β^- =100; β^- n=98 2; ...	*
$^{14}\text{Be}^p$	41470	60	1520 150			(2^+) 95Bo10		
^{14}B	23664	21	12.5 ms	0.5	2^-	01 95Re.A D	β^- =100; β^- n=6.04 23	
^{14}C	3019.893	0.004	5.70 ky	0.03	0^+	01	β^- =100	
^{14}N	2863.4170	0.0006	STABLE		1^+	01	IS=99.632 7	
^{14}O	8007.36	0.11	70.598 s	0.018	0^+	01 01Ga59 T	β^+ =100	*
^{14}F	32660#	400#			2^- #		p ?	
* ^{14}Be	D : ... ; β^- 2n=0.8 08; β^- 3n=0.2 2; β^- t=0.02 1; β^- n α <0.004							**
* ^{14}Be	D : supersedes 99Be53, same group							**
* ^{14}O	T : average 01Ga59=70.560(0.049) 78Wi04=70.613(0.025) 73Cl12=70.590(0.030)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁵ Be	49800#	500#	< 200 ns			03Ba47 I	n ?
¹⁵ B	28972	22	9.87 ms	0.07	3/2 ⁻	93 95Re.A TD	β^- =100; β^- n=93.6 12; β^- 2n=0.4 2 *
¹⁵ C	9873.1	0.8	2.449 s	0.005	1/2 ⁺	94	β^- =100
¹⁵ N	101.4380	0.0007	STABLE		1/2 ⁻	94	IS=0.368 7
¹⁵ O	2855.6	0.5	122.24 s	0.16	1/2 ⁻	94	β^+ =100
¹⁵ F	16780	130	410 ys	60	(1/2 ⁺)	93 01Ze.A T	p=100 *
* ¹⁵ B	D : β^- 2n intensity is from 89Re.A		J : given in 91Aj01				**
* ¹⁵ B	T : four other outweighed results, see ENSDF'93, ranging 10.1 - 10.8 ms **						
* ¹⁵ F	T : average 01Ze.A=1.23(0.22)MeV 78Be16=1.2(0.3) 78Ke06=0.8(0.3) **						
¹⁶ Be	57680#	500#	< 200 ns		0+	03Ba47 I	2n ? *
¹⁶ B	37080	60	< 190 ps		0-	99	n ?
¹⁶ C	13694	4	747 ms	8	0+	99 89Re.A D	β^- =100; β^- n=97.9 23
¹⁶ N	5683.7	2.6	7.13 s	0.02	2-	99 74Ne10 D	β^- =100; β^- α =0.00100 7
¹⁶ O	-4737.0014	0.0001	STABLE		0+	99	IS=99.757 16
¹⁶ F	10680	8	11 zs	6	0-	99	p=100
¹⁶ Ne	23996	20	9 zs		0+	99	2p=100
* ¹⁶ Be	I : 100 events expected, none observed **						
¹⁷ B	43770	170	5.08 ms	0.05	(3/2 ⁻)	99 88Du09 D	β^- =100; β^- n=63 1; ... *
¹⁷ C	21039	17	193 ms	5	(3/2 ⁺)	99 01Ma08 J	β^- =100; β^- n=28.4 13 *
¹⁷ N	7871	15	4.173 s	0.004	1/2 ⁻	99 94Do08 D	β^- =100; β^- n=95 1; ... *
¹⁷ O	-808.81	0.11	STABLE		5/2 ⁺	99	IS=0.038 1
¹⁷ F	1951.70	0.25	64.49 s	0.16	5/2 ⁺	99	β^+ =100
¹⁷ Ne	16461	27	109.2 ms	0.6	1/2 ⁻	99 88Bo39 D	β^+ =100; β^+ p=96.0 9; β^+ α =2.7 9
* ¹⁷ B	D : ... ; β^- 2n=11 7; β^- 3n=3.5 7; β^- 4n=0.4 3 **						
* ¹⁷ C	T : average 95Sc03=193(6) 95Re.A=188(10) 86Cu01=202(17) **						
* ¹⁷ C	D : β^- n intensity is from 95Re.A **						
* ¹⁷ N	D : ... ; β^- α =0.0025 4 **						
¹⁸ B	52320#	800#	< 26 ns		4-#	93Po.A I	n ?
¹⁸ C	24930	30	92 ms	2	0+	96	β^- =100; β^- n=31.5 15
¹⁸ N	13114	19	622 ms	9	1-	96 95Re.A D	β^- =100; β^- n=10.9 9; ... *
¹⁸ O	-781.5	0.6	STABLE		0+	96	IS=0.205 14
¹⁸ F	873.7	0.5	109.771 m	0.020	1+	96 02Un02 T	β^+ =100
¹⁸ F ^m	1995.1	0.5	1121.36	0.15	234 ns		5+
¹⁸ Ne	5317.17	0.28	1.672 s	0.008	0+	96	β^+ =100
¹⁸ Na	24190	50	1.3 zs	0.4	1-#	01Ze.A TD	p=?; β^+ ?
* ¹⁸ N	D : ... ; β^- α =12.2 6 **						
* ¹⁸ N	D : β^- n intensity is from 95Re.A; β^- α intensity from 89Zn04 **						
* ¹⁸ N	T : average 99Og03=620(14) 82O101=624(12) **						
¹⁹ B	59360#	400#	2.92 ms	0.13	3/2 ⁻ #	96 03Yo02 T	β^- =100; β^- n \approx 75; ... *
¹⁹ C	32420	100	46.2 ms	2.3	(1/2 ⁺)	96 88Du09 TD	β^- =100; β^- n=47.3; ... *
¹⁹ N	15862	16	271 ms	8	(1/2 ⁻)	96	β^- =100; β^- n=54.6 14 *
¹⁹ O	3334.9	2.8	26.464 s	0.009	5/2 ⁺	96 94It.A T	β^- =100
¹⁹ F	-1487.39	0.07	STABLE		1/2 ⁺	96	IS=100
¹⁹ Ne	1751.44	0.29	17.296 s	0.005	1/2 ⁺	96 94Ko.A T	β^+ =100
¹⁹ Na	12927	12	< 40 ns		5/2 ⁺ #	96 93Po.A I	p=100 *
¹⁹ Mg	33040	250			1/2 ⁻ #	96	2p ?
* ¹⁹ B	D : ... ; β^- 2n \approx 25 **						
* ¹⁹ B	T : others: 99Re16=4.5(1.5) 98Yo06=3.3(0.2) statistics + 2.0 systematics estimated by NUBASE **						
* ¹⁹ B	D : deduced from $P_n = \beta^- n + 2 \times \beta^- 2n + \dots = 125(32)\%$ in 98Yo06 and assuming **						
* ¹⁹ B	D : $\beta^- n + \beta^- 2n = 100\%$ **						
* ¹⁹ C	D : ... ; β^- 2n=7 3 **						
* ¹⁹ C	T : average 88Du09=49(4) 95Re.A=44(4) 95Oz02=45.5(4.0) **						
* ¹⁹ C	J : from 01Ma08, 99Na27 and 95Ba28 **						
* ¹⁹ N	J : 95Oz02=(1/2, 3/2, 5/2)- 89Ca25=(1/2-) **						
* ¹⁹ Na	D : most probably proton emitter from $S_p = -333(12)$ keV **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)		
²⁰ C	37560	240	16 ms	3	0 ⁺	98 90Mu06 T	β^- =100; β^- n=72 14	*	
²⁰ N	21770	60	130 ms	7		98 95Re.A TD	β^- =100; β^- n=57.0 25		
²⁰ O	3797.5	1.1	13.51 s	0.05	0 ⁺	98	β^- =100		
²⁰ F	-17.40	0.08	11.163 s	0.008	2 ⁺	98 98Ti06 T	β^- =100		
²⁰ Ne	-7041.9313	0.0018	STABLE		0 ⁺	98	IS=90.48 3		
²⁰ Na	6848	7	447.9 ms	2.3	2 ⁺	98 89Cl02 D	β^+ =100; β^+ α =25.0 4		
²⁰ Mg	17570	27	90 ms	6	0 ⁺	98 95Pi03 TD	β^+ =100; β^+ p=30.4 16	*	
* ²⁰ C	T : average 90Mu06=14(+6-5) 95Re.A 16.7(3.5)							**	
* ²⁰ Mg	T : average 95Pi03=95(3) 92Go10=82(4), with Birge ratio B=2.6							**	
²¹ C	45960#	500#	< 30 ns		1/2 ⁺ #	00 93Po.A I	n ?		
²¹ N	25250	100	87 ms	6	1/2 ⁻ #	00	β^- =100; β^- n=80 6		
²¹ O	8063	12	3.42 s	0.10	(1,3,5)/2 ⁺	00	β^- =100		
²¹ F	-47.6	1.8	4.158 s	0.020	5/2 ⁺	00	β^- =100		
²¹ Ne	-5731.78	0.04	STABLE		3/2 ⁺	00	IS=0.27 1		
²¹ Na	-2184.2	0.7	22.49 s	0.04	3/2 ⁺	00	β^+ =100		
²¹ Mg	10911	16	122 ms	2	(5/2,3/2) ⁺	00	β^+ =100; β^+ p=32.6 10; ...	*	
²¹ Al	26120#	300#	< 35 ns		1/2 ⁺ #	00 93Po.A I	p ?		
* ²¹ Mg	D : ... ; β^+ α <0.5							**	
* ²¹ Mg	J : from mirror ²¹ F, there is a preference for 5/2 ⁺							**	
²² C	53280#	900#	6.2 ms	1.3	0 ⁺	00 03Yo02 TD	β^- =100; β^- n=99 39; ...	*	
²² N	32040	190	13.9 ms	1.4		00 03Yo02 T	β^- =100; β^- n=35 5	*	
²² O	9280	60	2.25 s	0.15	0 ⁺	00	β^- =100; β^- n<22		
²² F	2793	12	4.23 s	0.04	4 ⁺ , (3 ⁺)	00	β^- =100; β^- n<11		
²² Ne	-8024.715	0.018	STABLE		0 ⁺	00	IS=9.25 3		
²² Na	-5182.4	0.4	2.6019 y	0.0004	3 ⁺	00	β^+ =100		
²² Na ^m	-4599.4	0.4	583.03	0.09	244 ns	6	1 ⁺	00 IT=100	
²² Mg	-397.0	1.3	3.857 s	0.009	0 ⁺	00	β^+ =100		
²² Al	18180#	90#	59 ms	3	(3) ⁺	00 97B103 D	β^+ =100; β^+ p=44 3; ...	*	
²² Si	32160#	200#	29 ms	2	0 ⁺	00 96B111 D	β^+ =100; β^+ p=32 4	*	
* ²² C	D : ... ; β^- 2n ? D : from 98Yo06							**	
* ²² N	D : from 90Mu06							**	
* ²² Al	D : ... ; β^+ 2p=0.9 5; β^+ α =0.31 9							**	
²³ N	38400#	300#	14.5 ms	2.4	1/2 ⁻ #	00 98Yo06 T	β^- =100; β^- n=80 21; β^- 2n ?	*	
²³ O	14610	120	90 ms	40	1/2 ⁺ #	00 90Mu06 T	β^- =100; β^- n=31 7		
²³ F	3330	80	2.23 s	0.14	(3/2,5/2) ⁺	00	β^- =100; β^- n<14		
²³ Ne	-5154.05	0.10	37.24 s	0.12	5/2 ⁺	00	β^- =100		
²³ Na	-9529.8536	0.0027	STABLE		3/2 ⁺	00	IS=100		
²³ Mg	-5473.8	1.3	11.317 s	0.011	3/2 ⁺	00	β^+ =100		
²³ Al	6770	19	470 ms	30	5/2 ⁺ #	00 95Ti08 D	β^+ =100; β^+ p=8 4	*	
²³ Si	23770#	200#	42.3 ms	0.4	3/2 ⁺ #	00 97B104 TD	β^+ =100; β^+ p \approx 88; ...	*	
* ²³ N	T : statistical error 1.4, systematics 2.0 estimated by NUBASE							**	
* ²³ Al	D : β^+ p=3.5(1.9)% from the IAS. Total=3.5 \times 4.8/2.2=7.6%							**	
* ²³ Si	D : ... ; β^+ 2p=3.6 3							**	
²⁴ N	47540#	400#	< 52 ns			00 93Po.A I	n ?		
²⁴ O	19070	240	65 ms	5	0 ⁺	00	β^- =100; β^- n=18 6		
²⁴ F	7560	70	400 ms	50	(1,2,3) ⁺	00	β^- =100; β^- n<5.9		
²⁴ Ne	-5951.5	0.4	3.38 m	0.02	0 ⁺	00	β^- =100		
²⁴ Na	-8418.11	0.08	14.9590 h	0.0012	4 ⁺	00	β^- =100		
²⁴ Na ^m	-7945.90	0.08	472.207	0.009	20.20 ms	0.07	1 ⁺	00 IT \approx 100; β^- =0.05	
²⁴ Mg	-13933.567	0.013	STABLE		0 ⁺	00	IS=78.99 4		
²⁴ Al	-56.9	2.8	2.053 s	0.004	4 ⁺	00	β^+ =100; β^+ α =0.035 6; ...	*	
²⁴ Al ^m	368.9	2.8	425.8	0.1	131.3 ms	2.5	1 ⁺	00 IT=82 3; β^+ =18 3; ...	*
²⁴ Si	10755	19	140 ms	8	0 ⁺	00 98Cz01 D	β^+ =100; β^+ p=37.6 25		
²⁴ P	32000#	500#			1 ⁺ #		p ?; β^+ ?		
* ²⁴ Al	D : ... ; β^+ p=0.0016 3							**	
* ²⁴ Al ^m	D : ... ; β^+ α =0.028 6							**	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)		
²⁵ N	56500#	500#	< 260 ns	1/2 ⁻ #		99Sa06 ID	n ?; 2n ?; β^- =0	*	
²⁵ O	27440#	260#	< 50 ns	3/2 ⁺ #	00	93Po.A I	n ?		
²⁵ F	11270	100	50 ms	6	5/2 ⁺ #	00	β^- =100; β^- n=14 5		
²⁵ Ne	-2108	26	602 ms	8	(3/2) ⁺	00	β^- =100		
²⁵ Na	-9357.8	1.2	59.1 s	0.6	5/2 ⁺	00	β^- =100		
²⁵ Mg	-13192.83	0.03	STABLE		5/2 ⁺	00	IS=10.00 1		
²⁵ Al	-8916.2	0.5	7.183 s	0.012	5/2 ⁺	00	β^+ =100		
²⁵ Si	3824	10	220 ms	3	5/2 ⁺	00	β^+ =100; β^+ p=36.81 5		
²⁵ P	18870#	200#	< 30 ns		1/2 ⁺ #	00	p ?		
* ²⁵ N	D : in 99Sa06 experiment, 240 ²⁵ N events expected, none observed							**	
²⁶ O	35710#	260#	< 40 ns	0 ⁺	00	93Po.A I	2n ?; n=30#; β^- =0	*	
²⁶ F	18270	170	10.2 ms	1.4	1 ⁺	00	β^- =100; β^- n=11 4	*	
²⁶ Ne	430	27	197 ms	1	0 ⁺	00	β^- =100; β^- n=0.13 3		
²⁶ Na	-6862	6	1.077 s	0.005	3 ⁺	00	β^- =100		
²⁶ Mg	-16214.582	0.027	STABLE		0 ⁺	00	IS=11.01 3		
²⁶ Al	-12210.31	0.06	717 ky	24	5 ⁺	00	β^+ =100		
²⁶ Al ^m	-11982.01	0.06	228.305	0.013	6.3452 s	0.0019	0 ⁺	00	
²⁶ Si	-7145	3	2.234 s	0.013	0 ⁺	00	β^+ =100		
²⁶ P	10970#	200#	30 ms	25	(3 ⁺)	00	β^+ =100; β^+ 2p≈1; ...	*	
²⁶ S	25970#	300#	10# ms		0 ⁺		2p ?		
* ²⁶ O	D : in 96Fa01 and 99Sa06, several 100s of ²⁶ O events expected, none observed							**	
* ²⁶ F	T : other not used 99DI01=9.6(0.8): same data							**	
* ²⁶ P	D : ... ; β^+ p≈0.9							**	
²⁷ O	44950#	500#	< 260 ns		3/2 ⁺ #	99Sa06 I	n ?; 2n ?		
²⁷ F	24930	380	4.9 ms	0.2	5/2 ⁺ #	01	β^- =100; β^- n=77 21	*	
²⁷ Ne	7070	110	32 ms	2	3/2 ⁺ #	01	β^- =100; β^- n=2.0 5		
²⁷ Na	-5517	4	301 ms	6	5/2 ⁺	01	β^- =100; β^- n=0.13 4		
²⁷ Mg	-14586.65	0.05	9.458 m	0.012	1/2 ⁺	01	β^- =100		
²⁷ Al	-17196.66	0.12	STABLE		5/2 ⁺	01	IS=100.		
²⁷ Si	-12384.30	0.15	4.16 s	0.02	5/2 ⁺	01	β^+ =100		
²⁷ P	-717	26	260 ms	80	1/2 ⁺	01	β^+ =100; β^+ p=0.07		
²⁷ S	17540#	200#	21 ms	4	(5/2 ⁺)	01	β^+ =100; β^+ 2p=2.0 10;...	*	
* ²⁷ F	T : others not used: 99Re16=6.5(1.1) and 97Ta22=5.3(0.9) outweighed; and							**	
* ²⁷ P	T : 99DI01=5.2(0.3) same data as in 99Re16							**	
* ²⁷ S	D : ... ; β^+ p=?							**	
²⁸ O	53850#	600#	< 100 ns		0 ⁺	98Po.A I	n ?; 2n ?; β^- =0	*	
²⁸ F	33230#	510#	< 40 ns			01	93Po.A I	n ?	
²⁸ Ne	11240	150	18.3 ms	2.2	0 ⁺	01	99Re16 T	β^- =100; β^- n=16 6	*
²⁸ Na	-989	13	30.5 ms	0.4	1 ⁺	01	β^- =100; β^- n=0.58 12		
²⁸ Mg	-15018.6	2.0	20.915 h	0.009	0 ⁺	01	β^- =100		
²⁸ Al	-16850.44	0.13	2.2414 m	0.0012	3 ⁺	01	β^- =100		
²⁸ Si	-21492.7968	0.0018	STABLE		0 ⁺	01	IS=92.2297 7		
²⁸ Si ^r	-8951.55	0.12	12541.25	0.12	RQ		3 ⁺	01	
²⁸ P	-7159	3	270.3 ms	0.5	3 ⁺	01	79Ho27 D	β^+ =100; β^+ p=0.0013 4;...	*
²⁸ S	4070	160	125 ms	10	0 ⁺	01	89Po10 D	β^+ =100; β^+ p=20.7 19	
²⁸ Cl	26560#	500#			1 ⁺ #		p ?		
* ²⁸ O	D : in 97Ta22 and 99Sa06, 11 and 37 ²⁸ O events expected, none observed							**	
* ²⁸ Ne	T : average 99Re16=18(3) 97Ta22=21(5) 92Te03=17(4). Others not used:							**	
* ²⁸ Ne	T : 95Re.A=8.2(2.5) at variance, 99DI01=20(3) same data as in 99Re16							**	
* ²⁸ P	D : ... ; β^+ α =0.00086 25							**	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
²⁹ F	40300#	580#	2.6 ms	0.3	5/2 ⁺ #	01 99Re16 T	β^- =100; β^- n=60.40; ... *	
²⁹ Ne	18060	270	15.6 ms	0.5	3/2 ⁺ #	01 01Be53 D	β^- =100; β^- n=19.4; ... *	
²⁹ Na	2665	13	44.9 ms	1.2	3/2 ⁺ (+)	01 95Re.A D	β^- =100; β^- n=25.9 23 *	
²⁹ Mg	-10619	14	1.30 s	0.12	3/2 ⁺	01	β^- =100	
²⁹ Al	-18215.3	1.2	6.56 m	0.06	5/2 ⁺	01	β^- =100	
²⁹ Si	-21895.046	0.021	STABLE		1/2 ⁺	01	IS=4.6832 5	
²⁹ P	-16952.6	0.6	4.142 s	0.015	1/2 ⁺	01	β^+ =100	
²⁹ S	-3160	50	187 ms	4	5/2 ⁺	01 79Vi01 D	β^+ =100; β^+ p=46.4 10	
²⁹ Cl	13140#	200#	< 20 ns		3/2 ⁺ #	01 93Po.A I	p ?	
* ²⁹ F	D : ... ; β^- 2n ?							**
* ²⁹ F	T : average 99Re16=2.9(0.8) 98No.A=2.6(0.4) 97Ta22=2.4(0.8). Others not							**
* ²⁹ F	T : used: 99Di01=2.4(0.4) same data as in 99Re16							**
* ²⁹ F	D : β^- n from 99Di01=100(80)%							**
* ²⁹ Ne	D : ... ; β^- 2n<2.2							**
* ²⁹ Ne	D : average 01Be53=17.5 99Re16=27.9; other not used: 99Di01=27(9)%, same							**
* ²⁹ Ne	D : data as in 99Re16. β^- 2n limit is from 01Be53							**
* ²⁹ Na	D : β^- n: average 95Re.A=27.1(1.6)% 84La03=21.5(3.0)%							**
³⁰ F	48900#	600#	< 260 ns			99Sa06 I	n ?	
³⁰ Ne	23100	570	5.8 ms	0.2	0 ⁺	01 99Di01 D	β^- =100; β^- n=13.8 *	
³⁰ Na	8361	25	48.4 ms	1.7	2 ⁺	01 99Di01 T	β^- =100; β^- n=30.4; ... *	
³⁰ Mg	-8911	8	335 ms	17	0 ⁺	01 84La03 D	β^- =100; β^- n<0.06	
³⁰ Al	-15872	14	3.60 s	0.06	3 ⁺	01	β^- =100	
³⁰ Si	-24432.928	0.030	STABLE		0 ⁺	01	IS=3.0872 5	
³⁰ P	-20200.6	0.3	2.498 m	0.004	1 ⁺	01	β^+ =100	
³⁰ S	-14063	3	1.178 s	0.005	0 ⁺	01	β^+ =100	
³⁰ Cl	4440#	200#	< 30 ns		3 ⁺ #	01 93Po.A I	p ?	
³⁰ Ar	20080#	300#	< 20 ns		0 ⁺	01 93Po.A I	2p ?	
* ³⁰ Ne	D : from 9(17)%							**
* ³⁰ Na	D : ... ; β^- 2n=1.17 16; β^- α =5.5e-5 20							**
* ³⁰ Na	T : average 99Di01=50(4) 97Ta22=48(5) 84La02=48(2)							**
* ³⁰ P	D : first observed radionuclide, in 1934							**
³¹ F	56290#	600#	1# ms (>260 ns)		5/2 ⁺ #	99Sa06 I	β^- ?; β^- n ?	
³¹ Ne	30840#	900#	3.4 ms	0.8	7/2 ⁻ #	01	β^- =100; β^- n ?	
³¹ Na	12650	210	17.0 ms	0.4	(3/2 ⁺)	01 93K102 J	β^- =100; β^- n=37.5; ... *	
³¹ Mg	-3217	12	230 ms	20	3/2 ⁺	01 95Re.A D	β^- =100; β^- n=6.2 20 *	
³¹ Al	-14954	20	644 ms	25	(5/2, 3/2) ⁺	01	β^- =100; β^- n<1.6 *	
³¹ Si	-22949.01	0.04	157.3 m	0.3	3/2 ⁺	01	β^- =100	
³¹ P	-24440.88	0.18	STABLE		1/2 ⁺	01	IS=100.	
³¹ S	-19044.6	1.5	2.572 s	0.013	1/2 ⁺	01	β^+ =100	
³¹ Cl	-7070	50	150 ms	25	3/2 ⁺	01 85Ay02 D	β^+ =100; β^+ p=0.7 *	
³¹ Ar	11290#	210#	14.4 ms	0.6	5/2 ⁺ (+)	01 00Fy01 T	β^+ =100; β^+ p=63.7; ... *	
* ³¹ Na	D : ... ; β^- 2n=0.9 2; β^- 3n<0.05							**
* ³¹ Na	D : all from 84Gu19							**
* ³¹ Mg	D : strongly conflicting with earlier 84La03=1.7(0.3)%							**
* ³¹ Al	J : from systematics there is a preference for 5/2 ⁺							**
* ³¹ Cl	D : β^+ p=0.44% for 986 keV protons. Total: 165/100×0.44=0.726%							**
* ³¹ Ar	D : ... ; β^+ 2p=7.2 11; β^+ 3p<1.4; β^+ p α <0.38; β^+ α <0.03							**
* ³¹ Ar	D : from 98Ax02							**
* ³¹ Ar	T : average 00Fy01=14.1(0.7) 92Ba01=15.1(+1.3-1.1) J : from 99Th09							**
³² Ne	37280#	800#	3.5 ms	0.9	0 ⁺	01	β^- =100; β^- n ?	
³² Na	19060	360	12.9 ms	0.7	(3 ⁻ , 4 ⁻)	01 93K102 J	β^- =100; β^- n=24.7; ... *	
³² Mg	-955	18	95 ms	16	0 ⁺	01	β^- =100; β^- n=2.4 5	
³² Al	-11060	90	31.7 ms	0.8	1 ⁺	01 95Re.A TD	β^- =100; β^- n=0.7 5	
³² Al ^m	-10100	90	955.7 0.4	200 ns	20	(4 ⁺)	01 96Ro02 ETJ	
³² Si	-24080.91	0.05	132 y	13	0 ⁺	01	β^- =100	
³² Si ^m	-18497.9	1.0	5583.0 1.0	27 ns	2	(5 ⁻)	97Fo01 ETJ	

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
³² P	-24305.22	0.19	14.263 d	0.003	1 ⁺	01 02Un02 T	$\beta^- = 100$	
³² S	-26015.70	0.14	STABLE		0 ⁺	01	IS=94.93 31	
³² Cl	-13330	7	298 ms	1	1 ⁺	01 79Ho27 D	$\beta^+ = 100; \beta^+ \alpha = 0.054 8; \dots$ *	
³² Ar	-2200.2	1.8	98 ms	2	0 ⁺	01	$\beta^+ = 100; \beta^+ p = 43.3$	
³² Ar ^m	3400#	100#	5600#	100#	5 ⁻ #		IT?	
³² K	20420#	500#			1 ⁺ #		p?	
³² K ^m	21370#	510#	950#	100#	4 ⁺ #		p?	
* ³² Na	D: ...; $\beta^- 2n = 8 2$							**
* ³² Na	T: average 98No.A=11.5(0.8) 84La03=13.2(0.4)							**
* ³² Cl	D: ...; $\beta^+ p = 0.026 5$							**
³³ Ne	46000#	800#	< 260 ns		7/2 ⁻ #	02No11 I	n?	
³³ Na	24890	870	8.2 ms	0.2	3/2 ⁺ #	01 02Ra16 TD	$\beta^- = 100; \beta^- n = 47.6; \dots$ *	
³³ Mg	4894	20	90.5 ms	1.6	7/2 ⁻ #	01 02Mo29 T	$\beta^- = 100; \beta^- n = 17.5$	
³³ Al	-8530	70	41.7 ms	0.2	5/2 ⁺ #	01 02Mo29 T	$\beta^- = 100; \beta^- n = 8.5 7$	
³³ Si	-20493	16	6.18 s	0.18	(3/2 ⁺)	01	$\beta^- = 100$	
³³ P	-26337.5	1.1	25.34 d	0.12	1/2 ⁺	01	$\beta^- = 100$	
³³ S	-26585.99	0.14	STABLE		3/2 ⁺	01	IS=0.76 2	
³³ Cl	-21003.4	0.5	2.511 s	0.003	3/2 ⁺	01	$\beta^+ = 100$	
³³ Ar	-9384.1	0.4	173.0 ms	2.0	1/2 ⁺	01	$\beta^+ = 100; \beta^+ p = 38.7 10$	
³³ K	6760#	200#	< 25 ns		3/2 ⁺ #	01 93Po.A I	p?	
* ³³ Ne	T: estimated half-life 1# ms for β^- decay I: also 02Le.A < 1.5 μ s							**
* ³³ Na	D: ...; $\beta^- 2n = 13 3$							**
³⁴ Ne	53120#	810#	1# ms (>1.5 μ s)		0 ⁺	02Le.A I	$\beta^- ?; \beta^- n ?$	
³⁴ Na	32760#	900#	5.5 ms	1.0	1 ⁺	01 ABBW D	$\beta^- = 100; \beta^- 2n \approx 50; \beta^- n \approx 15$ *	
³⁴ Mg	8810	230	20 ms	10	0 ⁺	01	$\beta^- = 100; \beta^- n ?$	
³⁴ Al	-2930	110	56.3 ms	0.5	4 ⁻ #	01 01Nu01 T	$\beta^- = 100; \beta^- n = 12.5 25$ *	
³⁴ Si	-19957	14	2.77 s	0.20	0 ⁺	01	$\beta^- = 100$	
³⁴ P	-24558	5	12.43 s	0.08	1 ⁺	01	$\beta^- = 100$	
³⁴ S	-29931.79	0.11	STABLE		0 ⁺	01	IS=4.29 28	
³⁴ Cl	-24439.78	0.18	1.5264 s	0.0014	0 ⁺	01	$\beta^+ = 100$	
³⁴ Cl ^m	-24293.42	0.18	146.36	0.03	32.00 m	0.04	3 ⁺	$\beta^+ = 55.4 6; IT = 44.6 6$
³⁴ Ar	-18377.2	0.4	845 ms	3	0 ⁺	01	$\beta^+ = 100$	
³⁴ K	-1480#	300#	< 40 ns		1 ⁺ #	01 93Po.A I	p?	
³⁴ Ca	13150#	300#	< 35 ns		0 ⁺	01 93Po.A I	2p?	
* ³⁴ Ne	I: also 02No11 > 260 ns							**
* ³⁴ Na	D: $\beta^- n \approx 15\%; \beta^- 2n \approx 50\%$ estimated from $P_n = \beta^- n + 2 \times \beta^- 2n = 115(20)\%$ in 84La03							**
* ³⁴ Na	D: assuming $\beta^- n / \beta^- 2n = 0.3$ from trends in the ³⁰ Na- ³³ Na series: 26 41 3 4							**
* ³⁴ Al	D: from 95Re.A; strongly conflicting with 89Ba50=27(5)% and 88Mu08=54(12)%							**
* ³⁴ Al	T: also 95Re.A=42(6) ms							**
³⁵ Na	39580#	950#	1.5 ms	0.5	3/2 ⁺ #	01	$\beta^- = 100; \beta^- n = ?$	
³⁵ Mg	16150#	400#	70 ms	40	7/2 ⁻ #	01 95Re.A D	$\beta^- = 100; \beta^- n = 52.46$	
³⁵ Al	-130	180	38.6 ms	0.4	5/2 ⁺ #	01 01Nu01 TD	$\beta^- = 100; \beta^- n = 41.13$ *	
³⁵ Si	-14360	40	780 ms	120	7/2 ⁻ #	01 95Re.A D	$\beta^- = 100; \beta^- n < 5$	
³⁵ P	-24857.7	1.9	47.3 s	0.7	1/2 ⁺	01	$\beta^- = 100$	
³⁵ S	-28846.36	0.10	87.51 d	0.12	3/2 ⁺	01	$\beta^- = 100$	
³⁵ Cl	-29013.54	0.04	STABLE		3/2 ⁺	01	IS=75.78 4	
³⁵ Ar	-23047.4	0.7	1.775 s	0.004	3/2 ⁺	01	$\beta^+ = 100$	
³⁵ K	-11169	20	178 ms	8	3/2 ⁺	01	$\beta^+ = 100; \beta^+ p = 0.37 15$	
³⁵ Ca	4600#	200#	25.7 ms	0.2	1/2 ⁺ #	01	$\beta^+ = 100; \beta^+ p = 95.7 14; \dots$ *	
* ³⁵ Al	T: also 95Re.A=30(4); both strongly conflicting with 89Le16=170(70) and							**
* ³⁵ Al	T: 88Mu08=130(+100-50)							**
* ³⁵ Al	D: also 95Re.A=26(4)% 89Le16=40(10)% and 88Mu08=87(+37-25)%							**
* ³⁵ Ca	D: ...; $\beta^+ 2p = 4.2 3$							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
³⁶ Na	47950#	950#	< 260 ns			02No11 I	n ?	*
³⁶ Mg	21420#	500#	5# ms(>200 ns)	0 ⁺	01	89Gu03 I	β^- ?	
³⁶ Al	5780	210	90 ms	40	01		β^- =100; β^- -n<30	
³⁶ Si	-12480	120	450 ms	60	01	95Re.A D	β^- =100; β^- -n=12.5	
³⁶ P	-20251	13	5.6 s	0.3	4 ⁻ #	01	β^- =100	
³⁶ S	-30664.07	0.19	STABLE		0 ⁺	01	IS=0.02 1	
³⁶ Cl	-29521.86	0.07	301 ky	2	2 ⁺	01	β^- =98.1 1; β^+ =1.9 1	
³⁶ Ar	-30231.540	0.027	STABLE		0 ⁺	01	IS=0.3365 30; 2 β^+ ?	
³⁶ K	-17426	8	342 ms	2	2 ⁺	01	β^+ =100; β^+ p=0.048 14; ...	*
³⁶ Ca	-6440	40	102 ms	2	0 ⁺	01	95Tr02 D β^+ =100; β^+ p=56.8 13	
³⁶ Sc	13900#	500#					p ?	
* ³⁶ Na	I : also 02Le.A < 1.5 μ s							**
* ³⁶ K	D : ... ; β^+ α =0.0034 13							**
³⁷ Na	55280#	960#	1# ms(>1.5 μ s)	3/2 ⁺ #		02Le.A I	β^- ?; β^- -n ?	*
³⁷ Mg	29250#	900#	40# ms(>260 ns)	7/2 ⁻ #	01	96Sa34 I	β^- ?; β^- -n ?	
³⁷ Al	9950	330	20# ms (>1 μ s)	3/2 ⁺ #	01	91Or01 I	β^- ?	
³⁷ Si	-6580	170	90 ms	60	7/2 ⁻ #	01	95Re.A D β^- =100; β^- -n=17 13	
³⁷ P	-18990	40	2.31 s	0.13	1/2 ⁺ #	01	β^- =100	
³⁷ S	-26896.36	0.20	5.05 m	0.02	7/2 ⁻	01	β^- =100	
³⁷ Cl	-31761.53	0.05	STABLE		3/2 ⁺	01	IS=24.22 4	
³⁷ Ar	-30947.66	0.21	35.04 d	0.04	3/2 ⁺	01	ϵ =100	
³⁷ K	-24800.20	0.09	1.226 s	0.007	3/2 ⁺	01	β^+ =100	
³⁷ Ca	-13162	22	181.1 ms	1.0	(3/2 ⁺)	01	95Tr03 D β^+ =100; β^+ p=82.1 7	
³⁷ Sc	2840#	300#			7/2 ⁻ #		p ?	
* ³⁷ Na	I : also 02No11 > 260 ns							**
³⁸ Mg	35000#	500#	1# ms(>260 ns)	0 ⁺	01	97Sa14 I	β^- ?	*
³⁸ Al	16050	730	40# ms(>200 ns)		01	89Gu03 I	β^- ?	
³⁸ Si	-4070	140	90# ms (>1 μ s)	0 ⁺	01	91Zh24 I	β^- ?; β^- -n ?	
³⁸ P	-14760	100	640 ms	140	01	95Re.A D	β^- =100; β^- -n=12.5	
³⁸ S	-26861	7	170.3 m	0.7	0 ⁺	01	β^- =100	
³⁸ Cl	-29798.10	0.10	37.24 m	0.05	2 ⁻	01	β^- =100	
³⁸ Cl ^m	-29126.74	0.10	671.361	0.008	715 ms	3	5 ⁻ 01 IT=100	
³⁸ Ar	-34714.6	0.3	STABLE		0 ⁺	01	IS=0.0632 5	
³⁸ K	-28800.7	0.4	7.636 m	0.018	3 ⁺	01	β^+ =100	
³⁸ K ^m	-28670.2	0.4	130.50	0.28	RQ	923.9 ms	0.6	0 ⁺ 01 β^+ =100
³⁸ K ⁿ	-25342.7	0.4	3458.0	0.2		21.98 μ s	0.11	(7 ⁺), (5 ⁺) 01 IT=100
³⁸ Ca	-22059	5	440 ms	8	0 ⁺	01	β^+ =100	
³⁸ Sc	-4940#	300#	< 300 ns		2 ⁻ #	01	94B110 I	p ?
³⁸ Sc ^m	-4270#	320#	670#	100#		5 ⁻ #	01	IT ?; p ?
³⁸ Ti	9100#	250#	< 120 ns		0 ⁺	01	96B121 I	2p ?
* ³⁸ Mg	I : 18 events reported							**
³⁹ Mg	43570#	510#	< 260 ns		7/2 ⁻ #	02No11 I	n ?	*
³⁹ Al	21400	1470	10# ms(>200 ns)	3/2 ⁺ #	01	89Gu03 I	β^- ?	
³⁹ Si	1930	340	90# ms (>1 μ s)	7/2 ⁻ #	01	90Au.A I	β^- ?	
³⁹ P	-12870	100	190 ms	50	1/2 ⁺ #	01	95Re.A TD β^- =100; β^- -n=26.8	
³⁹ S	-23160	50	11.5 s	0.5	(3,5,7)/2 ⁻	01	β^- =100	
³⁹ Cl	-29800.2	1.7	55.6 m	0.2	3/2 ⁺	01	β^- =100	
³⁹ Ar	-33242	5	269 y	3	7/2 ⁻	01	β^- =100	
³⁹ K	-33807.01	0.19	STABLE		3/2 ⁺	01	IS=93.2581 44	
³⁹ Ca	-27274.4	1.9	859.6 ms	1.4	3/2 ⁺	01	β^+ =100	
³⁹ Sc	-14168	24	< 300 ns		7/2 ⁻ #	01	94B110 I	p=100
³⁹ Ti	1500#	210#	31 ms	4	3/2 ⁺ #	01	90De43 TD β^+ =100; ...	*
* ³⁹ Mg	T : estimated half-life 1# ms for β^- decay							**
* ³⁹ Sc	D : most probably proton emitter from $S_p=-602(24)$ keV							**
* ³⁹ Ti	D : ... ; β^+ p=85 15; β^+ 2p=15# D : β^+ 2p decay observed by 92Mo15							**
* ³⁹ Ti	T : average 90De43=26(+8-7) 01Gi01=31(+6-4)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)		
^{40}Mg	50240#	900#	1# ms	0^+		02No11 I	$\beta^- ?; \beta^- n ?$	*	
^{40}Al	29300#	700#	10# ms (>260 ns)		02	97Sa14 I	$\beta^- ?; \beta^- n ?$	*	
^{40}Si	5470	560	20# ms (>200 ns)	0^+	02	89Gu03 I	$\beta^- ?; \beta^- n ?$	*	
^{40}P	-8110	140	153 ms 8	$(2^-, 3^-)$	02		$\beta^- =100; \dots$	*	
^{40}S	-22870	140	8.8 s 2.2	0^+	02		$\beta^- =100$		
^{40}Cl	-27560	30	1.35 m 0.02	2^-	02		$\beta^- =100$		
^{40}Ar	-35039.8960	0.0027	STABLE	0^+	02		IS=99.6003 30		
^{40}K	-33535.20	0.19	1.251 Gy 0.011	4^-	02		IS=0.0117 1; ...	*	
$^{40}\text{K}^m$	-31891.56	0.19	1643.639 0.011	336 ns 12	0^+	02	IT=100		
^{40}Ca	-34846.27	0.21	STABLE (>5.9Zy)	0^+	01	99Be64 T	IS=96.941 156; $2\beta^+ ?$	*	
^{40}Sc	-20523.2	2.8	182.3 ms 0.7	4^-	02		$\beta^+ =100; \dots$	*	
^{40}Ti	-8850	160	53.3 ms 1.5	0^+	02		$\beta^+ =100; \beta^+ p=100$		
^{40}V	10330#	500#		$2^- \#$			p ?		
* ^{40}Mg	I : one event expected, none observed; similar search in 02Le.A							**	
* ^{40}Al	I : 34 events reported in 97Sa14; also one event in 96Sa34							**	
* ^{40}P	D : ... ; $\beta^- n=15.8 21$							**	
* ^{40}K	D : ... ; $\beta^- =89.28 13; \beta^+ =10.72 13$							**	
* ^{40}Sc	D : ... ; $\beta^+ p=0.44 7; \beta^+ \alpha=0.017 5$							**	
^{41}Al	35700#	800#	2# ms (>260 ns)	$3/2^+ \#$	02	97Sa14 I	$\beta^- ?$	*	
^{41}Si	13560	1840	30# ms (>200 ns)	$7/2^- \#$	02	89Gu03 I	$\beta^- ?$		
^{41}P	-5280	220	150 ms 15	$1/2^+ \#$	02		$\beta^- =100; \beta^- n=30 10$		
^{41}S	-19020	120	1.99 s 0.05	$7/2^- \#$	02		$\beta^- =100; \beta^- n ?$		
^{41}Cl	-27310	70	38.4 s 0.8	$(1/2, 3/2^+)$	02		$\beta^- =100$		
^{41}Ar	-33067.5	0.3	109.61 m 0.04	$7/2^-$	02		$\beta^- =100$		
^{41}K	-35559.07	0.19	STABLE	$3/2^+$	02		IS=6.7302 44		
^{41}Ca	-35137.76	0.24	102 ky 7	$7/2^-$	02		$\epsilon=100$		
^{41}Sc	-28642.39	0.23	596.3 ms 1.7	$7/2^-$	02		$\beta^+ =100$		
$^{41}\text{Sc}^r$	-25760.10	0.23	2882.30 0.05 RQ	$7/2^+$	02		P=59 2; IT=41 2		
^{41}Ti	-15700#	100#	80.9 ms 1.2	$3/2^+$	02	98Bh12 T	$\beta^+ =100; \beta^+ p \approx 100$	*	
^{41}V	-210#	210#		$7/2^- \#$			p ?		
* ^{41}Al	I : reported 4 events							**	
* ^{41}Ti	T : average 98Bh12=81.3(2.0) 98Li46=82(3) 96Fa09=81(4) 74Se11=80(2)							**	
^{42}Al	43680#	900#	1# ms				$\beta^- ?; \beta^- n ?$		
^{42}Si	18430#	500#	5# ms (>200 ns)	0^+	01	90Le03 I	$\beta^- ?; \beta^- n ?$	*	
^{42}P	940	450	120 ms 30		01	89Le16 T	$\beta^- =100; \beta^- n=50 20$		
^{42}S	-17680	120	1.013 s 0.015	0^+	01		$\beta^- =100; \beta^- n < 4$		
^{42}Cl	-24910	140	6.8 s 0.3		01		$\beta^- =100$		
^{42}Ar	-34423	6	32.9 y 1.1	0^+	01		$\beta^- =100$		
^{42}K	-35021.56	0.22	12.360 h 0.012	2^-	01		$\beta^- =100$		
^{42}Ca	-38547.07	0.25	STABLE	0^+	01		IS=0.647 23		
^{42}Sc	-32121.24	0.27	681.3 ms 0.7	0^+	01		$\beta^+ =100$		
$^{42}\text{Sc}^m$	-31504.96	0.28	616.28 0.06	61.7 s 0.4	$(7, 5, 6)^+$	01	$\beta^+ =100$		
$^{42}\text{Sc}^r$	-26044.91	0.26	6076.33 0.08 RQ	$(1^+ \text{ to } 4^+)$	01		IT=100		
^{42}Ti	-25122	5	199 ms 6	0^+	01		$\beta^+ =100$		
^{42}V	-8170#	200#	< 55 ns	$2^- \#$	01	92Bo37 I	p ?		
^{42}Cr	5990#	300#	14 ms 3	0^+	01	01Gi01 TD	$\beta^+ \approx 100; \beta^+ p=?; 2p ?$		
* ^{42}Si	TD : ENSDF reports preliminary values from 98Yo.A: half-life=20 ms 10 and							**	
* ^{42}Si	TD : $\beta^- n=103 48$, subject to further analysis according to the authors							**	
^{43}Si	26700#	700#	15# ms (>260 ns)	$3/2^- \#$		02No11 I	$\beta^- ?; \beta^- n ?$		
^{43}P	5770	970	33 ms 3	$1/2^+ \#$	01		$\beta^- =100; \beta^- n=100$		
^{43}S	-11970	200	260 ms 15	$3/2^- \#$	01	98Wi.A T	$\beta^- =100; \beta^- n=40 10$		
$^{43}\text{S}^m$	-11650	200	319 5	480 ns 50	$(7/2^-)$	01	00Sa21 EJ	IT=100	*
^{43}Cl	-24170	160	3.07 s 0.07	$3/2^+ \#$	01		$\beta^- =100; \beta^- n ?$		
^{43}Ar	-32010	5	5.37 m 0.06	$(5/2^-)$	01		$\beta^- =100$		
^{43}K	-36593	9	22.3 h 0.1	$3/2^+$	01		$\beta^- =100$		
^{43}Ca	-38408.6	0.3	STABLE	$7/2^-$	01		IS=0.135 10		

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...							
^{43}Sc	-36187.9	1.9	3.891 h	0.012		7/2 ⁻ 01	$\beta^+=100$
$^{43}\text{Sc}^m$	-36036.5	1.9 151.4 0.2	438 μs	7		3/2 ⁺ 01	IT=100
^{43}Ti	-29321	7	509 ms	5		7/2 ⁻ 01	$\beta^+=100$
$^{43}\text{Ti}^m$	-29008	7 313.0 1.0	12.6 μs	0.6		(3/2 ⁺) 01	IT=100
$^{43}\text{Ti}^n$	-26255	7 3066.4 1.0	560 ns	6		(19/2 ⁻) 01	IT=100
^{43}V	-18020#	230#	80# ms			7/2 ⁻ # 01	$\beta^+?$ *
^{43}Cr	-2130#	220#	21.6 ms	0.7		(3/2 ⁺) 01	$\beta^+=100; \beta^+p=23\ 6; \dots$ *
$^{43}\text{S}^m$	J : from comparison of B(E2) and half-life with theoretical ones **						
^{43}V	T : >800 ms reported by 92Bo37 and adopted in ENSDF'01. To be confirmed. **						
^{43}Cr	D : ... ; $\beta^+2p=6\ 5; \beta^+\alpha?$ **						
^{44}Si	32840#	800#	10# ms			0 ⁺	$\beta^-?$; $\beta^-n?$
^{44}P	12100#	700#	30# ms (>200 ns)			99 89Gu03 I	$\beta^-?$
^{44}S	-9120	390	123 ms	10		99	$\beta^-=100; \beta^-n=18\ 3$
^{44}Cl	-20230	110	560 ms	110		99	$\beta^-=100; \beta^-n<8$
^{44}Ar	-32673.1	1.6	11.87 m	0.05		0 ⁺ 99	$\beta^-=100$
^{44}K	-35810	40	22.13 m	0.19		2 ⁻ 99	$\beta^-=100$
^{44}Ca	-41468.5	0.4	STABLE			0 ⁺ 99	IS=2.086 110
^{44}Sc	-37816.1	1.8	3.97 h	0.04		2 ⁺ 99	$\beta^+=100$
$^{44}\text{Sc}^m$	-37545.2	1.8 270.95 0.20	58.61 h	0.10		6 ⁺ 99	IT=98.80 7; $\beta^+=1.20\ 7$
$^{44}\text{Sc}^n$	-37669.9	1.8 146.224 0.022	50.4 μs	0.7		0 ⁻ 99	
^{44}Ti	-37548.5	0.7	60.0 y	1.1		0 ⁺ 99	$\epsilon=100$ *
^{44}V	-24120	120	* 111 ms	7		(2 ⁺) 99	$\beta^+=100; \beta^+\alpha=?$
$^{44}\text{V}^m$	-23850#	160# 270# 100#	* 150 ms	3		(6 ⁺) 99	$\beta^+=100$
$^{44}\text{V}^n$	-23970#	160# 150# 100#				0 ⁻ #	
^{44}Cr	-13460#	50#	54 ms	4		0 ⁺ 99 96Fa09 D	$\beta^+=100; \beta^+p=7\ 3$
^{44}Mn	6400#	500#	< 105 ns			2 ⁻ # 99	p ?
^{44}Ti	T : also 01Ha21=59(2) **						
^{45}P	17900#	800#	8# ms (>200 ns)			1/2 ⁺ # 93 90Le03 I	$\beta^-?$
^{45}S	-3250	1740	82 ms	13		3/2 ⁻ # 97	$\beta^-=100; \beta^-n=54$
^{45}Cl	-18360	120	400 ms	40		3/2 ⁺ # 95	$\beta^-=100; \beta^-n=24\ 4$
^{45}Ar	-29770.6	0.5	21.48 s	0.15		(1,3,5)/2 ⁻ 95	$\beta^-=100$ *
^{45}K	-36608	10	17.3 m	0.6		3/2 ⁺ 95	$\beta^-=100$
^{45}Ca	-40812.0	0.4	162.67 d	0.25		7/2 ⁻ 95 94Lo04 T	$\beta^-=100$
^{45}Sc	-41067.8	0.8	STABLE			7/2 ⁻ 95	IS=100.
$^{45}\text{Sc}^m$	-41055.4	0.8 12.40 0.05	318 ms	7		3/2 ⁺ 95	IT=100
^{45}Ti	-39005.7	1.0	184.8 m	0.5		7/2 ⁻ 95	$\beta^+=100$
^{45}V	-31880	17	547 ms	6		7/2 ⁻ 95	$\beta^+=100$
^{45}Cr	-18970	500	* 50 ms	6		7/2 ⁻ # 95	$\beta^+=100; \beta^+p>27$
$^{45}\text{Cr}^m$	-18920#	510# 50# 100#	* 1# ms			3/2 ⁺ #	IT ?; $\beta^+?$
^{45}Mn	-5110#	300#	< 70 ns			7/2 ⁻ # 97 92Bo37 I	p ?
^{45}Fe	13580#	220#	4.9 ms	1.5		3/2 ⁺ # 97 02Gi09 TD	2p=75 5; $\beta^+=25\ 5; \dots$ *
^{45}Ar	J : 7/2 ⁻ # is expected from theory and from systematics. See ENSDF. **						
^{45}Fe	D : ... ; $\beta^+p=25\ 5$ **						
^{45}Fe	T : average 02Gi09=4.7(+3.4-1.4) 02Pf02=3.2(+2.6-1.0) D : β^+p from 01Gi01 **						
^{46}P	25500#	900#	4# ms (>200 ns)			00 90Le03 I	$\beta^-?$
^{46}S	700#	700#	30# ms (>200 ns)			0 ⁺ 00 89Gu03 I	$\beta^-?$
^{46}Cl	-14710	720	220 ms	40		00	$\beta^-=100; \beta^-n=60\ 9$
^{46}Ar	-29720	40	8.4 s	0.6		0 ⁺ 00	$\beta^-=100$
^{46}K	-35418	16	105 s	10		2 ⁽⁻⁾ 00 82To02 J	$\beta^-=100$
^{46}Ca	-43135.1	2.3	STABLE (>100 Ey)			0 ⁺ 00 99Be64 T	IS=0.004 3; 2 $\beta^-?$ *
^{46}Sc	-41757.1	0.8	83.79 d	0.04		4 ⁺ 00	$\beta^-=100$
$^{46}\text{Sc}^m$	-41614.6	0.8 142.528 0.007	18.75 s	0.04		1 ⁻ 00	IT=100
^{46}Ti	-44123.4	0.8	STABLE			0 ⁺ 00	IS=8.25 3
^{46}V	-37073.0	1.0	422.50 ms	0.11		0 ⁺ 00	$\beta^+=100$
$^{46}\text{V}^m$	-36271.5	1.0 801.46 0.10	1.02 ms	0.07		3 ⁺ 00	IT=100
... A-group is continued on next page ...							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
⁴⁶ Cr	-29474 20		260 ms	60	0 ⁺	00	$\beta^+=100$	
⁴⁶ Mn	-12370# 110#	*	37 ms	3	(4 ⁺)	00	$\beta^+=100; \beta^+p=22\ 2; \dots$ *	
⁴⁶ Mn ^m	-12220# 150# 150# 100#	*	1# ms		1 ⁻ #		$\beta^+?$	
⁴⁶ Fe	760# 350#		9 ms	4	0 ⁺	00	$\beta^+=100; \beta^+p=36\ 20$	
* ⁴⁶ Ca	T: limit is for neutrinoless $\beta\beta$ decay							**
* ⁴⁶ Mn	D: ...; $\beta^+2p\approx 18; \beta^+\alpha?$							**
* ⁴⁶ Mn	T: average 92Bo37=41(+7-6) 01Gi01=34.0(+4.5-3.5)							**
* ⁴⁶ Mn	D: $\beta^+2p\approx 18\%$ estimated from $P_p = \beta^+p + 2\times\beta^+2p=58(9)\%$ in 01Gi01							**
⁴⁷ S	8000# 800#		20# ms (>200 ns)		3/2 ⁻ #	95	$\beta^-?$	
⁴⁷ Cl	-10520# 600#		200# ms (>200 ns)		3/2 ⁺ #	95	$\beta^-=100; \beta^-n<3$	
⁴⁷ Ar	-25910 100		580 ms	120	3/2 ⁻ #	95	$\beta^-=100; \beta^-n<1$ *	
⁴⁷ K	-35696 8		17.50 s	0.24	1/2 ⁺	95	$\beta^-=100$	
⁴⁷ Ca	-42340.1 2.3		4.536 d	0.003	7/2 ⁻	95	$\beta^-=100$	
⁴⁷ Sc	-44332.1 2.0		3.3492 d	0.0006	7/2 ⁻	95	$\beta^-=100$	
⁴⁷ Sc ^m	-43565.3 2.0	766.83 0.09	272 ns	8	(3/2 ⁺)	95	IT=100	
⁴⁷ Ti	-44932.4 0.8		STABLE		5/2 ⁻	95	IS=7.44 2	
⁴⁷ V	-42002.1 0.8		32.6 m	0.3	3/2 ⁻	95	$\beta^+=100$	
⁴⁷ Cr	-34558 14		500 ms	15	3/2 ⁻	95	$\beta^+=100$	
⁴⁷ Mn	-22260# 160#		100 ms	50	5/2 ⁻ #	95	$\beta^+=100; \beta^+p=3.4\ 9$	
⁴⁷ Fe	-6620# 260#		21.8 ms	0.7	7/2 ⁻ #	97	$\beta^+=100; \beta^+p=87\ 7$	
⁴⁷ Fe ^m	-5850# 280#	770# 100#			3/2 ⁺ #		IT?	
⁴⁷ Co	10700# 500#				7/2 ⁻ #		p?	
* ⁴⁷ Ar	D: from 95So03							**
⁴⁸ S	13200# 900#		10# ms (>200 ns)		0 ⁺	90Le03 I	$\beta^-?$	
⁴⁸ Cl	-4700# 700#		100# ms (>200 ns)			89Gu03 I	$\beta^-?$	
⁴⁸ Ar	-23720# 300#		500# ms		0 ⁺		$\beta^-?$	
⁴⁸ K	-32124 24		6.8 s	0.2	(2 ⁻)	95	$\beta^-=100; \beta^-n=1.14\ 15$	
⁴⁸ Ca	-44214 4		53 Ey	17	0 ⁺	95	IS=0.187 21; ... *	
⁴⁸ Sc	-44496 5		43.67 h	0.09	6 ⁺	95	$\beta^-=100$	
⁴⁸ Ti	-48487.7 0.8		STABLE		0 ⁺	95	IS=73.72 3	
⁴⁸ V	-44475.4 2.6		15.9735 d	0.0025	4 ⁺	95	$\beta^+=100$	
⁴⁸ Cr	-42819 7		21.56 h	0.03	0 ⁺	95	$\beta^+=100$	
⁴⁸ Mn	-29320 110		158.1 ms	2.2	4 ⁺	97	$\beta^+=100; \beta^+p=0.28\ 4; \dots$ *	
⁴⁸ Fe	-18160# 70#		44 ms	7	0 ⁺	95	$\beta^+=100; \beta^+p=3.6\ 11$	
⁴⁸ Co	1640# 400#				6 ⁺ #		p?	
⁴⁸ Ni	18400# 500#		10# ms (>500 ns)		0 ⁺	01	00Bi01 I	
* ⁴⁸ Ca	D: ...; $2\beta^-=?; \beta^-?$							**
* ⁴⁸ Ca	T: average 00Br63=42(33-13) 96Ba80=43(+24-11 statistics + 14 systematics)							**
* ⁴⁸ Ca	T: also $T>36$ Ey from 70Ba61. Single β^- decay: $T>6$ Ey (95% CL), from 85Al17							**
* ⁴⁸ Mn	D: ...; $\beta^+\alpha=6e-4$							**
* ⁴⁸ Mn	D: one $\beta^+\alpha$ event was observed, versus 437 β^+p , in fig.4 of 87Se07							**
⁴⁹ S	22000# 950#		< 200 ns		3/2 ⁻ #	97	n?	
⁴⁹ Cl	300# 800#		50# ms (>200 ns)		3/2 ⁺ #	95	$\beta^-?$	
⁴⁹ Ar	-18150# 500#		170 ms	50	3/2 ⁻ #	95	$\beta^-=100; \beta^-n=65\ 20$	
⁴⁹ K	-30320 70		1.26 s	0.05	(3/2 ⁺)	95	$\beta^-=100; \beta^-n=86\ 9$	
⁴⁹ Ca	-41289 4		8.718 m	0.006	3/2 ⁻	95	$\beta^-=100$	
⁴⁹ Sc	-46552 4		57.2 m	0.2	7/2 ⁻	95	$\beta^-=100$	
⁴⁹ Ti	-48558.8 0.8		STABLE		7/2 ⁻	95	IS=5.41 2	
⁴⁹ V	-47956.9 1.2		330 d	15	7/2 ⁻	95	$\epsilon=100$	
⁴⁹ Cr	-45330.5 2.4		42.3 m	0.1	5/2 ⁻	95	$\beta^+=100$	
⁴⁹ Mn	-37616 24		382 ms	7	5/2 ⁻	01	$\beta^+=100$	
⁴⁹ Fe	-24580# 150#		70 ms	3	(7/2 ⁻)	01	$\beta^+=100; \beta^+p=52\ 10$	
⁴⁹ Co	-9580# 260#		< 35 ns		7/2 ⁻ #	97	p?	
⁴⁹ Ni	9000# 400#		13 ms	4	7/2 ⁻ #	97	$\beta^+=100; \beta^+p=?$	
* ⁴⁹ S	I: statistics precludes any conclusion, say authors							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
⁵⁰ Cl	7300# 900#		20# ms				β^- ?
⁵⁰ Ar	-14500# 700#		85 ms 30	0 ⁺	95	03We09 TD	β^- =100; β^- n=35 10
⁵⁰ K	-25350 280		472 ms 4	(0 ⁻ , 1, 2 ⁻)	95		β^- =100; β^- n=29 3
⁵⁰ Ca	-39571 9		13.9 s 0.6	0 ⁺	95		β^- =100
⁵⁰ Sc	-44537 16		102.5 s 0.5	5 ⁺	95		β^- =100
⁵⁰ Sc ^m	-44280 16	256.895 0.010	350 ms 40	2 ⁺ , 3 ⁺	95		IT>97.5; β^- <2.5
⁵⁰ Ti	-51426.7 0.8		STABLE	0 ⁺	95		IS=5.18 2
⁵⁰ V	-49221.6 1.0		150 Py 40	6 ⁺	95		IS=0.250 4; β^+ =83 11; ... *
⁵⁰ Cr	-50259.5 1.0		STABLE (>1.3 Ey)	0 ⁺	95	03Bi05 T	IS=4.345 13; 2 β^+ ?
⁵⁰ Mn	-42626.8 1.0		283.9 ms 0.5	0 ⁺	95		β^+ =100
⁵⁰ Mn ^m	-42398 7	229 7	1.75 m 0.03	5 ⁺	95		β^+ =100
⁵⁰ Fe	-34480 60		155 ms 11	0 ⁺	01		β^+ =100; β^+ p≈0
⁵⁰ Co	-17200# 170#		44 ms 4	(6 ⁺)	01	96Fa09 JD	β^+ =100; β^+ p=54 12
⁵⁰ Ni	-3790# 260#		9.1 ms 1.8	0 ⁺	97	01Ma.A T	β^+ ?
* ⁵⁰ V	D : ... ; β^- =17 11						**
⁵¹ Cl	13500# 1000#		2# ms (>200 ns)	3/2 ⁺ #	97	90Le03 I	β^- ?
⁵¹ Ar	-7800# 700#		60# ms (>200 ns)	3/2 ⁻ #	97	89Gu03 I	β^- ?
⁵¹ K	-22000# 500#		365 ms 5	3/2 ⁺ #	97		β^- =100; β^- n=47 5
⁵¹ Ca	-35860 90		10.0 s 0.8	3/2 ⁻ #	97		β^- =100; β^- n ?
⁵¹ Sc	-43218 20		12.4 s 0.1	(7/2) ⁻	97		β^- =100
⁵¹ Ti	-49727.8 1.0		5.76 m 0.01	3/2 ⁻	97		β^- =100
⁵¹ V	-52201.4 1.0		STABLE	7/2 ⁻	97		IS=99.750 4
⁵¹ Cr	-51448.8 1.0		27.7025 d 0.0024	7/2 ⁻	97		ϵ =100
⁵¹ Mn	-48241.3 1.0		46.2 m 0.1	5/2 ⁻	97		β^+ =100
⁵¹ Fe	-40222 15		305 ms 5	5/2 ⁻	97		β^+ =100
⁵¹ Co	-27270# 150#		60# ms (>200 ns)	7/2 ⁻ #	97	87Po04 I	β^+ ?
⁵¹ Ni	-11440# 260#		30# ms (>200 ns)	7/2 ⁻ #	97	87Po04 I	β^+ ?
⁵² Ar	-3000# 900#		10# ms	0 ⁺	00		β^- ?
⁵² K	-16200# 700#		105 ms 5	2 ⁻ #	00	ABBW D	β^- =100; β^- n≈64; ... *
⁵² Ca	-32510 700		4.6 s 0.3	0 ⁺	00		β^- =100; β^- n<2
⁵² Sc	-40360 190		8.2 s 0.2	3 ⁽⁺⁾	00		β^- =100
⁵² Ti	-49465 7		1.7 m 0.1	0 ⁺	00		β^- =100
⁵² V	-51441.3 1.0		3.743 m 0.005	3 ⁺	00		β^- =100
⁵² Cr	-55416.9 0.8		STABLE	0 ⁺	00		IS=83.789 18
⁵² Mn	-50705.4 2.0		5.591 d 0.003	6 ⁺	00		β^+ =100
⁵² Mn ^m	-50327.7 2.0	377.749 0.005	21.1 m 0.2	2 ⁺	00		β^+ =98.25 5; IT=1.75 5
⁵² Fe	-48332 7		8.275 h 0.008	0 ⁺	00		β^+ =100
⁵² Fe ^m	-41520 130	6810 130 BD	45.9 s 0.6	12 ⁺ #	00		β^+ ≈100; IT<0.004
⁵² Co	-33920# 70#		115 ms 23	(6 ⁺)	00		β^+ =100
⁵² Co ^m	-33540# 120#	380# 100#	104 ms 11	2 ⁺ #		97Ha04 TD	β^+ =?; IT ? *
⁵² Ni	-22650# 80#		38 ms 5	0 ⁺	00		β^+ =100; β^+ p=17.0 14
⁵² Cu	-2630# 260#			3 ⁺ #	00		p ?
* ⁵² K	D : ... ; β^- 2n≈21						**
* ⁵² K	D : β^- n≈64%, β^- 2n≈21% estimated from $P_n = \beta^-n + 2 \times \beta^-2n = 107(20)\%$ in ⁸³ La23						**
* ⁵² K	D : and assuming $\beta^-n/\beta^-2n=3$ as in ³² Na						**
* ⁵² Co ^m	I : tentative: no specific evidence for ⁵² Co ^m , say authors in 97Ha04						**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)			
⁵³ Ar	4600#	1000#	3# ms	5/2 ⁻ #	99		β^- ?; β^-n ?			
⁵³ K	-12000#	700#	30 ms	5	3/2 ⁺ #	99 ABBW D	β^- =100; β^-n ≈67; ... *			
⁵³ Ca	-27900#	500#	90 ms	15	3/2 ⁻ #	99 83La23 D	β^- =100; β^-n >30 *			
⁵³ Sc	-37620#	300#	> 3 s		7/2 ⁻ #	99 98So03 TD	β^- =100; β^-n ?			
⁵³ Ti	-46830	100	32.7 s	0.9	(3/2) ⁻	99	β^- =100			
⁵³ V	-51849	3	1.60 m	0.04	7/2 ⁻	99	β^- =100			
⁵³ Cr	-55284.7	0.8	STABLE		3/2 ⁻	99	IS=9.501 17			
⁵³ Mn	-54687.9	0.8	3.7 My	0.4	7/2 ⁻	99	ϵ =100			
⁵³ Fe	-50945.3	1.8	8.51 m	0.02	7/2 ⁻	99	β^+ =100			
⁵³ Fe ^m	-47904.9	1.8	3040.4	0.3	2.526 m	0.024	19/2 ⁻	99 97Ge11 T	IT=100 *	
⁵³ Co	-42645	18	242 ms	8	7/2 ⁻ #	99	02Lo13 T	β^+ =100 *		
⁵³ Co ^m	-39447	22	3197	29	p	247 ms	12	(19/2 ⁻)	99	β^+ ≈98.5; p≈1.5
⁵³ Ni	-29370#	160#	45 ms	15	7/2 ⁻ #	99	76Vi02 D	β^+ =100; β^+p ≈45		
⁵³ Cu	-13460#	260#	< 300 ns		3/2 ⁻ #	99	93Bl.A I	p ?; β^+ ?		
* ⁵³ K	D : ... ; β^-2n ≈17							**		
* ⁵³ K	D : β^-n ≈67%, β^-2n ≈17% estimated from $P_n = \beta^-n + 2 \times \beta^-2n=100(30)\%$ in 83La23							**		
* ⁵³ K	D : and assuming $\beta^-n/\beta^-2n=4$ as in ³³ Na							**		
* ⁵³ Ca	D : $\beta^-n=40(10)\%$ is a lower limit (see ENSDF)							**		
* ⁵³ Ca	T : expected $T=2\#$ s from systematics of Ca isotopes							**		
* ⁵³ Fe ^m	T : average 97Ge11=2.48(0.05) 68De27=2.51(0.02) 67Es06=2.58(0.03)							**		
* ⁵³ Co	T : average 02Lo13=240(9) 89Ho13=240(20) 73Ko10=262(25)							**		
⁵⁴ K	-5400#	900#	10 ms	5	2 ⁻ #	01	β^- =100; β^-n =?			
⁵⁴ Ca	-23890#	700#	50# ms	(>300 ns)	0 ⁺	01	97Be70 I	β^- ?; β^-n ?		
⁵⁴ Sc	-34220	370	260 ms	30	3 ⁺ #	01	02Ja16 T	β^- =100; β^-n ? *		
⁵⁴ Sc ^m	-34110	370	110	3	7 μ s	5	(5 ⁺)	01 98Gr14 EJ	IT=100	
⁵⁴ Ti	-45590	120	1.5 s	0.4	0 ⁺	01	β^- =100			
⁵⁴ V	-49891	15	49.8 s	0.5	3 ⁺	01	β^- =100			
⁵⁴ V ^m	-49783	15	108	3	900 ns	500	(5 ⁺)	98Gr14 EJ	IT=100	
⁵⁴ Cr	-56932.5	0.8	STABLE		0 ⁺	01	IS=2.365 7			
⁵⁴ Mn	-55555.4	1.3	312.03 d	0.03	3 ⁺	01	02Un02 T	ϵ =100; β^- <2.9e-4; ... *		
⁵⁴ Fe	-56252.5	0.7	STABLE		0 ⁺	01	IS=5.845 35; 2 β^+ ?			
⁵⁴ Fe ^m	-49725.6	0.9	6526.9	0.6	364 ns	7	10 ⁺	01	IT=100	
⁵⁴ Co	-48009.5	0.7	193.23 ms	0.14	0 ⁺	01	β^+ =100			
⁵⁴ Co ^m	-47812.1	0.9	197.4	0.5	1.48 m	0.02	(7) ⁺	01	β^+ =100	
⁵⁴ Ni	-39210	50	104 ms	7	0 ⁺	01	02Lo13 T	β^+ =100 *		
⁵⁴ Cu	-21690#	210#	< 75 ns		3 ⁺ #	01	94Bl10 I	p ?		
⁵⁴ Zn	-6570#	400#			0 ⁺			2p ?		
* ⁵⁴ Sc	T : average 02Ja16=360(60) 98So03=225(40)							**		
* ⁵⁴ Mn	D : ... ; e^+ =1.28e-7 25							**		
* ⁵⁴ Mn	D : e^+ average 98Wu01=1.20(0.26) 97Za07=2.2(0.9)							**		
* ⁵⁴ Ni	T : average 02Lo13=103(9) 99Re06=106(12)							**		
⁵⁵ K	-270#	1000#	3# ms		3/2 ⁺ #		β^- ?; β^-n ?			
⁵⁵ Ca	-18120#	700#	30# ms	(>300 ns)	5/2 ⁻ #		97Be70 I	β^- ?		
⁵⁵ Sc	-29580	740	120 ms	40	7/2 ⁻ #	01	β^- =100; β^-n ?			
⁵⁵ Ti	-41670	150	490 ms	90	3/2 ⁻ #	01	98Am04 T	β^- =100 *		
⁵⁵ V	-49150	100	6.54 s	0.15	7/2 ⁻ #	01	β^- =100			
⁵⁵ Cr	-55107.5	0.8	3.497 m	0.003	3/2 ⁻	01	β^- =100			
⁵⁵ Mn	-57710.6	0.7	STABLE		5/2 ⁻	01	IS=100.			
⁵⁵ Fe	-57479.4	0.7	2.737 y	0.011	3/2 ⁻	01	ϵ =100			
⁵⁵ Co	-54027.6	0.7	17.53 h	0.03	7/2 ⁻	01	β^+ =100			
⁵⁵ Ni	-45336	11	204.7 ms	1.7	7/2 ⁻	01	02Lo13 T	β^+ =100 *		
⁵⁵ Cu	-31620#	300#	40# ms	(>200 ns)	3/2 ⁻ #	01	87Po04 I	β^+ ?; p ?		
⁵⁵ Zn	-14920#	250#	20# ms	(>1.6 μ s)	5/2 ⁻ #	01	01Gi10 I	β^+ ?		
* ⁵⁵ Ti	T : unweighed average 98Am04=320(60) 96Do23=600(40) and 95So.A=545(95)							**		
* ⁵⁵ Ti	T : (Birge ratio $B=2.75$)							**		
* ⁵⁵ Ni	T : average 02Lo13=196(5) 99Re06=204(3) 87Ha.A=212.1(3.8) 84Ay01=208(5)							**		
* ⁵⁵ Ni	T : and 77Ho25=189(5) 76Ed.A=219(6); 97Wo06=204(3) superseded by 99Re06							**		

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
⁵⁶ Ca	-13440#	900#	10# ms (>300 ns)	0 ⁺	99	97Be70 I	β^- ?	
⁵⁶ Sc	-25270#	700#	80# ms (>300 ns)	3 ⁺ #	99	97Be70 I	β^- ?	
⁵⁶ Ti	-38940	200	164 ms	24	0 ⁺	99 98Am04 TD	β^- =100; β^- n ? *	
⁵⁶ V	-46080	200	216 ms	4	(1 ⁺)	99 03Ma02 TJ	β^- =100; β^- n ?	
⁵⁶ Cr	-55281.2	1.9	5.94 m	0.10	0 ⁺	99	β^- =100	
⁵⁶ Mn	-56909.7	0.7	2.5789 h	0.0001	3 ⁺	99	β^- =100	
⁵⁶ Fe	-60605.4	0.7	STABLE		0 ⁺	99	IS=91.754 36	
⁵⁶ Co	-56039.4	2.1	77.23 d	0.03	4 ⁺	99	β^+ =100	
⁵⁶ Ni	-53904	11	6.075 d	0.010	0 ⁺	99	β^+ =100	
⁵⁶ Cu	-38600#	140#	93 ms	3	(4 ⁺)	99 01Bo54 TJD	β^+ =100; β^+ p=0.40 12	
⁵⁶ Zn	-25730#	260#	36 ms	10	0 ⁺	01 95Wa.A T	β^+ ?; β^+ p ? *	
⁵⁶ Ga	-4740#	260#			3 ⁺ #		p ?	
* ⁵⁶ Ti	T: average 98Am04=190(40) 96Do23=150(30)							**
* ⁵⁶ Zn	T: half-life is derived from experimental (p,n) cross sections							**
* ⁵⁶ Zn	I: identified by time-of-flight 01Gi10 with $T > 1.6 \mu$ s							**
⁵⁷ Ca	-7120#	1000#	5# ms		5/2 ⁻ #		β^- ?; β^- n ?	
⁵⁷ Sc	-20690#	700#	13 ms	4	7/2 ⁻ #	98 02So.A TD	β^- =100; β^- n=33#	
⁵⁷ Ti	-33540	460	60 ms	16	5/2 ⁻ #	98 99So20 T	β^- =100; β^- n=0.3# *	
⁵⁷ V	-44190	230	350 ms	10	(3/2 ⁻)	98 03Ma02 TJ	β^- =100; β^- n=0.4#	
⁵⁷ Cr	-52524.1	1.9	21.1 s	1.0	(3/2 ⁻)	98	β^- =100	
⁵⁷ Mn	-57486.8	1.8	85.4 s	1.8	5/2 ⁻	98	β^- =100	
⁵⁷ Fe	-60180.1	0.7	STABLE		1/2 ⁻	98	IS=2.119 10	
⁵⁷ Co	-59344.2	0.7	271.74 d	0.06	7/2 ⁻	98	ϵ =100	
⁵⁷ Ni	-56082.0	1.8	35.60 h	0.06	3/2 ⁻	98	β^+ =100	
⁵⁷ Cu	-47310	16	196.3 ms	0.7	3/2 ⁻	98	β^+ =100	
⁵⁷ Zn	-32800#	100#	38 ms	4	7/2 ⁻ #	98 02Lo13 T	β^+ =100; β^+ p \approx 65 *	
⁵⁷ Ga	-15900#	260#			1/2 ⁻ #		p ?	
* ⁵⁷ Ti	T: average 99So20=67(25) 96Do23=56(20); 98Am04=180(30) at variance not used							**
* ⁵⁷ Zn	T: average 02Lo13=37(5) 76Vi02=40(10)							**
⁵⁸ Sc	-15170#	800#	12 ms	5	3 ⁺ #	02So.A TD	β^- =100	
⁵⁸ Ti	-30770#	700#	54 ms	7	0 ⁺	97 99So20 TD	β^- =100	
⁵⁸ V	-40210	250	191 ms	8	3 ⁺ #	97 03Ma02 TD	β^- =100; β^- n ? *	
⁵⁸ Cr	-51830	200	7.0 s	0.3	0 ⁺	97	β^- =100	
⁵⁸ Mn	-55910	30	3.0 s	0.1	1 ⁺	97	β^- =100	
⁵⁸ Mn ^m	-55840	30	71.78	0.05	65.2 s	0.5	(4 ⁺) 97	β^- =?; IT=20#
⁵⁸ Fe	-62153.4	0.7	STABLE		0 ⁺	97	IS=0.282 4	
⁵⁸ Co	-59845.9	1.2	70.86 d	0.06	2 ⁺	00	β^+ =100	
⁵⁸ Co ^m	-59821.0	1.2	24.95	0.06	9.04 h	0.11	5 ⁺ 00	IT=100
⁵⁸ Co ⁿ	-59792.8	1.2	53.15	0.07	10.4 μ s	0.3	4 ⁺ 00	IT=100
⁵⁸ Ni	-60227.7	0.6	STABLE		(>700 Ey)	0 ⁺	01	IS=68.0769 89; 2 β^+ ? *
⁵⁸ Cu	-51662.1	1.6	3.204 s	0.007	1 ⁺	01	β^+ =100	
⁵⁸ Zn	-42300	50	84 ms	9	0 ⁺	99 02Lo13 T	β^+ =100; β^+ p<3 *	
⁵⁸ Ga	-23990#	210#			2 ⁺ #		p ?	
⁵⁸ Ga ^m	-23960#	230#	30#	100#	*		5 ⁺ #	
⁵⁸ Ge	-8370#	320#			0 ⁺		2p ?	
* ⁵⁸ Ti	T: average 02So.A=59(9) 99So20=47(10)							**
* ⁵⁸ V	T: average 03Ma02=185(10) 98Am04=200(20) 98So03=205(20)							**
* ⁵⁸ Ni	T: >400 Ey to 2 ⁺ level of ⁵⁸ Fe, >700 Ey to ground-state							**
* ⁵⁸ Zn	T: average 02Lo13=83(10) 98Jo18=86(18)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
⁵⁹ Sc	-10040# 900#		10# ms				β^- ?; β^-n ?	
⁵⁹ Ti	-25220# 700#		30 ms	3		02 02So.A T	β^- =100 *	
⁵⁹ V	-37070 310		75 ms	7		02	β^- =100; β^-n ?	
⁵⁹ Cr	-47890 240		460 ms	50		02	β^- =100	
⁵⁹ Cr ^m	-47390 240	503.0 1.7	96 μ s	20		02 (9/2 ⁺)	IT=100	
⁵⁹ Mn	-55480 30		4.59 s	0.05		02 (5/2) ⁻	β^- =100	
⁵⁹ Fe	-60663.1 0.7		44.495 d	0.009		02 3/2 ⁻	β^- =100	
⁵⁹ Co	-62228.4 0.6		STABLE			02 7/2 ⁻	IS=100.	
⁵⁹ Ni	-61155.7 0.6		101 ky	13		02 94Ru19 T	β^+ =100 *	
⁵⁹ Cu	-56357.2 0.8		81.5 s	0.5		02 3/2 ⁻	β^+ =100	
⁵⁹ Zn	-47260 40		182.0 ms	1.8		02 3/2 ⁻	β^+ =100; β^+p =0.10 3	
⁵⁹ Ga	-34120# 170#						β^+ ?	
⁵⁹ Ge	-17000# 280#						2p ?	
* ⁵⁹ Ti	T : supersedes 99So20=58(17) same group							**
* ⁵⁹ Ni	T : unweighed average 94Ru19=108(13) 94Ru19(meteorite)=120(22) 81Ni08=76(5)							**
* ⁵⁹ Ni	T : (Birge ratio B=2.05)							**
⁶⁰ Sc	-4000# 900#		3# ms				β^- ?	
⁶⁰ Ti	-21650# 800#		22 ms	2		02So.A TD	β^- =100	
⁶⁰ V	-32580 470		122 ms	18		97 99So20 TD	β^- =100; β^-n ? *	
⁶⁰ V ^m	-32580# 490#	0# 150#	40 ms	15		03So02 TD	β^- =?; IT ?	
⁶⁰ V ⁿ	-32480 470	101 1				99So20 EI	IT=100	
⁶⁰ Cr	-46500 210		560 ms	60		93 96Do23 T	β^- =100 *	
⁶⁰ Mn	-53180 90		51 s	6		94	β^- =100	
⁶⁰ Mn ^m	-52910 90	271.90 0.10	1.77 s	0.02		94 92Sc.A E	β^- =88.5 8; IT=11.5 8	
⁶⁰ Fe	-61412 3		1.5 My	0.3		93	β^- =100	
⁶⁰ Co	-61649.0 0.6		5.2713 y	0.0008		00 5 ⁺	β^- =100	
⁶⁰ Co ^m	-61590.4 0.6	58.59 0.01	10.467 m	0.006		00 2 ⁺	IT \approx 100; β^- =0.24 3	
⁶⁰ Ni	-64472.1 0.6		STABLE			96	IS=26.2231 77	
⁶⁰ Cu	-58344.1 1.7		23.7 m	0.4		93 2 ⁺	β^+ =100	
⁶⁰ Zn	-54188 11		2.38 m	0.05		02 0 ⁺	β^+ =100	
⁶⁰ Ga	-40000# 110#		70 ms	10		02 01Ma96 TJ	β^+ =100; β^+p =1.6 7; ... *	
⁶⁰ Ge	-27770# 230#		30# ms				β^+ ?	
⁶⁰ As	-6400# 600#						p ?	
⁶⁰ As ^m	-6340# 600#	60# 20#					p ?	
* ⁶⁰ V	T : also 98Am04=200(40), not used							**
* ⁶⁰ Cr	T : weighed average 96Do23=510(150) 88Bo06=570(60); other 95Am.A=380(30)							**
* ⁶⁰ Ga	D : ... ; $\beta^+ \alpha < 0.023$ 20							**
* ⁶⁰ Ga	T : average 02Lo13=70(13) 01Ma96=70(15)							**
⁶¹ Ti	-15650# 900#		10# ms	(>300 ns)		99 97Be70 I	β^- ?; β^-n ?	
⁶¹ V	-29360# 400#		47.0 ms	1.2		99 03So02 TD	β^- =100; $\beta^-n < 6$	
⁶¹ Cr	-42180 250		261 ms	15		99 99So20 TD	β^- =100; β^-n ? *	
⁶¹ Mn	-51560 230		670 ms	40		99 99Ha05 D	β^- =100; β^-n =?	
⁶¹ Fe	-58921 20		5.98 m	0.06		99 3/2 ⁻ , 5/2 ⁻	β^- =100	
⁶¹ Fe ^m	-58060 20	861 3	250 ns	10		99 98Gr14 E	IT=100	
⁶¹ Co	-62898.4 0.9		1.650 h	0.005		99 7/2 ⁻	β^- =100	
⁶¹ Ni	-64220.9 0.6		STABLE			99 3/2 ⁻	IS=1.1399 6	
⁶¹ Cu	-61983.6 1.0		3.333 h	0.005		99 3/2 ⁻	β^+ =100	
⁶¹ Zn	-56345 16		89.1 s	0.2		99 3/2 ⁻	β^+ =100	
⁶¹ Zn ^m	-56257 16	88.4 0.1	< 430 ms			99 1/2 ⁻	IT=100	
⁶¹ Zn ⁿ	-55927 16	418.10 0.15	140 ms	70		99 3/2 ⁻	IT=100	
⁶¹ Zn ^p	-55589 16	756.02 0.18	< 130 ms			99 5/2 ⁻	IT=100	
⁶¹ Ga	-47090 50		168 ms	3		99 3/2 ⁻	β^+ =100; $\beta^+p \approx 0$	
⁶¹ Ga ^m	-47000# 110#	90# 100#					1/2 ⁻ #	
⁶¹ Ge	-33730# 300#		39 ms	12		99 02Lo13 T	β^+ =100; $\beta^+p \approx 80$ *	
⁶¹ As	-18050# 600#						p ?	
* ⁶¹ Cr	T : average 99So20=251(22) 98Am04=270(20)							**
* ⁶¹ Ge	T : average 02Lo13=36(21) 87Ho01=40(15)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{62}Ti	-11650#	900#	10# ms	0^+			β^- ?	
^{62}V	-24420#	500#	33.5 ms	2.0	$3^+\#$	01 03So02 TD	β^- =100	
^{62}Cr	-40410	340	199 ms	9	0^+	01 02So.A TD	β^- =100; β^-n ? *	
^{62}Mn	-48040	220	671 ms	5	(3^+)	01 99Ha05 TD	β^- =100; $\beta^-n=?$ *	
$^{62}\text{Mn}^m$	-48040#	270#	0# 150#	92 ms	13	(1 ⁺)	99So20 TJD	β^- =100; $\beta^-n\approx 0$
^{62}Fe	-58901	14	68 s	2	0^+	01	β^- =100	
^{62}Co	-61432	20	1.50 m	0.04	2^+	01	β^- =100	
$^{62}\text{Co}^m$	-61410	21	22 5	13.91 m	0.05	5^+	01	β^- >99; IT<1
^{62}Ni	-66746.1	0.6	STABLE		0^+	01	IS=3.6345 17	
^{62}Cu	-62798	4	9.673 m	0.008	1^+	01 02Un02 T	β^+ =100 *	
^{62}Zn	-61171	10	9.186 h	0.013	0^+	01	β^+ =100	
^{62}Ga	-52000	28	115.99 ms	0.17	0^+	01 03Hy02 T	β^+ =100 *	
$^{62}\text{Ga}^m$	-51183	28	817.5 0.5	4.6 ns	0.5	(3 ⁺)	01 98Vi06 ETJ	IT=100
^{62}Ge	-42240#	140#	130 ms	40	0^+	01 02Lo13 TD	β^+ =100 *	
^{62}As	-24960#	300#			$1^+\#$	01	p ? *	
* ^{62}Cr	T : average 02So.A=209(12) 99So20=187(15) 98Am04=190(30)							**
* ^{62}Cu	T : others 97Zi06(LS method)=9.68(0.04) 97Zi06(IC method)=9.673(0.026)							**
* ^{62}Cu	T : 69Jo07=9.73(0.02) 69Bo11=9.7(0.1) 65Li11=9.79(0.06) 65Eb01=9.76(0.02)							**
* ^{62}Ga	T : average 03Hy02=115.84(0.25) 79Da04=116.34(0.35) 78Al23=115.95(0.30)							**
* ^{62}Ge	I : T=113(+6-5) ms in 93Wi03 (table 1) is a misprint for ^{62}Ga							**
* ^{62}As	D : p-unstable from estimated $S_p=-1476\#(422\#)$ keV							**
^{63}Ti	-5200#	1000#	3# ms	$1/2^-$			β^- ?; β^-n ?	
^{63}V	-20910#	600#	17 ms	3	$7/2^-$	01 03So02 TD	β^- =100; $\beta^-n<35$	
^{63}Cr	-35530#	300#	129 ms	2	$1/2^-$	01 02So.A TD	β^- =100; β^-n ? *	
^{63}Mn	-46350	260	275 ms	4	$5/2^-$	01 99Ha05 TD	β^- =100; $\beta^-n=?$ *	
^{63}Fe	-55550	170	6.1 s	0.6	$(5/2)^-$	01	β^- =100	
^{63}Co	-61840	20	26.9 s	0.4	$7/2^-$	01 94It.A T	β^- =100 *	
^{63}Ni	-65512.6	0.6	100.1 y	2.0	$1/2^-$	01	β^- =100	
$^{63}\text{Ni}^m$	-65425.5	0.6	87.15 0.11	1.67 μs	0.03	$5/2^-$	01	IT=100
^{63}Cu	-65579.5	0.6	STABLE		$3/2^-$	01	IS=69.17 3	
^{63}Zn	-62213.0	1.6	38.47 m	0.05	$3/2^-$	01	β^+ =100	
^{63}Ga	-56547.1	1.3	32.4 s	0.5	$(3/2)^-$	01	β^+ =100	
^{63}Ge	-46910#	200#	142 ms	8	$3/2^-$	01 02Lo13 TD	β^+ =100 *	
^{63}As	-33820#	500#			$3/2^-$	01	p ? *	
* ^{63}Cr	T : also 99So20=113(16) and 98Am04=110(70) outweighed, not used							**
* ^{63}Mn	T : also 99So20=322(23) 95Am.A=290(20) 85Bo49=250(40) outweighed, not used							**
* ^{63}Co	T : average 94It.A=26.41(0.27) 72Jo08=27.5(0.3) 69Wa15=26(1)							**
* ^{63}Ge	T : average 02Lo13=150(9) 93Wi03=95(+23-20)							**
* ^{63}As	D : p-unstable from estimated $S_p=-1132\#(522\#)$ keV							**
^{64}V	-15400#	700#	10# ms (>300 ns)			97 97Be70 I	β^- ?	
^{64}Cr	-33150#	400#	43 ms	1	0^+	02So.A TD	β^- =100 *	
^{64}Mn	-42620	270	88.8 ms	2.5	(1^+)	96 99So20 TJD	β^- =100; $\beta^-n=?$ *	
$^{64}\text{Mn}^m$	-42490	270	135 3	> 100 μs		98Gr14 ET	IT=100	
^{64}Fe	-54770	280	2.0 s	0.2	0^+	96	β^- =100	
^{64}Co	-59793	20	300 ms	30	1^+	96	β^- =100	
^{64}Ni	-67099.3	0.6	STABLE		0^+	96	IS=0.9256 9	
^{64}Cu	-65424.2	0.6	12.700 h	0.002	1^+	96	β^+ =61.0 3; β^- =39.0 3	
^{64}Zn	-66003.6	0.7	STABLE	(>2.3 Ey)	0^+	96 85No03 T	IS=48.63 60; $2\beta^+$?	
^{64}Ga	-58834.3	2.0	2.627 m	0.012	$0^{(+\#)}$	96	β^+ =100	
$^{64}\text{Ga}^m$	-58791.5	2.0	42.85 0.08	21.9 μs	0.7	2^+	96 99Ta29 TJ	IT=100
^{64}Ge	-54350	30	63.7 s	2.5	0^+	96	β^+ =100	
^{64}As	-39520#	360#	40 ms	30	$0^+\#$	02Lo13 TD	β^+ =100	
* ^{64}Cr	T : also 99So20=44(12) outweighed, not used							**
* ^{64}Mn	T : average 02So.A=91(4) 99So20=85(5) 99Ha05=89(4); 98Am04=140(30) not used							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{65}V	-11250# 800#		10# ms			5/2 ⁻ #	β^- ?; β^-n ?
^{65}Cr	-27800# 500#		27 ms	3		1/2 ⁻ # 97	β^- =100; β^-n ?
^{65}Mn	-40670 540		92 ms	1		5/2 ⁻ # 93	β^- =100; β^-n ? *
^{65}Fe	-50880 240		1.3 s	0.3		1/2 ⁻ # 93	β^- =100 *
$^{65}\text{Fe}^m$	-50520 240	364 3	430 ns	130		(5/2 ⁻) 98Gr14	IT=100
^{65}Co	-59170 13		1.20 s	0.06		(7/2 ⁻) 93	β^- =100
^{65}Ni	-65126.1 0.6		2.5172 h	0.0003		5/2 ⁻ 97	β^- =100
$^{65}\text{Ni}^m$	-64113.1 1.2	1013 1	26.7 ns	1.0		9/2 ⁺ 95Bi01	ETJ
^{65}Cu	-67263.7 0.7		STABLE			3/2 ⁻ 93	IS=30.83 3
^{65}Zn	-65911.6 0.7		244.06 d	0.10		5/2 ⁻ 00	β^+ =100
$^{65}\text{Zn}^m$	-65857.7 0.7	53.928 0.010	1.6 μs	0.6		(1/2 ⁻) 00	IT=100
^{65}Ga	-62657.2 0.8		15.2 m	0.2		3/2 ⁻ 93	β^+ =100
^{65}Ge	-56410 100		30.9 s	0.5		(3/2 ⁻) 93	β^+ =100; β^+p =0.011 3
^{65}As	-46980# 300#		170 ms	30		3/2 ⁻ # 93	β^+ =100 *
^{65}Se	-32920# 600#		< 50 ms			3/2 ⁻ # 93	β^+ =100; β^+p ? *
* ^{65}Mn	T : others 99Ha05=88(4) 99So20=100(8) 98Am04=110(20) outweighed, not used **						
* ^{65}Fe	T : 95Am.A=760(50) ms supersedes 94Cz02=450(150) from same group, none used **						
* ^{65}As	T : average 02Lo13=126(16) 95Mo26=190(11) with Birge ratio $B=3.3$ **						
* ^{65}Se	D : from 93Ba12 **						
^{66}Cr	-24800# 600#		10 ms	6	0 ⁺	98 02So.A	TD β^- =100
^{66}Mn	-36250# 400#		64.4 ms	1.8		98 02So.A	TD β^- =100; β^-n ? *
^{66}Fe	-49570 300		440 ms	40	0 ⁺	98 99So20	TD β^- =100; β^-n ? *
^{66}Co	-56110 250		194 ms	17	(3 ⁺)	98 00Mu10	TJ β^- =100 *
$^{66}\text{Co}^m$	-55940 250	175 3	1.21 μs	0.01		(5 ⁺) 98Gr14	ETJ IT=100
$^{66}\text{Co}^n$	-55470 250	642 5	> 100 μs			(8 ⁻) 98Gr14	ETJ IT=100
^{66}Ni	-66006.3 1.4		54.6 h	0.4	0 ⁺	98	β^- =100
^{66}Cu	-66258.3 0.7		5.120 m	0.014	1 ⁺	98	β^- =100
^{66}Zn	-68899.4 0.9		STABLE		0 ⁺	98	IS=27.90 27
^{66}Ga	-63724 3		9.49 h	0.07	0 ⁺	98	β^+ =100
^{66}Ge	-61620 30		2.26 h	0.05	0 ⁺	98	β^+ =100
^{66}As	-51500 680		95.77 ms	0.23	(0 ⁺)	98 98Gr12	J β^+ =100
$^{66}\text{As}^m$	-50140 680	1356.70 0.17	1.1 μs	0.1	(5 ⁺)	01Gr07	TJ IT=100 *
$^{66}\text{As}^n$	-48480 680	3023.9 0.3	8.2 μs	0.5	(9 ⁺)	01Gr07	TJ IT=100 *
^{66}Se	-41720# 300#		33 ms	12	0 ⁺	98 02Lo13	TD β^+ =100
* ^{66}Mn	T : average 02So.A=64(2) 99Ha05=66(4) **						
* ^{66}Mn	T : also 99So20=62(14) 98Am04=90(20) outweighed, not used **						
* ^{66}Fe	T : average 99So20=440(60) 98Am04=440(60) **						
* ^{66}Co	T : average 00Mu10=180(10) 94Cz02=240(30) 85Bo49=230(20) **						
* $^{66}\text{As}^m$	J : 3 ⁺ # from systematics **						
* $^{66}\text{As}^n$	T : supersedes 98Gr12=17.5(1.5) E : from 98Gr12 **						
^{67}Cr	-19050# 700#		10# ms (>300 ns)			1/2 ⁻ # 97Be70	I β^- ?
^{67}Mn	-33400# 500#		45 ms	3		5/2 ⁻ # 97	02So.A TD β^- =100; β^-n ? *
^{67}Fe	-45690 420		394 ms	9		1/2 ⁻ # 91	02So.A TD β^- =100; β^-n ? *
$^{67}\text{Fe}^m$	-45320 420	367 3	64 μs	17		(5/2 ⁻) 03Sa02	ET IT=100 *
^{67}Co	-55060 320		425 ms	20		7/2 ⁻ # 91	99We07 T β^- =100 *
^{67}Ni	-63742.7 2.9		21 s	1		1/2 ⁻ 01	00Ri14 J β^- =100
$^{67}\text{Ni}^m$	-62736 4	1007 3	13.3 μs	0.2		9/2 ⁺ 01	98Gr14 E IT=100
^{67}Cu	-67318.8 1.2		61.83 h	0.12		3/2 ⁻ 91	β^- =100
^{67}Zn	-67880.4 0.9		STABLE			5/2 ⁻ 91	IS=4.10 13
^{67}Ga	-66879.7 1.3		3.2612 d	0.0006		3/2 ⁻ 96	ϵ =100
^{67}Ge	-62658 5		18.9 m	0.3		1/2 ⁻ 91	β^+ =100
$^{67}\text{Ge}^m$	-62640 5	18.2 0.05	13.7 μs	0.9		5/2 ⁻ 91	IT=100
$^{67}\text{Ge}^n$	-61906 5	751.70 0.06	110.9 ns	1.4		91	IT=100
^{67}As	-56650 100		42.5 s	1.2		(5/2 ⁻) 91	β^+ =100

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...							
^{67}Se	-46490# 200#		133 ms	11	5/2 ⁻ #	97 95B123	TD $\beta^+=100; \beta^+p=0.5$ 1 *
^{67}Br	-32800# 500#				1/2 ⁻ #		p ?
* ^{67}Mn	T: average 02So.A=47(4) 99Ha05=42(4)						**
* ^{67}Fe	T: others 99So20=500(100) 98Am04=470(50) outweighed, not used						**
* $^{67}\text{Fe}^m$	T: average 03Sa02=75(21) 98Gr14=43(30), same authors, different experiment						**
* ^{67}Co	T: others 99Pr10=440(70) 99So20=440(80) 85Bo49=420(70) outweighed, not used						**
* ^{67}Co	T: and 95Am.A=310(20) at variance, not used						**
* ^{67}Se	T: average 02Lo13=136(12) 94Ba50=107(35)						**
* ^{67}Se	T: values from 95B123 for $^{67}\text{Se}=60(+17-11)$ and ^{71}Kr questioned by 97Oio1						**
^{68}Mn	-28600# 600#		28 ms	4		02 02So.A	T $\beta^-=100; \beta^-n=?$ *
^{68}Fe	-43130 700		187 ms	6	0 ⁺	02 02So.A	T $\beta^-=100; \beta^-n?$ *
^{68}Co	-51350 320		* 200 ms	21	(7 ⁻)	02 00Mu10	T $\beta^-=100$ *
$^{68}\text{Co}^m$	-51200# 350#	150# 150#	* 1.6 s	0.3	(3 ⁺)	02 00Mu10	JD $\beta^-=?; IT?$ *
^{68}Ni	-63463.8 3.0		29 s	2	0 ⁺	02	$\beta^-=100$
$^{68}\text{Ni}^m$	-61694 3	1770.0 1.0	276 ns	65	0 ⁺	02	IT=100
$^{68}\text{Ni}^n$	-60615 3	2849.1 0.3	860 μs	50	5 ⁻	02	IT=100
^{68}Cu	-65567.0 1.6		31.1 s	1.5	1 ⁺	02	$\beta^-=100$
$^{68}\text{Cu}^m$	-64845.4 1.7	721.6 0.7	3.75 m	0.05	(6 ⁻)	02	IT=84 1; $\beta^-=16$ 1
^{68}Zn	-70007.2 1.0		STABLE		0 ⁺	02	IS=18.75 51
^{68}Ga	-67086.1 1.5		67.71 m	0.09	1 ⁺	02	$\beta^+=100$
$^{68}\text{Ga}^m$	-65856.2 1.5	1229.87 0.04	62.0 ns	1.4	7 ⁻	02	IT=100
^{68}Ge	-66980 6		270.95 d	0.16	0 ⁺	02	$\epsilon=100$
^{68}As	-58900 40		151.6 s	0.8	3 ⁺	02	$\beta^+=100$
$^{68}\text{As}^m$	-58470 40	425.21 0.16	111 s	20	1 ⁺	02	IT=100
^{68}Se	-54210 30		35.5 s	0.7	0 ⁺	02	$\beta^+=100$
^{68}Br	-38640# 360#		< 1.5 μs		3 ⁺ #	02 95B106	I p ?
* ^{68}Mn	T: average 02So.A=28(8) 99Ha05=28(4)						**
* ^{68}Fe	T: others 99So20=155(50) 91Be33=100(60) outweighed, not used						**
* ^{68}Co	T: average 00Mu10=230(30) 99So20=170(30); not used 95Am.A=310(30)						**
* ^{68}Co	T: 95Am.A supersedes 91Be33=180(100) from same group						**
^{69}Mn	-25300# 800#		14 ms	4	5/2 ⁻ #	00	$\beta^-=100; \beta^-n=24$ # *
^{69}Fe	-38400# 500#		109 ms	9	1/2 ⁻ #	00 02So.A	T $\beta^-=100; \beta^-n=7$ # *
^{69}Co	-50000 340		227 ms	13	7/2 ⁻ #	00 02So.A	T $\beta^-=100; \beta^-n=1$ # *
^{69}Ni	-59979 4		11.5 s	0.3	9/2 ⁺	00 99Pr10	T $\beta^-=100$ *
$^{69}\text{Ni}^m$	-59658 4	321 2	3.5 s	0.4	(1/2 ⁻)	00 98Gr14	E $\beta^-\approx 100; IT?$ *
$^{69}\text{Ni}^n$	-57278 11	2701 10	439 ns	3	(17/2 ⁻)	00	IT=100
^{69}Cu	-65736.2 1.4		2.85 m	0.15	3/2 ⁻	00	$\beta^-=100$
$^{69}\text{Cu}^m$	-62994.4 1.7	2741.8 1.0	360 ns	30	(13/2 ⁺)	00	IT=100
^{69}Zn	-68418.0 1.0		56.4 m	0.9	1/2 ⁻	00	$\beta^-=100$
$^{69}\text{Zn}^m$	-67979.4 1.0	438.636 0.018	13.76 h	0.02	9/2 ⁺	00	IT \approx 100; $\beta^-=0.033$ 3
^{69}Ga	-69327.8 1.2		STABLE		3/2 ⁻	00	IS=60.108 9
^{69}Ge	-67100.6 1.3		39.05 h	0.10	5/2 ⁻	00	$\beta^+=100$
$^{69}\text{Ge}^m$	-67013.8 1.3	86.765 0.014	5.1 μs	0.2	1/2 ⁻	00	IT=100
$^{69}\text{Ge}^n$	-66702.7 1.3	397.944 0.018	2.81 μs	0.05	9/2 ⁺	00	IT=100
^{69}As	-63090 30		15.2 m	0.2	5/2 ⁻	00	$\beta^+=100$
^{69}Se	-56300 30		27.4 s	0.2	(1/2 ⁻)	00 95Po01	J $\beta^+=100; \beta^+p=0.045$ 10
$^{69}\text{Se}^m$	-56260 30	39.4 0.1	2.0 μs	0.2	5/2 ⁻	00	IT=100
$^{69}\text{Se}^n$	-55730 30	573.9 1.0	955 ns	16	9/2 ⁺	00 00Ch07	T IT=100 *
^{69}Br	-46480# 110#		* < 24 ns		1/2 ⁻ #	00 96Pr01	I p ? *
$^{69}\text{Br}^m$	-46440# 150#	40# 100#			5/2 ⁻ #		
$^{69}\text{Br}^n$	-45910# 150#	570# 100#			9/2 ⁺ #		
... A-group is continued on next page ...							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
⁶⁹ Kr	-32440# 400#		32 ms	10	5/2 ⁻ #	00	$\beta^+=100; \beta^+p=?$	
* ⁶⁹ Mn	D: β^-n observed by 99Ha05							**
* ⁶⁹ Co	T: average 02So.A=232(17) 99Mu17=220(20); other 99So20=190(40), not used							**
* ⁶⁹ Ni	T: average 99Pr10=11.7(0.6) 85Bo49=11.4(0.3); not used 98Fr15=11.2(0.9)							**
* ⁶⁹ Ni ^m	T: average 99Mu17=3.5(0.5) 99Pr10=3.4(0.7)							**
* ⁶⁹ Ni ^m	E: 9/2 ⁺ level in isotones: ⁷³ Ge=-66 ⁷¹ Zn=157(1) 69Ni=-321(2) exhibits							**
* ⁶⁹ Ni ^m	E: unusual strong variations							**
* ⁶⁹ Se ⁿ	T: average 00Ch07=950(21) 95Po01=960(23)							**
* ⁶⁹ Br	T: in contradiction with 450 keV protons, 50 < T < 100 μ s reported in 88Ho.A							**
⁷⁰ Fe	-35900# 600#		94 ms	17	0 ⁺	97 02So.A	TD $\beta^-=100$	
⁷⁰ Co	-45640 840		* 125 ms	7	(6 ⁻ , 7 ⁻)	93 00Mu10	TJD $\beta^-=100; \beta^-n?$	
⁷⁰ Co ^m	-45440# 860# 200# 200#		* 500 ms	180	(3 ⁺)	00Mu10	TJD $\beta^-\approx 100; IT?; \beta^-n?$	
⁷⁰ Ni	-59150 350		6.0 s	0.3	0 ⁺	03 98Fr15	TD $\beta^-=100$	
⁷⁰ Ni ^m	-56290 350 2860 2		232 ns	1	8 ⁺	03	IT=100	
⁷⁰ Cu	-62976.1 1.6		& 44.5 s	0.2	(6 ⁻)	93 02We03	TJ $\beta^-=100$	
⁷⁰ Cu ^m	-62875.4 2.0 100.7 2.6 MD		33 s	2	(3 ⁻)	02We03	TJ $\beta^-\approx 50; IT\approx 50$	
⁷⁰ Cu ⁿ	-62734.1 2.1 242.0 2.7 MD &		6.6 s	0.2	1 ⁺	93 02We03	TD $\beta^-\approx 95; IT\approx 5$	
⁷⁰ Zn	-69564.6 2.0		STABLE		0 ⁺	93	IS=0.62 3; 2 $\beta^-?$	
⁷⁰ Ga	-68910.1 1.2		21.14 m	0.03	1 ⁺	93	$\beta^-\approx 100; \epsilon=0.41$ 6	
⁷⁰ Ge	-70563.1 1.0		STABLE		0 ⁺	93	IS=20.84 87	
⁷⁰ As	-64340 50		52.6 m	0.3	4 ⁽⁺⁾ #	93	$\beta^+=100$	
⁷⁰ As ^m	-64310 50 32.06 0.03		96 μ s	3	2 ⁽⁺⁾	93	IT=100	
⁷⁰ Se	-62050 60		41.1 m	0.3	0 ⁺	93	$\beta^+=100$	
⁷⁰ Br	-51430# 310#		79.1 ms	0.8	0 ⁺ #	93	$\beta^+=100$	
⁷⁰ Br ^m	-49140# 310# 2292.2 0.8		2.2 s	0.2	(9 ⁺)	93 00Pi15	J $\beta^+=?; IT?$	
⁷⁰ Kr	-41680# 390#		57 ms	21	0 ⁺	97 00Oi02	TD $\beta^+?$	
* ⁷⁰ Co	T: average 02So.A=121(8) 98Am04=150(20); others 00Mu10=120(30) 99So20=92(25)							**
* ⁷⁰ Cu ⁿ	D: IT=0.3 percent E: post deadline 03Va.2 101.1(0.3) and 242.4(0.3)							**
* ⁷⁰ Zn	T: > 500 Ty in ENSDF is for 0v-2 β^- decay alone							**
* ⁷⁰ Br ⁿ	E: from 2002Je07							**
⁷¹ Fe	-31000# 800#		30# ms (>300 ns)		7/2 ⁺ #	97 97Be70	I $\beta^-?$	
⁷¹ Co	-43870 840		97 ms	2	7/2 ⁻ #	93 02So.A	T $\beta^-=100; \beta^-n?$	
⁷¹ Ni	-55200 370		2.56 s	0.03	1/2 ⁻ #	93 98Fr15	T $\beta^-=100$	
⁷¹ Cu	-62711.1 1.5		19.4 s	1.4	(3/2 ⁻)	93 99Pr10	T $\beta^-=100$	
⁷¹ Cu ^m	-59955 10 2756 10		271 ns	13	(19/2 ⁻)	98Gr14	ETJ IT=100	
⁷¹ Zn	-67327 10		2.45 m	0.10	1/2 ⁻	93	$\beta^-=100$	
⁷¹ Zn ^m	-67169 10 157.7 1.3		3.96 h	0.05	9/2 ⁺	93	$\beta^-\approx 100; IT\leq 0.05$	
⁷¹ Ga	-70140.2 1.0		STABLE		3/2 ⁻	93	IS=39.892 9	
⁷¹ Ge	-69907.7 1.0		11.43 d	0.03	1/2 ⁻	93	$\epsilon=100$	
⁷¹ Ge ^m	-69709.3 1.0 198.367 0.010		20.40 ms	0.17	9/2 ⁺	93	IT=100	
⁷¹ As	-67894 4		65.28 h	0.15	5/2 ⁻	93	$\beta^+=100$	
⁷¹ Se	-63120 30		4.74 m	0.05	5/2 ⁻	93	$\beta^+=100$	
⁷¹ Se ^m	-63070 30 48.79 0.05		5.6 μ s	0.7	1/2 ⁻ to 9/2 ⁻	93	IT=100	
⁷¹ Se ⁿ	-62860 30 260.48 0.10		19.0 μ s	0.5	(9/2 ⁺)	93 00Ch07	T IT=100	
⁷¹ Br	-57060 570		21.4 s	0.6	(5/2 ⁻)	93	$\beta^+=100$	
⁷¹ Kr	-46920 650		100 ms	3	(5/2 ⁻)	97 97Oi01	TJD $\beta^+=100; \beta^+p=2.1$ 7	
⁷¹ Rb	-32300# 500#		*		5/2 ⁻ #		p?	
⁷¹ Rb ^m	-32250# 510# 50# 100#		*		1/2 ⁻ #			
⁷¹ Rb ⁿ	-32040# 510# 260# 100#				9/2 ⁺ #			
* ⁷¹ Co	T: other not used: 98Am04=210(40)							**
* ⁷¹ Cu	T: average 99Pr10=19(3) 83Ru06=19.5(1.6)							**
* ⁷¹ Cu ^m	T: average 98Is11=250(30) 98Gr14=275(14)							**
* ⁷¹ Kr	T: average 97Oi01=100(3) 81Ew01=97(9); 95Bi23=64(+8-5) at variance not used							**
* ⁷¹ Kr	T: values from 95Bi23 for ⁶⁷ Se and ⁷¹ Kr questioned by 97Oi01							**
* ⁷¹ Kr	D: 95Bi23=5.2(0.6) at variance not used							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{72}Fe	-28300# 800#		10# ms (>300 ns)	0^+	97	97Be70 I	$\beta^- ?$	
^{72}Co	-39300# 600#		90 ms	20		98Am04 TD	$\beta^- = 100; \beta^- n ?$	
^{72}Ni	-53940 440		1.57 s	0.05	0^+	98Fr15 TD	$\beta^- = 100; \beta^- n ?$ *	
^{72}Cu	-59783.0 1.4		6.6 s	0.1	(1^+)	95	$\beta^- = 100$	
$^{72}\text{Cu}^m$	-59513 3 270	3	1.76 μs	0.03	(4^-)	98Gr14 ETJ	IT=100	
^{72}Zn	-68131 6		46.5 h	0.1	0^+	95	$\beta^- = 100$	
^{72}Ga	-68589.4 1.0		14.10 h	0.02	3^-	95	$\beta^- = 100$	
$^{72}\text{Ga}^m$	-68469.7 1.0 119.66	0.05	39.68 ms	0.13	(0^+)	95	IT=100	
^{72}Ge	-72585.9 1.6		STABLE		0^+	95	IS=27.54 34	
$^{72}\text{Ge}^m$	-71894.5 1.6 691.43	0.04	444.2 ns	0.8	0^+			
^{72}As	-68230 4		26.0 h	0.1	2^-	95	$\beta^+ = 100$	
^{72}Se	-67894 12		8.40 d	0.08	0^+	97	$\epsilon = 100$	
^{72}Br	-59020 60		78.6 s	2.4	1^+	95 03Pi03 J	$\beta^+ = 100$	
$^{72}\text{Br}^m$	-58920 60 100.92	0.03	10.6 s	0.3	1^-	95	IT=100; $\beta^+ = ?$	
^{72}Kr	-53941 8		17.16 s	0.18	0^+	95 03Pi03 T	$\beta^+ = 100$ *	
^{72}Rb	-38120# 500#		* < 1.5 μs		$3^+\#$	97 95Bi06 I	p ?	
$^{72}\text{Rb}^m$	-38020# 510# 100#	100#	* 1# μs		1^-		p ?	
* ^{72}Ni	T : not used 95Am.A=1.30(0.10) and 92Be.A=2.06(0.30) (the two of same group)							**
* ^{72}Kr	T : average 03Pi03=17.1(0.2) 73Da22=17.4(0.4)							**
^{73}Co	-37040# 700#		80# ms (>300 ns)	$7/2^- \#$	02	97Be70 I	$\beta^- ?$	
^{73}Ni	-49860# 300#		840 ms	30	$(9/2^+)$	02	$\beta^- = 100; \beta^- n ?$	
^{73}Cu	-58987 4		4.2 s	0.3	$(3/2^-)$	02 98Fr15 J	$\beta^- = 100; \beta^- n ?$	
^{73}Zn	-65410 40		23.5 s	1.0	$(1/2^-)$	02	$\beta^- = 100$	
$^{73}\text{Zn}^m$	-65210 40 195.5	0.2	13.0 ms	0.2	$(5/2^+)$	02	IT=100	
$^{73}\text{Zn}^m$	-65170 40 237.6	2.0	EU 5.8 s	0.8	$(7/2^+)$	02	IT=?; $\beta^- = ?$ *	
^{73}Ga	-69699.3 1.7		4.86 h	0.03	$3/2^-$	02	$\beta^- = 100$	
^{73}Ge	-71297.5 1.6		STABLE		$9/2^+$	02	IS=7.73 5	
$^{73}\text{Ge}^m$	-71284.2 1.6 13.2845	0.0015	2.92 μs	0.03	$5/2^+$	02	IT=100	
$^{73}\text{Ge}^n$	-71230.8 1.6 66.726	0.009	499 ms	11	$1/2^-$	02	IT=100	
^{73}As	-70957 4		80.30 d	0.06	$3/2^-$	93	$\epsilon = 100$	
^{73}Se	-68218 11		7.15 h	0.08	$9/2^+$	03	$\beta^+ = 100$	
$^{73}\text{Se}^m$	-68192 11 25.71	0.04	39.8 m	1.3	$3/2^-$	03	IT=72.6 3; $\beta^+ = 27.4 3$	
^{73}Br	-63630 50		3.4 m	0.2	$1/2^-$	02	$\beta^+ = 100$	
^{73}Kr	-56552 7		28.6 s	0.6	$3/2^-$	02 99Mi17 T	$\beta^+ = 100; \beta^+ p = 0.25 3$ *	
$^{73}\text{Kr}^m$	-56118 7 433.66	0.12	107 ns	10	$(9/2^+)$	03	IT=100	
^{73}Rb	-46050# 150#		< 30 ns		$3/2^- \#$	03 96Pf01 I	p ?	
$^{73}\text{Rb}^m$	-45620# 180# 430#	100#			$9/2^+ \#$			
^{73}Sr	-31700# 600#		> 25 ms		$1/2^- \#$	03	$\beta^+ = 100; \beta^+ p = ?$	
* $^{73}\text{Zn}^n$	E : if 42.1 keV γ feeds $^{73}\text{Zn}^m$, EU: see discussion in ENSDF'02							**
* ^{73}Kr	T : average 99Mi17=29.0(1.0) 81Ha44=28.4(0.7); 73Da22=25.9(0.6) at variance,							**
* ^{73}Kr	T : not used							**
^{74}Co	-32250# 800#		50# ms (>300 ns)		03	97Be70 I	$\beta^- ?$	
^{74}Ni	-48370# 400#		680 ms	120	0^+	03 98Fr15 T	$\beta^- = 100; \beta^- n ?$ *	
^{74}Cu	-56006 6		1.594 s	0.010	$1^+\#$	95	$\beta^- = 100$	
^{74}Zn	-65710 50		95.6 s	1.2	0^+	95	$\beta^- = 100$	
^{74}Ga	-68050 4		8.12 m	0.12	(3^-)	95	$\beta^- = 100$	
$^{74}\text{Ga}^m$	-67990 4 59.571	0.014	9.5 s	1.0	(0)	95	IT=?; $\beta^- = 25\#$	
^{74}Ge	-73422.4 1.6		STABLE		0^+	95	IS=36.28 73	
^{74}As	-70860.0 2.3		17.77 d	0.02	2^-	95	$\beta^+ = 66 2; \beta^- = 34 2$	
^{74}Se	-72212.7 1.7		STABLE		0^+	95	IS=0.89 4; $2\beta^+ ?$	
^{74}Br	-65306 15		25.4 m	0.3	(0^-)	95	$\beta^+ = 100$	
$^{74}\text{Br}^m$	-65292 15 13.58	0.21	46 m	2	$4^{(++)}$	95	$\beta^+ = 100$	
^{74}Kr	-62331.5 2.0		11.50 m	0.11	0^+	95	$\beta^+ = 100$	
$^{74}\text{Kr}^m$	-61824 10 508	10	29 ns	6	0^+	00Ch07 ETJ	IT=100	
^{74}Rb	-51917 4		64.76 ms	0.03	(0^+)	95 01Ba12 T	$\beta^+ = 100$	
^{74}Sr	-40700# 500#		50# ms (>1.5 μs)		0^+	97 95Bi06 I	$\beta^+ ?$	
* ^{74}Ni	T : average 98Fr15=900(200) 98Am04=540(160)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
⁷⁵ Co	-29500# 800#		40# ms (>300 ns)	7/2 ⁻ #	99	97Be70 I	β^- ?	
⁷⁵ Ni	-43900# 400#		600 ms	200	7/2 ⁺ #	99 85Re01 D	β^- =100; β^- -n=1.6# *	
⁷⁵ Cu	-54120 980		1.224 s	0.003	3/2 ⁻ #	99	β^- =100; β^- -n=3.5 6	
⁷⁵ Zn	-62470 70		10.2 s	0.2	7/2 ⁺ #	99	β^- =100	
⁷⁵ Ga	-68464.6 2.4		126 s	2	(3/2) ⁻	99	β^- =100	
⁷⁵ Ge	-71856.4 1.6		82.78 m	0.04	1/2 ⁻	99	β^- =100	
⁷⁵ Ge ^m	-71716.7 1.6	139.69 0.03	47.7 s	0.5	7/2 ⁺	99	IT \approx 100; β^- =0.030 6	
⁷⁵ As	-73032.4 1.8		STABLE		3/2 ⁻	99	IS=100.	
⁷⁵ As ^m	-72728.5 1.8	303.9241 0.0007	17.62 ms	0.23	9/2 ⁺	99	IT=100	
⁷⁵ Se	-72169.0 1.7		119.779 d	0.004	5/2 ⁺	99	ϵ =100	
⁷⁵ Br	-69139 14		96.7 m	1.3	3/2 ⁻	99	β^+ =100	
⁷⁵ Kr	-64324 8		4.29 m	0.17	5/2 ⁺	99	β^+ =100	
⁷⁵ Rb	-57222 7		19.0 s	1.2	(3/2 ⁻)	99	β^+ =100	
⁷⁵ Sr	-46620 220		88 ms	3	(3/2 ⁻)	99 03Hu01 TJD	β^+ =100; β^+ -p=5.2 9	
* ⁷⁵ Ni	D : β^- -n=1.6%# estimated by 85Re01							**
⁷⁶ Ni	-41610# 900#		470 ms	390	0 ⁺	97 98Am04 T	β^- =100; β^- -n ?	
⁷⁶ Cu	-50976 7		* 641 ms	6	(3,5)	95 90Wi12 J	β^- =100; β^- -n=3 2	
⁷⁶ Cu ^m	-50980# 200#	0# 200#	* 1.27 s	0.30	(1,3)	95 90Wi12 J	β^- =100	
⁷⁶ Zn	-62140 80		5.7 s	0.3	0 ⁺	95	β^- =100	
⁷⁶ Ga	-66296.6 2.0		32.6 s	0.6	(2 ⁺ , 3 ⁺)	95	β^- =100	
⁷⁶ Ge	-73213.0 1.7		1.58 Zy	0.17	0 ⁺	95 01K111 T	IS=7.61 38; 2 β^- =100 *	
⁷⁶ As	-72289.5 1.8		1.0778 d	0.0020	2 ⁻	95	β^- \approx 100; ϵ <0.02	
⁷⁶ As ^m	-72245.1 1.8	44.425 0.001	1.84 μ s	0.06	(1) ⁺			
⁷⁶ Se	-75252.1 1.7		STABLE		0 ⁺	95	IS=9.37 29	
⁷⁶ Br	-70289 9		16.2 h	0.2	1 ⁻	95	β^+ =100	
⁷⁶ Br ^m	-70186 9	102.58 0.03	1.31 s	0.02	(4) ⁺	95	IT>99.4; β^+ <0.6	
⁷⁶ Kr	-69014 4		14.8 h	0.1	0 ⁺	95	β^+ =100	
⁷⁶ Rb	-60479.8 1.9		36.5 s	0.6	1 ⁽⁻⁾	95 78Ha08 D	β^+ =100; β^+ - α =3.8e-7 10	
⁷⁶ Rb ^m	-60162.9 1.9	316.93 0.08	3.050 μ s	0.007	(4) ⁺	95 00Ch07 T	IT=100	
⁷⁶ Sr	-54240 40		8.9 s	0.3	0 ⁺	95	β^+ =100	
⁷⁶ Y	-38700# 500#		500# ns (>170 ns)			00We.A I	β^+ ?; p ? *	
* ⁷⁶ Ge	T : from 01K111=1.55(+0.19-0.15); other results from same group:							**
* ⁷⁶ Ge	T : ⁹⁷ Ga13=1.77(+0.13-0.11) ⁹⁴ Ba15=1.42(0.13)							**
* ⁷⁶ Ge	T : other groups ⁹³ Br22=0.84(+0.10-0.08)(2 σ) ⁹⁰ Va18=0.90(0.10)							**
* ⁷⁶ Ge	T : and ⁹⁰ Mi23=1.1(+0.6-0.3)(2 σ)							**
* ⁷⁶ Ge	TD : claim for 0 ν - $\beta\beta$ 01K113=15 Yy not trusted. See also 02Aa.1 and 02Zd02							**
* ⁷⁶ Y	I : also 01Ki13>200 ns, same group							**
⁷⁷ Ni	-36750# 500#		300# ms (>300 ns)	9/2 ⁺ #	97	97Be70 I	β^- ?	
⁷⁷ Cu	-48580# 400#		469 ms	8	3/2 ⁻ #	97	β^- =100	
⁷⁷ Zn	-58720 120		2.08 s	0.05	7/2 ⁺ #	97	β^- =100	
⁷⁷ Zn ^m	-57950 120	772.39 0.12	1.05 s	0.10	1/2 ⁻ #	97	IT>50; β^- <50	
⁷⁷ Ga	-65992.3 2.4		13.2 s	0.2	(3/2 ⁻)	97	β^- =100	
⁷⁷ Ge	-71214.0 1.7		11.30 h	0.01	7/2 ⁺	97	β^- =100	
⁷⁷ Ge ^m	-71054.3 1.7	159.70 0.10	52.9 s	0.6	1/2 ⁻	97	β^- =81 2; IT=19 2	
⁷⁷ As	-73916.6 2.3		38.83 h	0.05	3/2 ⁻	97	β^- =100	
⁷⁷ As ^m	-73441.2 2.3	475.443 0.016	114.0 μ s	2.5	9/2 ⁺	97	IT=100	
⁷⁷ Se	-74599.6 1.7		STABLE		1/2 ⁻	97	IS=7.63 16	
⁷⁷ Se ^m	-74437.7 1.7	161.9223 0.0007	17.36 s	0.05	7/2 ⁺	97	IT=100	
⁷⁷ Br	-73235 3		57.036 h	0.006	3/2 ⁻	97	β^+ =100	
⁷⁷ Br ^m	-73129 3	105.86 0.08	4.28 m	0.10	9/2 ⁺	97	IT=100	
⁷⁷ Kr	-70169.4 2.0		74.4 m	0.6	5/2 ⁺	97	β^+ =100	
⁷⁷ Rb	-64825 7		3.77 m	0.04	3/2 ⁻	97	β^+ =100	
⁷⁷ Sr	-57804 9		9.0 s	0.2	5/2 ⁺	97	β^+ =100; β^+ -p<0.25	
⁷⁷ Y	-46910# 60#		63 ms	17	5/2 ⁺ #	97 01Ki13 T	β^+ =?; β^+ -p ?; p<10 *	
* ⁷⁷ Y	D : limit for p is from 00We.A							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{78}Ni	-34300# 1100#		200#	ms (>300 ns)	0 ⁺	97 97Be70 I	β^- ?	
^{78}Cu	-44750# 400#		342	ms	11	97 91Kr15 T	β^- =100	
^{78}Zn	-57340 90		1.47	s	0.15	91	β^- =100	
$^{78}\text{Zn}^m$	-54670 90	2673 1	319	ns	9 (8 ⁺)	00Da07 ET	IT=100	
^{78}Ga	-63706.6 2.4		5.09	s	0.05 (3 ⁺)	91	β^- =100	
^{78}Ge	-71862 4		88	m	1	91	β^- =100	
^{78}As	-72817 10		90.7	m	0.2	2- 91	β^- =100	
^{78}Se	-77026.1 1.7		STABLE			0+ 91	IS=23.77 28	
^{78}Br	-73452 4		6.46	m	0.04	1+ 91	β^+ ≈100; β^- <0.01	
$^{78}\text{Br}^m$	-73271 4	180.82 0.13	119.2	μs		4+		
^{78}Kr	-74179.7 1.1		STABLE		(>110 Ey)	0+ 91	94Sa31 T IS=0.35 1; 2 β^+ ?	
^{78}Rb	-66936 7		17.66	m	0.08	0(+) 91	β^+ =100	
$^{78}\text{Rb}^m$	-66825 7	111.20 0.10	5.74	m	0.05	4(-) 91	91Mc.A E β^+ =90 2; IT=10 2	
$^{78}\text{Rb}^x$	-66862 14	74 12	R = 2.0 0.5			spmix		
^{78}Sr	-63174 7		159	s	8	0+ 91	92Gr09 T β^+ =100	
^{78}Y	-52530# 400#		54	ms	5 (0 ⁺)	97 01Ga24 TJD	β^+ =100; β^+ p ?	
$^{78}\text{Y}^m$	-52530# 640#	0# 500#	5.8	s	0.5	5+#	01Ki13 TD β^+ =100; β^+ p ?	
^{78}Zr	-41700# 500#		50#	ms (>170 ns)	0 ⁺	00We.A I	β^+ ?; β^+ p ?	
* ^{78}Br	D : β^- branch is uncertain. See ENSDF							**
* ^{78}Kr	T : limit given here is for the K-e ⁺ decay (theoretically faster)							**
* ^{78}Y	T : average 01Ga24=50(8) 01Ki13=55(+9-6)							**
* $^{78}\text{Y}^m$	T : average 01Ki13=5.7(0.7) 98Uu01=5.8(0.6)							**
* ^{78}Zr	I : also 01Ki13>200 ns same group							**
^{79}Cu	-42330# 500#		188	ms	25	3/2-# 02	β^- =100; β^- n=55 17	
^{79}Zn	-53420# 260#		995	ms	19	(9/2 ⁺) 02	β^- =100; β^- n=1.3 4	
^{79}Ga	-62510 100		2.847	s	0.003	3/2-# 02	β^- =100; β^- n=0.089 19	
^{79}Ge	-69490 90		18.98	s	0.03	(1/2)- 02	β^- =100	
$^{79}\text{Ge}^m$	-69300 90	185.95 0.04	39.0	s	1.0	7/2 ⁺ # 02	β^- =96 1; IT=4 1	
^{79}As	-73637 6		9.01	m	0.15	3/2- 02	β^- =100	
$^{79}\text{As}^m$	-72864 6	772.81 0.06	1.21	μs	0.01	(9/2 ⁺) 02	98Gr14 T IT=100	
^{79}Se	-75917.6 1.7		295	ky	38	7/2 ⁺ 02	β^- =100	
$^{79}\text{Se}^m$	-75821.8 1.7	95.77 0.03	3.92	m	0.01	1/2- 02	IT≈100; β^- =0.056 11	
^{79}Br	-76068.5 2.0		STABLE			3/2- 02	IS=50.69 7	
$^{79}\text{Br}^m$	-75860.9 2.0	207.61 0.09	4.86	s	0.04	(9/2 ⁺) 02	IT=100	
^{79}Kr	-74443 4		35.04	h	0.10	1/2- 02	β^+ =100	
$^{79}\text{Kr}^m$	-74313 4	129.77 0.05	50	s	3	7/2 ⁺ 02	IT=100	
$^{79}\text{Kr}^n$	-74296 4	147.06 0.06	78.7	ns	1.0	(5/2-) 02	IT=100	
^{79}Rb	-70803 6		22.9	m	0.5	5/2 ⁺ 02	β^+ =100	
^{79}Sr	-65477 8		2.25	m	0.10	3/2(-) 02	β^+ =100	
^{79}Y	-58360 450		14.8	s	0.6	5/2 ⁺ # 02	β^+ =100; β^+ p ?	
^{79}Zr	-47360# 400#		56	ms	30	5/2 ⁺ # 02	β^+ =100; β^+ p ?	
* $^{79}\text{As}^m$	T : 98Ho15=0.87(0.06) outweighed, not used							**
^{80}Cu	-36450# 600#		100#	ms (>300 ns)		97 97Be70 I	β^- ?	
^{80}Zn	-51840 170		545	ms	16	0+ 92	β^- =100; β^- n=1.0 5	
^{80}Ga	-59140 120		1.697	s	0.011	(3) 92	93Ru01 D β^- =100; β^- n=0.89 6	
^{80}Ge	-69515 28		29.5	s	0.4	0+ 92	β^- =100	
^{80}As	-72159 23		15.2	s	0.2	1+ 92	β^- =100	
^{80}Se	-77759.9 2.0		STABLE			0+ 92	IS=49.61 41; 2 β^- ?	
^{80}Br	-75889.5 2.0		17.68	m	0.02	1+ 92	β^- =91.7 2; β^+ =8.3 2	
$^{80}\text{Br}^m$	-75803.7 2.0	85.843 0.004	4.4205	h	0.0008	5- 92	IT=100	
^{80}Kr	-77892.5 1.5		STABLE			0+ 92	IS=2.28 6	
^{80}Rb	-72173 7		33.4	s	0.7	1+ 92	93Al03 T β^+ =100	
$^{80}\text{Rb}^m$	-71679 7	494.4 0.5	1.6	μs	0.02	6+	92Do10 E	
^{80}Sr	-70308 7		106.3	m	1.5	0+ 99	β^+ =100	
^{80}Y	-61220 180		30.1	s	0.5	4- 92	98Do04 TJ β^+ =100	
$^{80}\text{Y}^m$	-60990 180	228.5 0.1	4.8	s	0.3	(1-) 98Do04 ETJ	IT=81 2; β^+ =19 2	
^{80}Zr	-60910 180	312.5 1.0	4.7	μs	0.3	(2+) 00Ch07 ETJ	IT=100	

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
^{80}Zr	-5520	1490	4.6 s	0.6	0 ⁺	92 01Ki13 T	$\beta^+=100; \beta^+p?$ *	
* ^{80}Y	T : differences with $^{82}\text{De}36=38(1)$ $^{81}\text{Li}12=33.8(0.6)$ explained in 98Do04 **							
* $^{80}\text{Y}^m$	T : average $^{01}\text{No}07=5.0(0.5)$ $^{98}\text{Do}04=4.7(0.3)$ D : from 98Do04 **							
* $^{80}\text{Y}^m$	E : $^{00}\text{Ch}07=84(1)$ above 228.5 level **							
* ^{80}Zr	T : average $^{01}\text{Ki}13=5.3(+1.1-0.9)$ $^{00}\text{Re}03=4.1(+0.8-0.6)$ **							
^{81}Zn	-46130#	300#	290 ms	50	$5/2^+\#$	97	$\beta^-=100; \beta^-n=7.5$ 30	
^{81}Ga	-57980	190	1.217 s	0.005	$(5/2^-)$	97	$\beta^-=100; \beta^-n=11.9$ 7	
^{81}Ge	-66300	120	8 s	2	$9/2^+\#$	97	$\beta^-=100$ *	
$^{81}\text{Ge}^m$	-65620	120	679.13	0.04	8 s	2	$(1/2^+)$ 97	
^{81}As	-72533	6	33.3 s	0.8	$3/2^-$	97	$\beta^-=100$	
^{81}Se	-76389.5	2.0	18.45 m	0.12	$1/2^-$	97	$\beta^-=100$	
$^{81}\text{Se}^m$	-76286.5	2.0	102.99	0.06	57.28 m	0.02	$7/2^+$ 97	
^{81}Br	-77974.8	2.0	STABLE		$3/2^-$	97	$\text{IT}\approx 100; \beta^-=0.052$ 14	
$^{81}\text{Br}^m$	-77438.6	2.0	536.20	0.09	34.6 μs		$\text{IS}=49.31$ 7	
^{81}Kr	-77694.0	2.0	229 ky	11	$7/2^+$	97	$\epsilon=100$	
$^{81}\text{Kr}^m$	-77503.4	2.0	190.62	0.04	13.10 s	0.03	$\text{IT}\approx 100; \epsilon=0.0025$ 4	
^{81}Rb	-75455	6	4.576 h	0.005	$3/2^-$	97	$\beta^+=100$	
$^{81}\text{Rb}^m$	-75369	6	86.31	0.07	30.5 m	0.3	$\text{IT}=97.6$ 6; $\beta^+=2.4$ 6	
^{81}Sr	-71528	6	22.3 m	0.4	$1/2^-$	99	$\beta^+=100$	
^{81}Y	-66020	60	70.4 s	1.0	$(5/2^+)$	98	$\beta^+=100$	
^{81}Zr	-58490	170	5.5 s	0.4	$3/2^-$	# 00	$\beta^+=100; \beta^+p=0.12$ 2	
^{81}Nb	-47480#	1500#	< 44 ns		$3/2^-$	# 97	00We.A I $p?; \beta^+?; \beta^+p?$ *	
* ^{81}Ge	T : derived from $7.6(0.6)$, for mixture of ground-state and isomer with almost same half-life **							
* ^{81}Nb	I : also $^{99}\text{Ja}02<80$ $^{01}\text{Ki}13<200$ ns T : estimated half-life for β^+ : 100 ms **							
^{82}Zn	-42460#	500#	100# ms	(>300 ns)	0 ⁺	03 97Be70 I	$\beta^-?$	
^{82}Ga	-53100#	300#	599 ms	2	$(1, 2, 3)$	03 93Ru01 D	$\beta^-=100; \beta^-n=21.3$ 13 *	
^{82}Ge	-65620	240	4.55 s	0.05	0 ⁺	03	$\beta^-=100$	
^{82}As	-70320	200	19.1 s	0.5	(1^+)	03	$\beta^-=100$	
$^{82}\text{As}^m$	-70075	25	250	200	BD *	13.6 s	0.4	(5^-) 03
^{82}Se	-77594.0	2.0	97 Ey	5	0 ⁺	03 99Pi08 T	$\text{IS}=8.73$ 22; $2\beta^-=100$ *	
^{82}Br	-77496.5	1.9	35.282 h	0.007	5 ⁻	03	$\beta^-=100$	
$^{82}\text{Br}^m$	-77450.6	1.9	45.9492	0.0010	6.13 m	0.05	2^- 03	
^{82}Kr	-80589.5	1.8	STABLE		0 ⁺	03	$\text{IT}=97.6$ 3; $\beta^-=2.4$ 3	
^{82}Rb	-76188.2	2.8	1.273 m	0.002	1 ⁺	03	$\beta^+=100$	
$^{82}\text{Rb}^m$	-76119.1	2.4	69.1	1.5	MD	6.472 h	0.006	5 ⁻ 03
^{82}Sr	-76008	6	25.36 d	0.03	0 ⁺	03 87Ho06 T	$\beta^+\approx 100; \text{IT}<0.33$	
^{82}Y	-68190	100	8.30 s	0.20	1 ⁺	03	$\epsilon=100$ *	
$^{82}\text{Y}^m$	-67790	100	402.63	0.14	268 ns	25	4 ⁻ 03	
^{82}Zr	-64190#	230#	32 s	5	0 ⁺	03	$\beta^+=100$	
^{82}Nb	-52970#	300#	51 ms	5	0 ⁺	03 01Ga24 T	$\beta^+=100; \beta^+p?$ *	
* ^{82}Ga	D : average $^{93}\text{Ru}01=31.1(4.4)$ $^{86}\text{Wa}17=19.8(1.7)$ $^{80}\text{Lu}04=21.4(2.2)$ **							
* ^{82}Se	T : average $^{99}\text{Pi}08=83(+9-7)$ $^{98}\text{Ar}10=83(12)$ $^{92}\text{El}07=108(+26-6)$ $^{88}\text{Li}11=120(10)$ **							
* ^{82}Sr	T : average $^{87}\text{Ho}06=25.36(0.03)$ $^{87}\text{Ju}02=25.342(0.053)$ **							
* ^{82}Nb	T : average $^{01}\text{Ga}24=52(6)$ $^{01}\text{Ki}13=48(+8-6)$ **							
^{83}Zn	-36300#	500#	80# ms	(>300 ns)	$5/2^+\#$	01 97Be70 I	$\beta^-?$	
^{83}Ga	-49390#	300#	308 ms	1	$3/2^-$	# 01	$\beta^-=100; \beta^-n=37$ 17	
^{83}Ge	-60900#	200#	1.85 s	0.06	$5/2^+\#$	01	$\beta^-=100$	
^{83}As	-69880	220	13.4 s	0.3	$3/2^-$	# 01	$\beta^-=100$	
^{83}Se	-75341	4	22.3 m	0.3	$9/2^+$	01	$\beta^-=100$	
$^{83}\text{Se}^m$	-75113	4	228.50	0.20	70.1 s	0.4	$1/2^-$ 01	
^{83}Br	-79009	4	2.40 h	0.02	$3/2^-$	01	$\beta^-=100$	
$^{83}\text{Br}^m$	-75940	4	3068.8	0.6	700 ns	100	$(19/2^-)$ 01	
^{83}Kr	-79981.7	2.8	STABLE		$9/2^+$	01	$\text{IS}=11.49$ 6	
$^{83}\text{Kr}^m$	-79972.3	2.8	9.4053	0.0008	154.4 ns	1.1	$7/2^+$ 01	
$^{83}\text{Kr}^n$	-79940.1	2.8	41.5569	0.0010	1.83 h	0.02	$1/2^-$ 01	
... A-group is continued on next page ...								

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...									
^{83}Rb	-79075	6		86.2	d	0.1	5/2 ⁻	01	$\epsilon=100$
$^{83}\text{Rb}^m$	-79033	6	42.11 0.04	7.8	ms	0.7	9/2 ⁺	01 68Et01 T	IT=100
^{83}Sr	-76795	10		32.41	h	0.03	7/2 ⁺	01	$\beta^+=100$
$^{83}\text{Sr}^m$	-76536	10	259.15 0.09	4.95	s	0.12	1/2 ⁻	01	IT=100
^{83}Y	-72330	40		7.08	m	0.06	9/2 ⁺	01 92Bu10 J	$\beta^+=100$
$^{83}\text{Y}^m$	-72270	40	61.98 0.11	2.85	m	0.02	(3/2 ⁻)	01	$\beta^+=60.5$; IT=40.5
^{83}Zr	-66460	100		41.6	s	2.4	1/2 ⁻	# 01	$\beta^+=100$; $\beta^+p=?$
$^{83}\text{Zr}^m$	-66410	100	52.72 0.05	530	ns	0.12	(5/2 ⁻)	01	IT=100
$^{83}\text{Zr}^i$			non existent	8	s	1	high	01	$\beta^+=100$; $\beta^+p=?$
^{83}Nb	-58960	310		4.1	s	0.3	(5/2 ⁺)	01	$\beta^+=100$
^{83}Mo	-47750#	500#		23	ms	19	3/2 ⁻	# 01 01Ki13 TD	$\beta^+=100$; $\beta^+p?$
$^{83}\text{Zr}^i$	D : 6(4)% of total β^+p go to first excited state in ^{82}Sr								
$^{83}\text{Zr}^i$	I : misassigned: absence of radiations suggests no isomer with E>18 keV								
^{84}Ga	-44110#	400#		85	ms	10		97	$\beta^-=100$; $\beta^-n=70.15$
^{84}Ge	-58250#	300#		954	ms	14	0 ⁺	97 93Ru01 T	$\beta^-=100$; $\beta^-n=10.8.6$
^{84}As	-66080#	300#		4.02	s	0.03	(3)(+)	97 93Ru01 T	$\beta^-=100$; $\beta^-n=0.28.4$
$^{84}\text{As}^m$	-66080#	320#	0# 100#	650	ms	150		97	$\beta^-=100$
^{84}Se	-75952	15		3.1	m	0.1	0 ⁺	97	$\beta^-=100$
^{84}Br	-77799	15		31.80	m	0.08	2 ⁻	97	$\beta^-=100$
$^{84}\text{Br}^m$	-77460	100	340 100	6.0	m	0.2	(6 ⁻)	97	$\beta^-=100$
$^{84}\text{Br}^i$	-77391	15	408.2 0.4	< 140	ns		1 ⁺	97	IT=100
^{84}Kr	-82431.0	2.8		STABLE			0 ⁺	97	IS=57.00.4
$^{84}\text{Kr}^m$	-79195.0	2.8	3236.02 0.18	1.89	μs	0.04	8 ⁺	97	IT=100
^{84}Rb	-79750.0	2.8		32.77	d	0.14	2 ⁻	97	$\beta^+=96.2.5$; $\beta^-=3.8.5$
$^{84}\text{Rb}^m$	-79286.4	2.8	463.62 0.09	20.26	m	0.04	6 ⁻	97	IT \approx 100; $\beta^+=0.0012$
^{84}Sr	-80644	3		STABLE			0 ⁺	97	IS=0.56.1; 2 $\beta^+?$
^{84}Y	-74160	90		4.6	s	0.2	1 ⁺	97	$\beta^+=100$
$^{84}\text{Y}^m$	-74230	170	-80 190	39.5	m	0.8	(5 ⁻)	97	$\beta^+=100$
^{84}Zr	-71490#	200#		25.9	m	0.7	0 ⁺	97	$\beta^+=100$
^{84}Nb	-61880#	300#		9.8	s	0.9	3 ⁺	97 03Do01 T	$\beta^+=100$; $\beta^+p?$
$^{84}\text{Nb}^m$	-61540#	300#	338 10	103	ns	19	(5 ⁻)	00Ch07 ETJ	IT=100
^{84}Mo	-55810#	400#		3.8	ms	0.9	0 ⁺	97 01Ki13 T	$\beta^+=100$; $\beta^+p?$
^{84}Ge	T : average 93Ru01=947(11) 91Kr15=984(23)								
^{84}Nb	T : average 03Do01=9.5(1.0) 77Ko05=12(3)								
^{85}Ga	-40050#	500#		50#	ms (>300 ns)		3/2 ⁻	# 97 97Be70 I	$\beta^-?$
^{85}Ge	-53070#	400#		540	ms	50	5/2 ⁺	# 97	$\beta^-=100$; $\beta^-n=14.3$
^{85}As	-63320#	200#		2.021	s	0.010	3/2 ⁻	# 97	$\beta^-=100$; $\beta^-n=59.4.24$
^{85}Se	-72428	30		31.7	s	0.9	5/2 ⁺	# 97	$\beta^-=100$
^{85}Br	-78610	19		2.90	m	0.06	3/2 ⁻	91	$\beta^-=100$
^{85}Kr	-81480.3	1.9		10.776	y	0.003	9/2 ⁺	91 02Un02 T	$\beta^-=100$
$^{85}\text{Kr}^m$	-81175.4	1.9	304.871 0.020	4.480	h	0.008	1/2 ⁻	91	$\beta^-=78.6.4$; IT=21.4.4
$^{85}\text{Kr}^i$	-79488.5	2.3	1991.8 1.3	1.6	μs	0.7	(17/2 ⁺)	91	IT=100
^{85}Rb	-82167.331	0.011		STABLE			5/2 ⁻	91	IS=72.17.2
^{85}Sr	-81102.6	2.8		64.853	d	0.008	9/2 ⁺	91 02Un02 T	$\epsilon=100$
$^{85}\text{Sr}^m$	-80863.9	2.8	238.66 0.06	67.63	m	0.04	1/2 ⁻	91	IT=86.6.4; $\beta^+=13.4.4$
^{85}Y	-77842	19		2.68	h	0.05	(1/2 ⁻)	94	$\beta^+=100$
$^{85}\text{Y}^m$	-77822	19	19.8 0.5	4.86	h	0.13	9/2 ⁺	94	$\beta^+\approx 100$; IT<0.002
^{85}Zr	-73150	100		7.86	m	0.04	7/2 ⁺	94	$\beta^+=100$
$^{85}\text{Zr}^m$	-72860	100	292.2 0.3	10.9	s	0.3	(1/2 ⁻)	94	IT \leq 92; $\beta^+>8$
^{85}Nb	-67150	220		20.9	s	0.7	(9/2 ⁺)	91	$\beta^+=100$
$^{85}\text{Nb}^m$	-66390	220	759.0 1.0	12	s	5	(1/2 ⁻)	91 98Oi.A ETJ	$\beta^+=100$
^{85}Mo	-59100#	280#		3.2	s	0.2	1/2 ⁻	# 97 97Hu15 TD	$\beta^+=100$; $\beta^+p=?$
^{85}Tc	-47670#	400#		< 110	ns		1/2 ⁻	# 00We.A I	$p?$; $\beta^+?$; $\beta^+p?$
^{85}Tc	I : also 99Ja02<100 ns T : estimated half-life for β^+ decay: 100#ms								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{86}Ga	-34350#	800#	30# ms (>300 ns)		01	97Be70 I	β^- ?
^{86}Ge	-49840#	500#	300# ms (>300 ns)	0^+	01	94Be24 I	β^- ?; β^-_n ?
^{86}As	-59150#	300#	945 ms	8	01		β^- =100; β^-_n =33.4
^{86}Se	-70541	16	15.3 s	0.9	0^+	01	β^- =100
^{86}Br	-75640	11	55.1 s	0.4	(2^-)	01	β^- =100
^{86}Kr	-83265.57	0.10	STABLE		0^+	01	IS=17.30 22; $2\beta^-$?
^{86}Rb	-82747.02	0.20	18.642 d	0.018	2^-	01	$\beta^- \approx 100$; $\epsilon=0.0052$ 5
$^{86}\text{Rb}^m$	-82190.97	0.27	556.05	0.18	6-	01	IT \approx 100; $\beta^- < 0.3$
^{86}Sr	-84523.6	1.1	STABLE		0^+	01	IS=9.86 1
$^{86}\text{Sr}^m$	-81567.9	1.1	2955.68	0.21	455 ns	7	IT=100
^{86}Y	-79284	14	14.74 h	0.02	4-	97	β^+ =100
$^{86}\text{Y}^m$	-79066	14	218.30	0.20	48 m	1	IT=99.31 4; $\beta^+=0.69$ 4
$^{86}\text{Y}^n$	-78982	14	302.2	0.5	125 ns	6	IT=100
^{86}Zr	-77800	30	16.5 h	0.1	0^+	01	β^+ =100
^{86}Nb	-69830	90	88 s	1	(6^+)	01	β^+ =100
$^{86}\text{Nb}^m$	-69580#	180#	250#	160#	* 56 s	8	high 01 94Sh07 JD β^+ =100
^{86}Mo	-64560	440	19.6 s	1.1	0^+	01	β^+ =100
^{86}Tc	-53210#	300#	55 ms	6	(0^+)	01	01Ga24 TJ β^+ =100; β^+p ?
$^{86}\text{Tc}^m$	-51710#	340#	1500	150	1.11 μs	0.21	$(5^+, 5^-)$ 01 00Ch07 EJ IT=100
$^{86}\text{Nb}^m$	I : existence considered as uncertain in ENSDF'01; needs confirmation						
^{86}Tc	T : average 01Ga24=44(12) 01Ki13=59(+8-7)						
$^{86}\text{Tc}^m$	E : above the 4^+ state at 1328 or 1445 keV						
^{87}Ge	-44240#	500#	150# ms (>300 ns)	$5/2^+$	# 02	97Be70 I	β^- ?; β^-_n ?
^{87}As	-55980#	300#	610 ms	120	$3/2^-$	# 02	93Ru01 T β^- =100; β^-_n =15.4 22
^{87}Se	-66580	40	5.50 s	0.12	$5/2^+$	# 02	β^- =100; β^-_n =0.20 4
^{87}Br	-73857	18	55.65 s	0.13	$3/2^-$	02	β^- =100; β^-_n =2.60 4
^{87}Kr	-80709.43	0.27	76.3 m	0.5	$5/2^+$	02	β^- =100
^{87}Rb	-84597.795	0.012	49.23 Gy	0.22	$3/2^-$	02	82Mi14 T IS=27.83 2; β^- =100
^{87}Sr	-84880.4	1.1	STABLE		$9/2^+$	02	IS=7.00 1
$^{87}\text{Sr}^m$	-84491.9	1.1	388.533	0.003	2.815 h	0.012	$1/2^-$ 02 IT \approx 100; $\epsilon=0.30$ 8
^{87}Y	-83018.7	1.6	79.8 h	0.3	$1/2^-$	02	β^+ =100
$^{87}\text{Y}^m$	-82637.9	1.6	380.82	0.07	13.37 h	0.03	$9/2^+$ 02 IT=98.43 10; $\beta^+=1.57$ 10
^{87}Zr	-79348	8	1.68 h	0.01	$(9/2)^+$	02	β^+ =100
$^{87}\text{Zr}^m$	-79012	8	335.84	0.19	14.0 s	0.2	$(1/2)^-$ 02 IT=100
^{87}Nb	-74180	60	3.75 m	0.09	$(1/2^-)$	02	β^+ =100
$^{87}\text{Nb}^m$	-74180	60	3.84	0.14	2.6 m	0.1	$9/2^+$ 02 β^+ =100
^{87}Mo	-67690	220	14.05 s	0.23	$7/2^+$	# 02	97Hu07 TD β^+ =100; β^+p =15.5
^{87}Tc	-59120#	300#	2.18 s	0.16	$1/2^-$	# 02	00We.A TD β^+ =100; β^+p ?
$^{87}\text{Tc}^m$	-59100#	310#	20#	60#	* 2# s		$9/2^+$ 02 β^+ ?; IT ?
^{87}Ru	-47340#	600#	50# ms (>1.5 μs)	$1/2^-$	# 02	95Ry03 I	β^+ ?
^{87}As	T : unweighed average 93Ru01=485(40) 78Cr03=730(60) (Birge ratio $B=3.4$)						
^{87}Rb	T : average 82Mi14=49.44(0.28) 74Ne14=48.8(0.8) 77Da22=48.9(0.4) obtained by						
^{87}Rb	T : three methods, respectively: geochronology, decay counting, chemical						
^{87}Rb	T : 77Da22 supersedes 66Mc12=47.2(0.4) using the same material						
^{87}Mo	T : average 97Hu07=13.6(1.1) 91Mi15=14.5(0.3) 83Ha06=13.3(0.4)						
^{87}Mo	D : average 97Hu07=15(6)% (through 3 levels) 83Ha06=15(8)% first 2^+ state						

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
⁸⁸ Ge	-40140#	700#		80# ms (>300 ns)	0 ⁺	97	97Be70 I	β^- ?
⁸⁸ As	-51290#	500#		300# ms (>300 ns)		97	94Be24 I	β^- ?; β^-_n ?
⁸⁸ Se	-63880	50		1.53 s	0 ⁺	97		β^- =100; β^-_n =0.99 10
⁸⁸ Br	-70730	40		16.36 s	(2 ⁻ , 1 ⁺)	98	93Ru01 T	β^- =100; β^-_n =6.58 18 *
⁸⁸ Br ^m	-70460	40	272.7	5.4 μ s		98		IT=100
⁸⁸ Kr	-79692	13	0.3	2.84 h	0 ⁺	88		β^- =100
⁸⁸ Rb	-82609.00	0.16		17.78 m	2 ⁻	88		β^- =100
⁸⁸ Sr	-87921.7	1.1		STABLE	0 ⁺	88		IS=82.58 1
⁸⁸ Y	-84299.1	1.9		106.65 d	4 ⁻	88		β^+ =100
⁸⁸ Y ^m	-83624.6	1.9	674.55	13.9 ms	(8) ⁺	88		IT=100
⁸⁸ Y ⁿ	-83906.2	1.9	392.86	300 μ s	1 ⁺	88		
⁸⁸ Zr	-83623	10		83.4 d	0 ⁺	88		ϵ =100
⁸⁸ Nb	-76070	100		* 14.5 m	(8 ⁺)	88		β^+ =100
⁸⁸ Nb ^m	-76030	100	40	7.8 m	(4 ⁻)	88		β^+ =100
⁸⁸ Mo	-72700	20		8.0 m	0 ⁺	97		β^+ =100
⁸⁸ Tc	-62710#	200#		* 5.8 s	(2, 3)	97		β^+ =100
⁸⁸ Tc ^m	-62710#	360#	0#	6.4 s	(6, 7, 8)	97		β^+ =100
⁸⁸ Ru	-55650#	400#	300#	1.3 s	0 ⁺	97	01Ki13 TD	β^+ =100; β^+_p ?
* ⁸⁸ Br	T : average 93Ru01=16.34(0.08) 74Gr29=16.5(0.2)				J : systematics prefers (2 ⁻)			**
⁸⁹ Ge	-33690#	900#		50# ms (>300 ns)	3/2 ⁺ #	98	97Be70 I	β^- ?
⁸⁹ As	-47140#	500#		200# ms (>300 ns)	3/2 ⁻ #	98	94Be24 I	β^- ?
⁸⁹ Se	-59200#	300#		410 ms	5/2 ⁺ #	98		β^- =100; β^-_n =7.8 25
⁸⁹ Br	-68570	60		4.40 s	(3/2 ⁻ , 5/2 ⁻)	98		β^- =100; β^-_n =13.8 4 *
⁸⁹ Kr	-76730	50		3.15 m	3/2 ⁽⁺⁾ #	98	95Ke04 J	β^- =100
⁸⁹ Rb	-81713	5		15.15 m	3/2 ⁻	98		β^- =100
⁸⁹ Sr	-86209.1	1.1		50.53 d	5/2 ⁺	98		β^- =100
⁸⁹ Y	-87701.7	2.6		STABLE	1/2 ⁻	98		IS=100.
⁸⁹ Y ^m	-86792.7	2.6	908.97	15.663 s	9/2 ⁺	98	94It.A T	IT=100
⁸⁹ Zr	-84869	4		78.41 h	9/2 ⁺	98		β^+ =100
⁸⁹ Zr ^m	-84281	4	587.82	4.161 m	1/2 ⁻	98		IT=93.77 12; ... *
⁸⁹ Nb	-80650	27		* 2.03 h	(9/2 ⁺)	98		β^+ =100
⁸⁹ Nb ^m	-80650#	40#	0#	1.10 h	(1/2) ⁻	98		β^+ =100
⁸⁹ Mo	-75004	15		2.11 m	(9/2 ⁺)	98		β^+ =100
⁸⁹ Mo ^m	-74617	15	387.5	190 ms	(1/2 ⁻)	98		IT=100
⁸⁹ Tc	-67840#	200#		12.8 s	(9/2 ⁺)	98		β^+ =100
⁸⁹ Tc ^m	-67780#	200#	62.6	12.9 s	(1/2 ⁻)	98		β^+ \approx 100; IT<0.01
⁸⁹ Ru	-59510#	500#		1.38 s	(7/2) ⁽⁺⁾ #	98	00We.A T	β^+ =100; β^+_p ? *
⁸⁹ Rh	-47660#	450#		10# ms (>1.5 μ s)	7/2 ⁺ #	98	95Ry03 I	β^+ ? *
* ⁸⁹ Br	T : ENSDF averages 8 values. Also 93Ru01=4.348(0.022)							**
* ⁸⁹ Zr ^m	D : ... ; β^+ =6.23 12							**
* ⁸⁹ Ru	T : average 00We.A=1.45(0.13) 99Li33=1.2(0.2); same group 01Ki13=1.5(0.2)							**
* ⁸⁹ Rh	I : unobserved in 00We.A, at detection limit							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
⁹⁰ As	-41450# 800#		80# ms (>300 ns)			97Be70 I	β^- ?	
⁹⁰ Se	-55930# 400#		300# ms (>300 ns)	0 ⁺		94Be24 I	β^- ?; β^-n ?	
⁹⁰ Br	-64620 80		1.910 s 0.010		98	93Ru01 T	β^- =100; β^-n =25.2 9 *	
⁹⁰ Kr	-74970 19		32.32 s 0.09	0 ⁺	98		β^- =100	
⁹⁰ Rb	-79362 7		158 s 5	0 ⁻	98		β^- =100	
⁹⁰ Rb ^m	-79255 7	106.90 0.03	258 s 4	3 ⁻	98		β^- =97.4 4; IT=2.6 4	
⁹⁰ Rb ^x	-79291 14	71 12	R = 2 1					
⁹⁰ Sr	-85941.6 2.9		28.79 y 0.06	0 ⁺	98		β^- =100	
⁹⁰ Y	-86487.5 2.6		64.00 h 0.21	2 ⁻	98		β^- =100	
⁹⁰ Y ^m	-85805.8 2.6	681.67 0.10	3.19 h 0.06	7 ⁺	98		IT≈100; β^- =0.0018 2	
⁹⁰ Zr	-88767.3 2.4		STABLE	0 ⁺	98		IS=51.45 40	
⁹⁰ Zr ^m	-86448.3 2.4	2319.000 0.010	809.2 ms 2.0	5 ⁻	98		IT=100	
⁹⁰ Zr ⁿ	-85177.9 2.4	3589.419 0.016	131 ns 4	8 ⁺	98		IT=100	
⁹⁰ Nb	-82656 5		14.60 h 0.05	8 ⁺	98		β^+ =100	
⁹⁰ Nb ^m	-82534 5	122.370 0.022	63 μ s 2	6 ⁺	98		IT=100	
⁹⁰ Nb ⁿ	-82531 5	124.67 0.25	18.81 s 0.06	4 ⁻	98		IT=100	
⁹⁰ Nb ^p	-82485 5	171.10 0.10	< 1 μ s	7 ⁺	98		IT=100	
⁹⁰ Nb ^q	-82274 5	382.01 0.25	6.19 ms 0.08	1 ⁺	98		IT=100	
⁹⁰ Nb ^r	-80776 5	1880.21 0.20	472 ns 13	(11 ⁻)	98		IT=100	
⁹⁰ Mo	-80167 6		5.56 h 0.09	0 ⁺	98		β^+ =100	
⁹⁰ Mo ^m	-77292 6	2874.73 0.15	1.12 μ s 0.05	8 ⁺ #	98		IT=100	
⁹⁰ Tc	-71210 240		* & 8.7 s 0.2	1 ⁺	98		β^+ =100	
⁹⁰ Tc ^m	-70900 300	310 390	BD * & 49.2 s 0.4	(8 ⁺)	98	93Ru03 J	β^+ =100 *	
⁹⁰ Ru	-65310# 300#		11 s 3	0 ⁺	98		β^+ =100	
⁹⁰ Rh	-53220# 500#		* 15 ms 7	0 ⁺ #	98	01Ki13 TD	β^+ =100; β^+p ?	
⁹⁰ Rh ^m	-53220# 710#	0# 500#	* 1.1 s 0.3	9 ⁺ #		01Ki13 TD	β^+ =100; β^+p ?	
* ⁹⁰ Br	T : supersedes 80A115=1.92(0.02) from same group							**
* ⁹⁰ Tc ^m	E : arguments are given in 93Ru03 for the (8 ⁺) level to be the ground-state							**
⁹¹ As	-36860# 900#		50# ms (>300 ns)	3/2 ⁻ #	99	97Be70 I	β^- ?	
⁹¹ Se	-50340# 500#		270 ms 50	1/2 ⁺ #	99		β^- =100; β^-n =21 10	
⁹¹ Br	-61510 70		541 ms 5	3/2 ⁻ #	99		β^- =100; β^-n =20 3	
⁹¹ Kr	-71310 60		8.57 s 0.04	5/2 ⁽⁺⁾	01		β^- =100	
⁹¹ Rb	-77745 8		58.4 s 0.4	3/2 ⁽⁻⁾	99		β^- =100	
⁹¹ Sr	-83645 5		9.63 h 0.05	5/2 ⁺	01		β^- =100	
⁹¹ Sr ^x	-83599 11	47 11	R = 6					
⁹¹ Y	-86345.0 2.9		58.51 d 0.06	1/2 ⁻	99		β^- =100	
⁹¹ Y ^m	-85789.4 2.9	555.58 0.05	49.71 m 0.04	9/2 ⁺	99		IT>98.5; β^- <1.5	
⁹¹ Zr	-87890.4 2.3		STABLE	5/2 ⁺	01		IS=11.22 5	
⁹¹ Zr ^m	-84723.1 2.3	3167.3 0.4	4.35 μ s 0.14	(21/2 ⁺)	01		IT=100	
⁹¹ Nb	-86632 4		680 y 130	9/2 ⁺	99	91Hi.A D	ϵ ≈100; e^+ =0.0138 25	
⁹¹ Nb ^m	-86527 4	104.60 0.05	60.86 d 0.22	1/2 ⁻	99	91Hi.A D	IT=96.6 5; ϵ =3.4 5; ... *	
⁹¹ Nb ⁿ	-84598 4	2034.35 0.19	3.76 μ s 0.12	(17/2 ⁻)	99		IT=100	
⁹¹ Mo	-82204 11		15.49 m 0.01	9/2 ⁺	99		β^+ =100	
⁹¹ Mo ^m	-81551 11	653.01 0.09	64.6 s 0.6	1/2 ⁻	99		IT=50.0 16; β^+ =50.0 16	
⁹¹ Tc	-75980 200		3.14 m 0.02	(9/2 ⁺)	99		β^+ =100	
⁹¹ Tc ^m	-75840 200	139.3 0.3	3.3 m 0.1	(1/2 ⁻)	99		β^+ >99; IT<1	
⁹¹ Ru	-68660# 580#		* 9 s 1	(9/2 ⁺)	99		β^+ =100	
⁹¹ Ru ^m	-68580 500	80# 300#	* 7.6 s 0.8	(1/2 ⁻)	99		β^+ ≈100; β^+p ?; IT ?	
⁹¹ Rh	-59100# 400#		1.74 s 0.14	7/2 ⁺ #	99	00We.A TD	β^+ =100; β^+p ?	
⁹¹ Pd	-47400# 570#		10# ms (>1.5 μ s)	7/2 ⁺ #	99	95Ry03 I	β^+ ?	
* ⁹¹ Nb ^m	D : ... ; e^+ =0.0028 2							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
⁹² As	-30930#	900#	30# ms (>300 ns)		01	97Be70 I	β^- ?	
⁹² Se	-46650#	600#	100# ms (>300 ns)	0 ⁺	01	97Be70 I	β^- ?	
⁹² Br	-56580	50	343 ms 15	(2 ⁻)	01		β^- =100; β^- n=33.1 25	
⁹² Kr	-68785	12	1.840 s 0.008	0 ⁺	01		β^- =100; β^- n=0.0332 25	
⁹² Rb	-74772	6	4.492 s 0.020	0 ⁻	01		β^- =100; β^- n=0.0107 5	
⁹² Sr	-82868	3	2.66 h 0.04	0 ⁺	03		β^- =100	
⁹² Y	-84813	9	3.54 h 0.01	2 ⁻	01		β^- =100	
⁹² Zr	-88453.9	2.3	STABLE	0 ⁺	01		IS=17.15 8	
⁹² Nb	-86448.3	2.8	34.7 My 2.4	(7 ⁺)	01		β^+ ≈100; β^- <0.05	
⁹² Nb ^m	-86312.8	2.8 135.5 0.4	10.15 d 0.02	(2 ⁺)	01		β^+ =100	
⁹² Nb ⁿ	-86222.6	2.8 225.7 0.4	5.9 μs 0.2	(2 ⁻)	01		IT=100	
⁹² Nb ^p	-84245.0	2.8 2203.3 0.4	167 ns 4	(11 ⁻)	01		IT=100	
⁹² Mo	-86805	4	STABLE (>190 Ey)	0 ⁺	01	97Ba35 T	IS=14.84 35; 2 β^+ ?	
⁹² Mo ^m	-84045	4 2760.46 0.16	190 ns 3	8 ⁺	01		IT=100	
⁹² Tc	-78935	26	4.25 m 0.15	(8 ⁺)	01		β^+ =100	
⁹² Tc ^m	-78665	26 270.15 0.11	1.03 μs 0.07	(4 ⁺)	01		IT=100	
⁹² Ru	-74410#	300#	3.65 m 0.05	0 ⁺	01		β^+ =100	
⁹² Rh	-63360#	400#	4.3 s 1.3	(6 ⁺)	01	01Xu05 TJD	β^+ =100; β^+ p=?	
⁹² Pd	-55500#	500#	1.1 s 0.3	0 ⁺	01	01Ki13 TD	β^+ =100; β^+ p ?	
* ⁹² Mo	T : T > 190 Ey (2σ)							**
* ⁹² Rh	T : unweighed average 01Xu05=3.0(0.8) 01Ki13=5.6(0.5) (Birge ratio B=2.76)							**
* ⁹² Rh	J : from 97Ka07; 01Xu05>4							**
⁹³ Se	-40720#	800#	50# ms (>300 ns)	1/2 ⁺ #	97	97Be70 I	β^- ?	
⁹³ Br	-53050#	300#	102 ms 10	3/2 ⁻ #	01		β^- =100; β^- n=68 7	
⁹³ Kr	-64020	100	1.286 s 0.010	1/2 ⁺	01		β^- =100; β^- n=1.95 11	
⁹³ Rb	-72618	8	5.84 s 0.02	5/2 ⁻	97		β^- =100; β^- n=1.39 7	
⁹³ Rb ^m	-72365	8 253.38 0.03	57 μs 15	(3/2 ⁻ , 5/2 ⁻)	97		IT=100	
⁹³ Sr	-80085	8	7.423 m 0.024	5/2 ⁺	97		β^- =100	
⁹³ Y	-84223	11	10.18 h 0.08	1/2 ⁻	97		β^- =100	
⁹³ Y ^m	-83464	11 758.719 0.021	820 ms 40	7/2 ⁺	97		IT=100	
⁹³ Zr	-87117.0	2.3	1.53 My 0.10	5/2 ⁺	97		β^- =100	
⁹³ Nb	-87208.3	2.4	STABLE	9/2 ⁺	97		IS=100.	
⁹³ Nb ^m	-87177.5	2.4 30.77 0.02	16.13 y 0.14	1/2 ⁻	97		IT=100	
⁹³ Mo	-86803	4	4.0 ky 0.8	5/2 ⁺	97		ϵ =100	
⁹³ Mo ^m	-84378	4 2424.89 0.03	6.85 h 0.07	21/2 ⁺	97		IT≈100; β^+ =-0.12 1	
⁹³ Tc	-83603	4	2.75 h 0.05	9/2 ⁺	01		β^+ =100	
⁹³ Tc ^m	-83211	4 391.84 0.08	43.5 m 1.0	1/2 ⁻	01		IT=76.6 11; β^+ =23.4 11	
⁹³ Tc ⁿ	-81418	4 2185.16 0.15	10.2 μs 0.3	(17/2 ⁻)	01			
⁹³ Ru	-77270	90	59.7 s 0.6	(9/2 ⁺)	97		β^+ =100	
⁹³ Ru ^m	-76540	90 734.40 0.10	10.8 s 0.3	(1/2 ⁻)	97	83Ay01 D	β^+ =78.0 23; ...	
⁹³ Ru ⁿ	-75190	90 2082.6 0.9	2.20 μs 0.17	(21/2 ⁺)	97		IT=100	
⁹³ Rh	-69170#	400#	13.9 s 1.6	9/2 ⁺ #	01	01Ki13 TD	β^+ =100; β^+ p ?	
⁹³ Pd	-59700#	400#	1.07 s 0.12	(9/2 ⁺)	01	01Ki13 TJD	β^+ =100; β^+ p=?	
⁹³ Ag	-46780#	600#	5# ms (>1.5 μs)	9/2 ⁺ #	97	95Ry03 I	p ?; β^+ ?	
* ⁹³ Ru ^m	D : ... ; IT=22.0 23; β^+ p=0.027 5							**
* ⁹³ Pd	T : average 01Ki13=1000(200) 01Xu05=1300(200) 00Sc31=900(200)							**
* ⁹³ Ag	I : the few events reported in 94He28 are not trusted by NUBASE							**
* ⁹³ Ag	T : estimated half-life is for β^+ decay; p-decay would be much shorter							**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)	
⁹⁴ Se	-36800#	800#		20#	ms (>300 ns)	0 ⁺	97	97Be70 I	β^- ?	
⁹⁴ Br	-47800#	400#		70	ms	20	92		β^- =100; β^- -n=70 15	
⁹⁴ Kr	-61140#	300#		210	ms	4	0 ⁺	01 03Be05 TD	β^- =100; β^- -n=1.11 7 *	
⁹⁴ Rb	-68553	8		2.702	s	0.005	3 ⁽⁻⁾	92 93Ru01 D	β^- =100; β^- -n=10.01 23	
⁹⁴ Sr	-78840	7		75.3	s	0.2	0 ⁺	92	β^- =100	
⁹⁴ Y	-82348	7		18.7	m	0.1	2 ⁻	92	β^- =100	
⁹⁴ Zr	-87266.8	2.4		STABLE		(>110 Py)	0 ⁺	92 99Ar25 T	IS=17.38 28; 2 β^- ?	
⁹⁴ Nb	-86364.5	2.4		20.3	ky	1.6	(6) ⁺	92	β^- =100	
⁹⁴ Nb ^m	-86323.6	2.4	40.902	0.012	6.263	m	0.004	3 ⁺	92 IT=99.50 6; β^- =0.50 6	
⁹⁴ Mo	-88409.7	1.9		STABLE			0 ⁺	97	IS=9.25 12	
⁹⁴ Tc	-84154	4		293	m	1	7 ⁺	92	β^+ =100	
⁹⁴ Tc ^m	-84079	4	75.5	1.9	52.0	m	1.0	(2) ⁺	92 β^+ ≈100; IT<0.1	
⁹⁴ Ru	-82568	13		51.8	m	0.6	0 ⁺	92	β^+ =100	
⁹⁴ Ru ^m	-79923	13	2644.55	0.25	71	μ s	4	(8) ⁺	92 IT=100	
⁹⁴ Rh	-72940#	450#		*	70.6	s	0.6	(2 ⁺ , 4 ⁺)	92 β^+ =100; β^+ -p=1.8 5	
⁹⁴ Rh ^m	-72640	400	300#	200#	*	25.8	s	0.2	(8) ⁺	92 β^+ =100
⁹⁴ Pd	-66350#	400#		9.0	s	0.5	0 ⁺	02	β^+ =100	
⁹⁴ Pd ^m	-61470#	400#	4884.4	0.5	530	ns	10	(14) ⁺	02 IT=100	
⁹⁴ Ag	-53300#	500#		37	ms	18	0 ⁺ #	02	β^+ =100; β^+ -p ?	
⁹⁴ Ag ^m	-51950#	640#	1350#	400#	422	ms	16	(7) ⁺	02 β^+ =100; β^+ -p=? *	
⁹⁴ Ag ⁿ	-46800#	500#	6500#	2000#	300	ms	200	(21) ⁺	02 β^+ =100; β^+ -p=? *	
* ⁹⁴ Kr	T : average 03Be05=212(5) 72Am01=200(10); others outweighed not used: **									
* ⁹⁴ Kr	T : 03Be05=210(20) 75As04=220(20) and 96Me09=330(100) **									
* ⁹⁴ Ag ^m	T : average 02La18=360(30) 01Ki13=450(20) 94Sc35=420(50) **									
⁹⁵ Br	-43900#	500#		50#	ms (>300 ns)	3/2 ⁻ #	97	97Be70 I	β^- ?	
⁹⁵ Kr	-56040#	400#		114	ms	3	1/2 ⁽⁺⁾	95 03Be05 TD	β^- =100; β^- -n=2.87 18 *	
⁹⁵ Rb	-65854	21		377.5	ms	0.8	5/2 ⁻	95	β^- =100; β^- -n=8.73 20	
⁹⁵ Sr	-75117	7		23.90	s	0.14	1/2 ⁺	94	β^- =100	
⁹⁵ Y	-81207	7		10.3	m	0.1	1/2 ⁻	94	β^- =100	
⁹⁵ Zr	-85657.8	2.4		64.032	d	0.006	5/2 ⁺	00	β^- =100	
⁹⁵ Nb	-86781.9	2.0		34.991	d	0.006	9/2 ⁺	00	β^- =100	
⁹⁵ Nb ^m	-86546.2	2.0	235.690	0.020	3.61	d	0.03	1/2 ⁻	00 IT=94.4 6; β^- =5.6 6	
⁹⁵ Mo	-87707.5	1.9		STABLE			5/2 ⁺	00	IS=15.92 13	
⁹⁵ Tc	-86017	5		20.0	h	0.1	9/2 ⁺	95	β^+ =100	
⁹⁵ Tc ^m	-85978	5	38.89	0.05	61	d	2	1/2 ⁻	95 β^+ =96.12 32; IT=3.88 32	
⁹⁵ Ru	-83450	12		1.643	h	0.014	5/2 ⁺	94	β^+ =100	
⁹⁵ Rh	-78340	150		5.02	m	0.10	(9/2) ⁺	94	β^+ =100	
⁹⁵ Rh ^m	-77800	150	543.3	0.3	1.96	m	0.04	(1/2) ⁻	94 IT=88 5; β^+ =12 5	
⁹⁵ Pd	-70150#	400#		10#	s		9/2 ⁺ #	95 97Sc30 TD	β^+ =100 *	
⁹⁵ Pd ^m	-68290	300	1860#	500#	13.3	s	0.3	(21/2 ⁺)	95 β^+ =?; IT=5#; ... *	
⁹⁵ Ag	-60100#	400#		1.74	s	0.13	(9/2 ⁺)	95 94Sc35 TJD	β^+ =100; β^+ -p=? *	
⁹⁵ Ag ^m	-59760#	400#	344.2	0.3	< 0.5	s	(1/2 ⁻)	03Do.1 ETJ	IT=100	
⁹⁵ Ag ⁿ	-57570#	400#	2531	1	< 16	ms	(23/2 ⁺)	03Do.1 ETJ	IT=100	
⁹⁵ Ag ^p	-55240#	400#	4859	1	< 40	ms	(37/2 ⁺)	03Do.1 ETJ	IT=100	
⁹⁵ Cd	-46700#	600#		5#	ms		9/2 ⁺ #		β^+ ?; β^+ -p ?	
* ⁹⁵ Kr	J : from 95Ke04 **									
* ⁹⁵ Pd	T : 1.35(0.26) s in 97Sc30, if the 1219.3 keV γ originates from ground-state; **									
* ⁹⁵ Pd	T : 1.7 < T < 7.5 s in Schmidt's thesis 1995 cited in 97Sc30t **									
* ⁹⁵ Pd ^m	D : ... ; β^+ -p=0.90 16 **									
* ⁹⁵ Ag	T : from 97Sc30 for β^+ γ activity; supersedes 94Sc35=2.0(0.1) by same authors **									
* ⁹⁵ Ag	T : also 03Do.1=1.85(0.34), same group **									

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
⁹⁶ Br	-38630# 700#				20# ms (>300 ns)		97	97Be70 I	β^- ?
⁹⁶ Kr	-53030# 500#				80 ms 7	0 ⁺	97	03Be05 TD	β^- =100; β^- n=3.7 4
⁹⁶ Rb	-61225 29			*	203 ms 3	2 ⁺	95	93Ru01 D	β^- =100; β^- n=13.4 4 *
⁹⁶ Rb ^m	-61230# 200#	0#	200#	*	200# ms (>1 ms)	1 ^(-#)		81Bo30 JI	β^- ?; IT ?; β^- n ? *
⁹⁶ Sr	-72939 27				1.07 s 0.01	0 ⁺	93		β^- =100
⁹⁶ Y	-78347 23				5.34 s 0.05	0 ⁻	93		β^- =100
⁹⁶ Y ^m	-77206 21	1140	30	BD	9.6 s 0.2	(8) ⁺	93		β^- =100
⁹⁶ Zr	-85442.8 2.8				24 Ey 6	0 ⁺	98	99Ar25 T	IS=2.80 9; 2 β^- =100 *
⁹⁶ Nb	-85604 4				23.35 h 0.05	6 ⁺	93		β^- =100
⁹⁶ Mo	-88790.5 1.9				STABLE	0 ⁺	93		IS=16.68 2
⁹⁶ Tc	-85817 5				4.28 d 0.07	7 ⁺	93		β^+ =100
⁹⁶ Tc ^m	-85783 5	34.28	0.07		51.5 m 1.0	4 ⁺	93		IT=98.0 5; β^+ =2.0 5
⁹⁶ Ru	-86072 8				STABLE (>67 Py)	0 ⁺	01	85No03 T	IS=5.54 14; 2 β^+ ?
⁹⁶ Rh	-79679 13				9.90 m 0.10	(6 ⁺)	93		β^+ =100
⁹⁶ Rh ^m	-79627 13	52.0	0.1		1.51 m 0.02	(3 ⁺)	93		IT=60 5; β^+ =40 5
⁹⁶ Pd	-76230 150				122 s 2	0 ⁺	93		β^+ =100
⁹⁶ Pd ^m	-73700 150	2530.8	0.1		1.81 μ s 0.01	8 ⁺	93	98Gr.B TD	IT=100 *
⁹⁶ Ag	-64570# 400#			*	4.45 s 0.04	(8 ⁺)	93	03Ba39 TJ	β^+ =100; β^+ p=9.7 17 *
⁹⁶ Ag ^m	-64570# 400#	0#	50#	*	6.9 s 0.6	(2 ⁺)		03Ba39 TJD	β^+ =100; β^+ p=18 5
⁹⁶ Ag ⁿ	-64570# 400#				700 ns 200			97Gr02 T	IT ?
⁹⁶ Cd	-56100# 500#				1# s	0 ⁺			β^+ ?
* ⁹⁶ Rb	T : ENSDF average of 8 values. There is also 93Ru01=201(1) **								
* ⁹⁶ Rb ^m	I : non-observation by 81Th04 is not in contradiction with 81Bo30 experiment **								
* ⁹⁶ Rb ^m	I : existence of this isomer is discussed in ENSDF **								
* ⁹⁶ Zr	T : from 21(+8-4 statistics + 2 systematics); other 93Ka12=39(9) in geochemical **								
* ⁹⁶ Zr	T : experiment, not used: observation of 2 β^- decay questioned by 96Ba37 **								
* ⁹⁶ Pd ^m	T : supersedes 97Gr02=1.7(0.1); other 83Gr01=2.2(0.3) outweighed **								
* ⁹⁶ Ag	T : average 03Ba39=4.40(0.06) 97Sc30=4.50(0.06) **								
* ⁹⁶ Ag	D : average β^+ p 97Sc30=11.9(2.6) 82Ku15=8.0(2.3); 96He25=3.7(0.9) not used **								
⁹⁷ Br	-34650# 800#				10# ms (>300 ns)	3/2 ⁻ #	97	97Be70 I	β^- ?
⁹⁷ Kr	-47920# 500#				63 ms 4	3/2 ⁺ #		03Be05 TD	β^- =100; β^- n=6.7 6
⁹⁷ Rb	-58360 30				169.9 ms 0.7	3/2 ⁺	93	93Ru01 D	β^- =100; β^- n=25.7 8
⁹⁷ Sr	-68788 19				429 ms 5	1/2 ⁺	93		β^- =100; β^- n<0.05
⁹⁷ Sr ^m	-68480 19	308.13	0.11		170 ns 10	(7/2) ⁺	93		IT=100
⁹⁷ Sr ⁿ	-67957 19	830.8	0.2		255 ns 10	11/2 ⁻ #	93		IT=100
⁹⁷ Y	-76258 12				3.75 s 0.03	(1/2 ⁻)	93	93Ru01 D	β^- =100; β^- n=0.058 7
⁹⁷ Y ^m	-75590 12	667.51	0.23		1.17 s 0.03	(9/2) ⁺	93		β^- >99.3; IT<0.7; ... *
⁹⁷ Y ⁿ	-72735 12	3523.3	0.4		142 ms 8	(27/2 ⁻)	93		IT \geq 80; β^- \leq 20
⁹⁷ Zr	-82946.6 2.8				16.90 h 0.05	1/2 ⁺	93		β^- =100
⁹⁷ Nb	-85605.6 2.6				72.1 m 0.7	9/2 ⁺	93		β^- =100
⁹⁷ Nb ^m	-84862.3 2.6	743.35	0.03		52.7 s 1.8	1/2 ⁻	93		IT=100
⁹⁷ Mo	-87540.4 1.9				STABLE	5/2 ⁺	93		IS=9.55 8
⁹⁷ Tc	-87220 5				2.6 My 0.4	9/2 ⁺	93		ϵ =100
⁹⁷ Tc ^m	-87123 5	96.56	0.06		90.1 d 1.0	1/2 ⁻	93		IT \approx 100; ϵ <0.34
⁹⁷ Ru	-86112 8				2.9 d 0.1	5/2 ⁺	93		β^+ =100
⁹⁷ Rh	-82590 40				30.7 m 0.6	9/2 ⁺	93		β^+ =100
⁹⁷ Rh ^m	-82330 40	258.85	0.17		46.2 m 1.6	1/2 ⁻	93		β^+ =94.4 6; IT=5.6 6
⁹⁷ Pd	-77800 300				3.10 m 0.09	5/2 ⁺ #	01		β^+ =100
⁹⁷ Ag	-70820 320				25.3 s 0.3	(9/2 ⁺)	93	97Sc30 T	β^+ =100
⁹⁷ Ag ^m	-68480 320	2343	49		5 ns	(21/2 ⁺)			
⁹⁷ Cd	-60600# 400#				2.8 s 0.6	9/2 ⁺ #	93	97Sc30 T	β^+ =100; β^+ p=?
⁹⁷ In	-47000# 600#				5# ms	9/2 ⁺ #			p ?; β^+ ? *
* ⁹⁷ Y ^m	D : ... ; β^- n<0.08 **								
* ⁹⁷ In	T : estimated half-life is for β^+ decay; p-decay would be much shorter **								

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
⁹⁸ Kr	-44800# 600#				46 ms	8	0 ⁺	03	β^- =100; β^- -n=7.0 10
⁹⁸ Rb	-54220				114 ms	5	(0, 1) ^(-#)	03	β^- =100; β^- -n=13.8 6; ... *
⁹⁸ Rb ^m	-53940	120	290	130	96 ms	3	(3, 4) ^(+#)	03	β^- =100
⁹⁸ Sr	-66646	26			653 ms	2	0 ⁺	03	β^- =100; β^- -n=0.25 5
⁹⁸ Y	-72467	25			548 ms	2	(0) ⁻	03	β^- =100; β^- -n=0.331 24
⁹⁸ Y ^m	-72050	30	410	30	2.0 s	0.2	(5 ⁺ , 4 ⁻)	03	β^- =?; IT=10#; ... *
⁹⁸ Y ⁿ	-71971	25	496.19	0.15	7.6 μ s	0.4	(2 ⁻)	03	IT=100
⁹⁸ Y ^p	-72296	25	170.74	0.6	620 ns	80	(2) ⁻	03	IT=100
⁹⁸ Zr	-81287	20			30.7 s	0.4	0 ⁺	03	β^- =100
⁹⁸ Nb	-83529	6			2.86 s	0.06	1 ⁺	03	β^- =100
⁹⁸ Nb ^m	-83445	7	84	4	51.3 m	0.4	(5 ⁺)	03	β^- \approx 100; IT=0.1#
⁹⁸ Mo	-88111.7	1.9			STABLE (>100 Ty)	0 ⁺	03	52Fr23 T	IS=24.13 31; 2 β^- ? *
⁹⁸ Tc	-86428	4			4.2 My	0.3	(6) ⁺	03	β^- =100; β^+ =0
⁹⁸ Tc ^m	-86337	4	90.76	0.16	14.7 μ s	3	(2) ⁻	03	IT=100
⁹⁸ Ru	-88224	6			STABLE	0 ⁺	03		IS=1.87 3
⁹⁸ Rh	-83175	12			* 8.72 m	0.12	(2) ⁺	03	β^+ =100
⁹⁸ Rh ^m	-83120#	50#	60#	50#	* 3.6 m	0.2	(5 ⁺)	03	IT=89 5; β^+ =11 5
⁹⁸ Pd	-81300	21			17.7 m	0.3	0 ⁺	03	β^+ =100
⁹⁸ Ag	-73060	70			47.5 s	0.3	(5 ⁺)	03	ABBW03 J β^+ =100; β^+ -p=0.0012 5 *
⁹⁸ Ag ^m	-72890	70	167.83	0.15	220 ns	20	(3 ⁺)	03	98Gr.B ETD IT=100
⁹⁸ Cd	-67630	80			9.2 s	0.3	0 ⁺	03	β^+ =100; β^+ -p<0.025
⁹⁸ Cd ^m	-65200	80	2427.5	0.6	190 ns	20	8 ⁺ #	98 98Gr.B TD	IT=100 *
⁹⁸ In	-53900# 200#				* 45 ms	23	0 ⁺ #	03	01Ki13 TD β^+ =100; β^+ ?
⁹⁸ In ^m	-53900# 540#	0#	500#		* 1.7 s	0.8		03	01Ki13 TD β^+ =100; β^+ ?
* ⁹⁸ Rb	D : ... ; β^- -2n=0.051 7 **								
* ⁹⁸ Y ^m	D : ... ; β^- -n=3.4 10 **								
* ⁹⁸ Y ^m	J : 94St31=(5 ⁺) 95Ha.B=(4-) **								
* ⁹⁸ Mo	T : limit given here is for 0v-2 β^- decay (theoretically faster, see text) **								
* ⁹⁸ Ag	J : (5 ⁺) with experimental basis preferred to (6 ⁺), see discussion in ENSDF **								
* ⁹⁸ Cd ^m	T : supersedes 97Gr02=200(+300-170); other 97Go18=480(160) outweighed **								
⁹⁹ Kr	-39500# 600#				40 ms	11	3/2 ⁺ #	97 03Be05 TD	β^- =100; β^- -n=11 7
⁹⁹ Rb	-50880	130			50.3 ms	0.7	(5/2 ⁺)	98	β^- =100; β^- -n=15.9 20
⁹⁹ Sr	-62190	80			269 ms	1	3/2 ⁺	95	β^- =100; β^- -n=0.100 19
⁹⁹ Y	-70201	24			1.470 s	0.007	(5/2 ⁺)	95	β^- =100; β^- -n=1.9 4
⁹⁹ Y ^m	-68059	24	2141.65	0.19	8.6 μ s	0.8	(17/2 ⁺)	95	IT=100
⁹⁹ Zr	-77768	20			2.1 s	0.1	1/2 ⁺	95 02Ca37 J	β^- =100
⁹⁹ Nb	-82327	13			15.0 s	0.2	9/2 ⁺	95	β^- =100
⁹⁹ Nb ^m	-81962	13	365.29	0.14	2.6 m	0.2	1/2 ⁻	95	β^- =?; IT<3.8
⁹⁹ Mo	-85965.8	1.9			65.94 h	0.01	1/2 ⁺	95	β^- =100
⁹⁹ Mo ^m	-85868.0	1.9	97.785	0.003	15.5 μ s	0.2	5/2 ⁺	95	IT=100
⁹⁹ Tc	-87323.1	2.0			211.1 ky	1.2	9/2 ⁺	01	β^- =100
⁹⁹ Tc ^m	-87180.4	2.0	142.6832	0.0011	6.015 h	0.009	1/2 ⁻	01	IT \approx 100; β^- =0.0037 6
⁹⁹ Ru	-87617.0	2.0			STABLE		5/2 ⁺	95	IS=12.76 14
⁹⁹ Rh	-85574	7			16.1 d	0.2	(1/2 ⁻)	95	β^+ =100
⁹⁹ Rh ^m	-85510	7	64.3	0.4	4.7 h	0.1	9/2 ⁺	95	β^+ \approx 100; IT<0.16
⁹⁹ Pd	-82188	15			21.4 m	0.2	(5/2 ⁺)	95	β^+ =100
⁹⁹ Ag	-76760	150			124 s	3	(9/2 ⁺)	95	β^+ =100
⁹⁹ Ag ^m	-76250	150	506.1	0.4	10.5 s	0.5	(1/2 ⁻)	95	IT=100
⁹⁹ Cd	-69850# 210#				16 s	3	(5/2 ⁺)	95	β^+ =100; β^+ -p=0.21 8;... *
⁹⁹ In	-61270# 400#				3.1 s	0.8	9/2 ⁺ #	97 01Ki13 TD	β^+ =100; β^+ ?
⁹⁹ In ^m	-60870# 430#	400#	150#		1# s		1/2 ⁻ #		β^+ ?; IT ?
⁹⁹ Sn	-47200# 600#				5# ms		9/2 ⁺ #		β^+ ?; β^+ p ? *
⁹⁹ Sn ^m	-46800# 610#	400#	100#				1/2 ⁻ #		
* ⁹⁹ Cd	D : ... ; β^+ α <1e-4 **								
* ⁹⁹ Sn	I : the 3 events reported in 95Ry03 are not trusted by NUBASE **								

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁰⁰ Kr	-36200#	500#		10# ms (>300 ns)	0 ⁺	97	97Be70 I	β^- ?	
¹⁰⁰ Rb	-46700#	300#		51 ms	8	(3 ⁺)	97 93Ru01 D	β^- =100; β^- -n=5.6 12;... *	
¹⁰⁰ Sr	-60220	130		202 ms	3	0 ⁺	97	β^- =100; β^- -n=0.78 13	
¹⁰⁰ Y	-67290	80		735 ms	7	1 ⁻ , 2 ⁻	97	β^- =100; β^- -n=0.92 8	
¹⁰⁰ Y ^m	-67090#	220#	200#	* 940 ms	30	(3, 4, 5) ⁺ #	97	β^- =100	
¹⁰⁰ Zr	-76600	40		7.1 s	0.4	0 ⁺	97	β^- =100	
¹⁰⁰ Nb	-79939	26		1.5 s	0.2	1 ⁺	97	β^- =100	
¹⁰⁰ Nb ^m	-79471	28	470 40	BD 2.99 s	0.11	(4 ⁺ , 5 ⁺)	97	β^- =100	
¹⁰⁰ Mo	-86184	6		8.5 Ey	0.5	0 ⁺	97 97A102 T	IS=9.63 23; 2 β^- =100 *	
¹⁰⁰ Tc	-86016.2	2.2		15.8 s	0.1	1 ⁺	97	β^- ≈100; ϵ =0.0018 9	
¹⁰⁰ Tc ^m	-85815.5	2.2	200.67 0.04	8.32 μ s	0.14	(4) ⁺	97		
¹⁰⁰ Tc ⁿ	-85772.2	2.2	243.96 0.04	3.2 μ s	0.2	(6) ⁺	97		
¹⁰⁰ Ru	-89219.0	2.0		STABLE		0 ⁺	97	IS=12.60 7	
¹⁰⁰ Rh	-85584	18		20.8 h	0.1	1 ⁻	97	β^+ =100	
¹⁰⁰ Rh ^m	-85476	18	107.6 0.2	4.6 m	0.2	(5 ⁺)	97	IT≈98.3; β^+ ≈1.7	
¹⁰⁰ Pd	-85226	11		3.63 d	0.09	0 ⁺	97	ϵ =100	
¹⁰⁰ Ag	-78150	80		2.01 m	0.09	(5) ⁺	97	β^+ =100	
¹⁰⁰ Ag ^m	-78130	80	15.52 0.16	2.24 m	0.13	(2) ⁺	97	β^+ =?; IT ?	
¹⁰⁰ Cd	-74250	100		49.1 s	0.5	0 ⁺	97	β^+ =100	
¹⁰⁰ Cd ^m	-71700	100	2548.6 0.5	60 ns	3	(8) ⁺	97	IT=100	
¹⁰⁰ In	-64170	250		5.9 s	0.2	(6, 7) ⁺	97 02PI03 TJ	β^+ =100; β^+ p>3.9 *	
¹⁰⁰ Sn	-56780	710		1.1 s	0.4	0 ⁺	97	β^+ =100; β^+ p<17 *	
* ¹⁰⁰ Rb	D : ... ; β^- 2n=0.15 5							**	
* ¹⁰⁰ Rb	T : ENSDF average of 3 values. See also 53(2) of 85Pf.A				J : from 95Pf04			**	
* ¹⁰⁰ Rb	D : β^- 2n intensity is derived from β^- 2n/ β^- n=0.027(7), in 81Jo.A							**	
* ¹⁰⁰ Mo	T : average 97A102=7.6(+2.2-1.4) 97De40=6.82(+0.38-0.53 statistics + 0.68 systematics)							**	
* ¹⁰⁰ Mo	T : 95Da37=9.5(0.9) 91Ej02=11.5(+3-2) and 91El04=11.6(+3.4-0.8)							**	
* ¹⁰⁰ In	T : others: 95Sz01=6.1(0.9) 95Fa.A=6.3(+1.0-9); 95Fa.A supersedes 95Sc33=7.8(.8)							**	
* ¹⁰⁰ Sn	D : from 97Su06 β^+ p/ β^+ <20%							**	
¹⁰¹ Rb	-43600	170		32 ms	4	3/2 ⁺ #	98	β^- =100; β^- -n=28 4	
¹⁰¹ Sr	-55410	120		118 ms	3	(5/2 ⁻)	98	β^- =100; β^- -n=2.37 14	
¹⁰¹ Y	-64910	100		426 ms	20	(5/2 ⁺)	98 96Me09 T	β^- =100; β^- -n=1.94 18 *	
¹⁰¹ Zr	-73460	30		2.3 s	0.1	3/2 ⁺	98 02Ca37 J	β^- =100	
¹⁰¹ Nb	-78942	19		7.1 s	0.3	(5/2#) ⁺	98	β^- =100	
¹⁰¹ Mo	-83511	6		14.61 m	0.03	1/2 ⁺	98	β^- =100	
¹⁰¹ Tc	-86336	24		14.22 m	0.01	9/2 ⁺	98	β^- =100	
¹⁰¹ Tc ^m	-86128	24	207.53 0.04	636 μ s	8	1/2 ⁻	98	IT=100	
¹⁰¹ Ru	-87949.7	2.0		STABLE		5/2 ⁺	98	IS=17.06 2	
¹⁰¹ Ru ^m	-87422.2	2.0	527.5 0.4	17.5 μ s	0.4	11/2 ⁻	98	IT=100	
¹⁰¹ Rh	-87408	17		3.3 y	0.3	1/2 ⁻	98	ϵ =100	
¹⁰¹ Rh ^m	-87251	17	157.32 0.04	4.34 d	0.01	9/2 ⁺	98	ϵ =93.6 2; IT=6.4 2	
¹⁰¹ Pd	-85428	18		8.47 h	0.06	5/2 ⁺	98	β^+ =100	
¹⁰¹ Ag	-81220	100		11.1 m	0.3	9/2 ⁺	98	β^+ =100	
¹⁰¹ Ag ^m	-80950	100	274.1 0.3	3.10 s	0.10	1/2 ⁻	98	IT=100	
¹⁰¹ Cd	-75750	150		1.36 m	0.05	(5/2 ⁺)	98	β^+ =100	
¹⁰¹ In	-68610#	300#		15.1 s	1.1	9/2 ⁺ #	98	β^+ =100; β^+ p=?	
¹⁰¹ In ^m	-68060#	320#	550# 100#	10# s		1/2 ⁻ #		β^+ =95#; IT=5#	
¹⁰¹ Sn	-59560#	300#		3 s	1	5/2 ⁺ #	98	β^+ =100; β^+ p=?	
* ¹⁰¹ Y	T : average 96Me09=400(20) 86Wa17=440(20) and 83Wo10=500(50)							**	
* ¹⁰¹ Y	T : 93Ru01=279(9) at variance, not used							**	

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{102}Rb	-38310#	500#			37 ms	5		98		$\beta^- = 100; \beta^- n = 18.8$
^{102}Sr	-53080	110			69 ms	6	0^+	98	93Ru01 D	$\beta^- = 100; \beta^- n = 5.5, 15$
^{102}Y	-61890	90			* & 300 ms	10	low	98		$\beta^- = 100; \beta^- n = 4.9, 12$
$^{102}\text{Y}^m$	-61690#	220#	200#	200#	* & 360 ms	40	high	98		$\beta^- = 100; \beta^- n = 4.9, 12$
^{102}Zr	-71740	50			2.9 s	0.2	0^+	98		$\beta^- = 100$
^{102}Nb	-76350	40			1.3 s	0.2	1^+	98		$\beta^- = 100$
$^{102}\text{Nb}^m$	-76220	50	130	50	BD 4.3 s	0.4	high	98		$\beta^- = 100$
^{102}Mo	-83557	21			11.3 m	0.2	0^+	01		$\beta^- = 100$
^{102}Tc	-84566	9			* 5.28 s	0.15	1^+	98		$\beta^- = 100$
$^{102}\text{Tc}^m$	-84546	13	20	10	* 4.35 m	0.07	(4,5)	98		$\beta^- = 98.2; IT = 2.2$
^{102}Ru	-89098.0	2.0			STABLE		0^+	98		IS=31.55, 14
^{102}Rh	-86775	5			207.0 d	1.5	($1^-, 2^-$)	98	98Sh21 T	$\beta^+ = 78.5; \beta^- = 22.5$ *
$^{102}\text{Rh}^m$	-86634	5	140.75	0.08	3.742 y	0.010	6^+	98	98Sh21 T	$\beta^+ \approx 100; IT = 0.233, 24$ *
^{102}Pd	-87925.1	3.0			STABLE		0^+	98		IS=1.02, 1; $2\beta^+ ?$
^{102}Ag	-82265	28			12.9 m	0.3	5^+	98		$\beta^+ = 100$
$^{102}\text{Ag}^m$	-82256	28	9.3	0.4	7.7 m	0.5	2^+	98		$\beta^+ = 51.5; IT = 49.5$
^{102}Cd	-79678	29			5.5 m	0.5	0^+	98		$\beta^+ = 100$
^{102}In	-70710	110			23.3 s	0.1	(6^+)	98	03Gi06 T	$\beta^+ = 100; \beta^+ p = 0.0093, 13$ *
^{102}Sn	-64930	130			4.6 s	1.4	0^+	98	95Fa.A T	$\beta^+ = 100; \beta^+ p ?$ *
$^{102}\text{Sn}^m$	-62910	130	2017	2	720 ns	220	(6^+)	98	98Li50 EJT	IT=100 *
* ^{102}Rh	T : average 98Sh21=207.3(1.7) 61Hi06=206(3) **									
* $^{102}\text{Rh}^m$	J : from 99Gi14 **									
* ^{102}In	J : from 95Sz01 **									
* ^{102}Sn	T : 95Fa.A, supersedes 95Sc28=4.5(0.7), preliminary from same group **									
* $^{102}\text{Sn}^m$	T : average 98Li50=620(+430-190) 97Gr02=300(+500-200) 96Li50=1000(500) **									
^{103}Sr	-47550#	500#			50# ms (>300 ns)			01	97Be70 I	$\beta^- ?$
^{103}Y	-58940#	300#			224 ms	19	$5/2^+ \#$	01	96Me09 T	$\beta^- = 100; \beta^- n = 8.3$ *
^{103}Zr	-68370	110			1.3 s	0.1	($5/2^-$)	01		$\beta^- = 100$
^{103}Nb	-75320	70			1.5 s	0.2	($5/2^+$)	01		$\beta^- = 100$
^{103}Mo	-80850	60			67.5 s	1.5	($3/2^+$)	01		$\beta^- = 100$
^{103}Tc	-84597	10			54.2 s	0.8	$5/2^+$	01		$\beta^- = 100$
^{103}Ru	-87258.8	2.0			39.26 d	0.02	$3/2^+$	01		$\beta^- = 100$
$^{103}\text{Ru}^m$	-87020.6	2.1	238.2	0.7	1.69 ms	0.07	$11/2^-$	01		IT=100
^{103}Rh	-88022.2	2.8			STABLE		$1/2^-$	01		IS=100.
$^{103}\text{Rh}^m$	-87982.4	2.8	39.756	0.006	56.114 m	0.009	$7/2^+$	01		IT=100
^{103}Pd	-87479.1	2.9			16.991 d	0.019	$5/2^+$	01		$\epsilon = 100$
$^{103}\text{Pd}^m$	-86694.3	2.9	784.79	0.10	25 ns	2	$11/2^-$	01		IT=100
^{103}Ag	-84791	17			65.7 m	0.7	$7/2^+$	01		$\beta^+ = 100$
$^{103}\text{Ag}^m$	-84657	17	134.45	0.04	5.7 s	0.3	$1/2^-$	01		IT=100
^{103}Cd	-80649	15			7.3 m	0.1	$5/2^+$	01		$\beta^+ = 100$
^{103}In	-74599	25			60 s	1	$9/2^+ \#$	01	97Sz04 T	$\beta^+ = 100$
$^{103}\text{In}^m$	-73967	25	631.7	0.1	34 s	2	$1/2^- \#$	01	97Sz04 ETD	$\beta^+ = 67; IT = 33$
^{103}Sn	-66970#	300#			7 s	3	$5/2^+ \#$	01		$\beta^+ = 100; \beta^+ p = ?$
^{103}Sb	-56180#	300#			100# ms (>1.5 μs)		$5/2^+ \#$	01	95Ry03 I	$\beta^+ ?$
* ^{103}Y	T : average 96Me09=230(20) 96Lh04=190(50) **									

Nuclide	Mass excess (keV)	Excitation energy (keV)				Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{104}Sr	-44400# 700#					30# ms (>300 ns)	0^+	00	97Be70 I	β^- ?
^{104}Y	-54910# 400#					180 ms	60	00	99Wa09 D	β^- =100; β^- -n=?
^{104}Zr	-66340# 400#					1.2 s	0.3	0^+	00	β^- =100
^{104}Nb	-72220 100					4.9 s	0.3	(1^+)	00	β^- =100; β^- -n=0.06 3 *
$^{104}\text{Nb}^m$	-72010 100	220	120		BD *	940 ms	40	high	00	β^- =100; β^- -n=0.05 3
^{104}Mo	-80330 50					60 s	2	0^+	00	β^- =100
^{104}Tc	-82490 50					18.3 m	0.3	3^+ #	00	β^- =100
$^{104}\text{Tc}^m$	-82420 50	69.7	0.2			3.5 μ s	0.3	$2^{(+)}$	00	IT=100
^{104}Ru	-88089 3					STABLE		0^+	00	IS=18.62 27; $2\beta^-$?
^{104}Rh	-86949.8 2.8					42.3 s	0.4	1^+	00	β^- \approx 100; β^+ =0.45 10
$^{104}\text{Rh}^m$	-86820.8 2.8	128.967	0.004			4.34 m	0.03	5^+	00	IT \approx 100; β^- =0.13 1
^{104}Pd	-89390 4					STABLE		0^+	00	IS=11.14 8
^{104}Ag	-85111 6					69.2 m	1.0	5^+	00	β^+ =100
$^{104}\text{Ag}^m$	-85104 6	6.9	0.4			33.5 m	2.0	2^+	00	β^+ \approx 100; IT<0.07
^{104}Cd	-83975 9					57.7 m	1.0	0^+	00	β^+ =100
^{104}In	-76110 80					1.80 m	0.03	$5, 6^{(+)}$	00	β^+ =100
$^{104}\text{In}^m$	-76020 80	93.48	0.10			15.7 s	0.5	(3^+)	00	IT=80; β^+ =20
^{104}Sn	-71590 100					20.8 s	0.5	0^+	00	β^+ =100
^{104}Sb	-59180# 360#					470 ms	130		00	95Fa.A D β^+ =?; β^+ p<7; p<7; α ? *
* ^{104}Nb	D : β^- -n=0.71% of 83En03, at variance, not used **									
* ^{104}Sb	D : 95Fa.A supersedes 95Sc28 p<1 **									
^{105}Sr	-38580# 700#					20# ms (>300 ns)		97	97Be70 I	β^- ?
^{105}Y	-51350# 500#					60# ms (>300 ns)	$5/2^+$ #	97	94Be24 I	β^- ?
^{105}Zr	-62360# 400#					600 ms	100	97		β^- =100; β^- -n ?
^{105}Nb	-70850 100					2.95 s	0.06	$5/2^+$ #	94	96Me09 D β^- =100; β^- -n=1.7 9
^{105}Mo	-77340 70					35.6 s	1.6	$(5/2^-)$	93	β^- =100
^{105}Tc	-82290 60					7.6 m	0.1	$(3/2^-)$	93	β^- =100
^{105}Ru	-85928 3					4.44 h	0.02	$3/2^+$	93	β^- =100
^{105}Rh	-87846 4					35.36 h	0.06	$7/2^+$	93	β^- =100
$^{105}\text{Rh}^m$	-87716 4	129.781	0.004			45 s		$1/2^-$	93	IT=100 *
^{105}Pd	-88413 4					STABLE		$5/2^+$	93	IS=22.33 8
^{105}Ag	-87068 11					41.29 d	0.07	$1/2^-$	93	β^+ =100
$^{105}\text{Ag}^m$	-87043 11	25.465	0.012			7.23 m	0.16	$7/2^+$	93	IT \approx 100; β^+ =0.34 7
^{105}Cd	-84330 12					55.5 m	0.4	$5/2^+$	93	β^+ =100
^{105}In	-79481 17					5.07 m	0.07	$9/2^+$	93	87Eb02 J β^+ =100
$^{105}\text{In}^m$	-78807 17	674.1	0.3			48 s	6	$(1/2^-)$	93	IT=?; β^+ =25#
^{105}Sn	-73260 80					34 s	1	$(5/2^+)$	93	95Pf01 T β^+ =100; β^+ p=? *
^{105}Sb	-63820 100					1.12 s	0.16	$(5/2^+)$	02	β^+ ?; p \approx 1; β^+ p ?
^{105}Te	-52500# 500#					1# μ s		$5/2^+$ #		α ?; β^+ ? *
* $^{105}\text{Rh}^m$	T : no error given; other value: 30 s (see ENSDF: remeasurement recommended) **									
* ^{105}Sn	J : from 85De08 **									
* ^{105}Te	I : the 3 events reported in 95Ry03 are not trusted by NUBASE **									
^{106}Y	-46770# 700#					50# ms (>300 ns)		97	97Be70 I	β^- ?
^{106}Zr	-59700# 500#					200# ms (>300 ns)	0^+	97	94Be24 I	β^- ? *
^{106}Nb	-67100# 200#					920 ms	40	2^+ #	94	96Me09 TD β^- =100; β^- -n=4.5 3 *
^{106}Mo	-76255 18					8.73 s	0.12	0^+	94	95Jo02 T β^- =100
^{106}Tc	-79775 13					35.6 s	0.6	$(1, 2)$	94	β^- =100
^{106}Ru	-86322 8					373.59 d	0.15	0^+	94	β^- =100
^{106}Rh	-86361 8					29.80 s	0.08	1^+	94	β^- =100
$^{106}\text{Rh}^m$	-86225 11	136	12	BD		131 m	2	(6^+)	94	β^- =100
^{106}Pd	-89902 4					STABLE		0^+	94	IS=27.33 3
^{106}Ag	-86937 5					23.96 m	0.04	1^+	94	β^+ =?; β^- \approx 0.5
$^{106}\text{Ag}^m$	-86847 5	89.66	0.07			8.28 d	0.02	6^+	94	β^+ =100; IT \leq 4.2e-6

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
¹⁰⁶ Cd	-87132	6	STABLE	(>410 Ey)	0 ⁺	94 02Tr04 T	IS=1.25 6; 2 β^+ ?	
¹⁰⁶ In	-80606	12	6.2 m	0.1	7 ⁺	94	β^+ =100	
¹⁰⁶ In ^m	-80577	12	28.6	0.3	5.2 m	0.1 (3 ⁺)	94 β^+ =100	
¹⁰⁶ Sn	-77430	50	1.92 m	0.08	0 ⁺	94	β^+ =100	
¹⁰⁶ Sb	-66330#	310#	600 ms	200	(4 ⁺)	97 94Se01 J	β^+ =100 *	
¹⁰⁶ Sb ^m	-65330#	590#	1000#	500#	220 ns	20	98Li50 T IT=100	
¹⁰⁶ Te	-58210	130	70 μ s	20	0 ⁺	94 94Pa11 T	α =100 *	
* ¹⁰⁶ Zr	I : and $T > 240$ ns in 97So07							**
* ¹⁰⁶ Nb	T : average 96Me09=900(20) 83Sh06=1020(50)							**
* ¹⁰⁶ Sb	T : from 95Le.C, Fig. 4, preliminary							**
* ¹⁰⁶ Te	T : average 94Pa11=60(+40-20) 81Sc17=60(+30-10)							**
¹⁰⁷ Y	-42720#	500#	30# ms	(>300 ns)	5/2 ⁺ #	00 97Be70 I	β^- ?	
¹⁰⁷ Zr	-55190#	300#	150# ms	(>300 ns)		00 94Be24 I	β^- ? *	
¹⁰⁷ Nb	-64920#	400#	300 ms	9	5/2 ⁺ #	00 96Me09 TD	β^- =100; β^- n=6.0 15 *	
¹⁰⁷ Mo	-72940	160	3.5 s	0.5	(7/2 ⁻)	00	β^- =100	
¹⁰⁷ Mo ^m	-72870	160	66.3	0.2	470 ns	30 (5/2 ⁻)	00 IT=100	
¹⁰⁷ Tc	-79100	150	21.2 s	0.2	(3/2 ⁻)	00	β^- =100	
¹⁰⁷ Tc ^m	-79030	150	65.7	1.0	184 ns	3 (5/2 ⁻)	00 IT=100	
¹⁰⁷ Ru	-83920	120	3.75 m	0.05	(5/2 ⁺)	00	β^- =100	
¹⁰⁷ Rh	-86863	12	21.7 m	0.4	7/2 ⁺	00	β^- =100	
¹⁰⁷ Rh ^m	-86595	12	268.36	0.04	> 10 μ s	1/2 ⁻	00 IT=100	
¹⁰⁷ Pd	-88368	4	6.5 My	0.3	5/2 ⁺	00	β^- =100	
¹⁰⁷ Pd ^m	-88153	4	214.6	0.3	21.3 s	0.5 (11/2 ⁻)	00 IT=100	
¹⁰⁷ Ag	-88402	4	STABLE		1/2 ⁻	00	IS=51.839 8	
¹⁰⁷ Ag ^m	-88309	4	93.125	0.019	44.3 s	0.2 (7/2 ⁺)	00 IT=100	
¹⁰⁷ Cd	-86985	6	6.50 h	0.02	5/2 ⁺	00	β^+ =100	
¹⁰⁷ In	-83560	11	32.4 m	0.3	9/2 ⁺	00	β^+ =100	
¹⁰⁷ In ^m	-82882	11	678.5	0.3	50.4 s	0.6 (1/2 ⁻)	00 IT=100	
¹⁰⁷ Sn	-78580	80	2.90 m	0.05	(5/2 ⁺)	00	β^+ =100	
¹⁰⁷ Sb	-70650#	300#	4.6 s	0.8	5/2 ⁺ #	00	β^+ =100	
¹⁰⁷ Te	-60540#	300#	3.1 ms	0.1	5/2 ⁺ #	00	α =70 30; β^+ =30 30	
* ¹⁰⁷ Zr	I : and $T > 240$ ns in 97So07							**
* ¹⁰⁷ Nb	T : average 96Me09=300(30) 91Hi02=300(10)							**
¹⁰⁸ Y	-37740#	800#	20# ms	(>300 ns)		00 95Cz.A I	β^- ?; β^- n ?	
¹⁰⁸ Zr	-52200#	600#	80# ms	(>300 ns)	0 ⁺	00 97Be70 I	β^- ?; β^- n ?	
¹⁰⁸ Nb	-60700#	300#	193 ms	17	(2 ⁺)	00	β^- =100; β^- n=6.2 5	
¹⁰⁸ Mo	-71300#	200#	1.09 s	0.02	0 ⁺	00	β^- =100	
¹⁰⁸ Tc	-75950	130	5.17 s	0.07	(2 ⁺)	00	β^- =100	
¹⁰⁸ Ru	-83670	120	4.55 m	0.05	0 ⁺	00	β^- =100	
¹⁰⁸ Rh	-85020	110	16.8 s	0.5	1 ⁺	00	β^- =100	
¹⁰⁸ Rh ^m	-85080	40	-60	110	BD *	6.0 m	0.3 (5) ^(+#) 00 β^- =100	
¹⁰⁸ Pd	-89524	3	STABLE		0 ⁺	00	IS=26.46 9	
¹⁰⁸ Ag	-87602	4	2.37 m	0.01	1 ⁺	00	β^- =97.15 20; β^+ =2.85 20	
¹⁰⁸ Ag ^m	-87493	4	109.440	0.007	418 y	21	6 ⁺ 00 β^+ =91.3 9; IT=8.7 9 *	
¹⁰⁸ Cd	-89252	6	STABLE	(>410 Py)	0 ⁺	02 95Ge14 T	IS=0.89 3; 2 β^+ ?	
¹⁰⁸ In	-84116	10	58.0 m	1.2	7 ⁺	00	β^+ =100	
¹⁰⁸ In ^m	-84086	10	29.75	0.05	39.6 m	0.7	2 ⁺ 00 β^+ =100	
¹⁰⁸ Sn	-82041	20	10.30 m	0.08	0 ⁺	00	β^+ =100	
¹⁰⁸ Sb	-72510#	210#	7.4 s	0.3	(4 ⁺)	00	β^+ =100; β^+ p ?	
¹⁰⁸ Te	-65720	100	2.1 s	0.1	0 ⁺	00 85Ti02 D	β^+ =51 4; α =49 4; ... *	
¹⁰⁸ I	-52650#	360#	36 ms	6	1 ⁺ #	00 94Pa12 D	α =?; β^+ =9#; p<1 *	
* ¹⁰⁸ Ag ^m	T : discrepant results: 418(7) 310(130) 127(21), see ENSDF							**
* ¹⁰⁸ Te	D : ... ; β^+ p=2.4 10; β^+ α <0.065							**
* ¹⁰⁸ I	D : β^+ =9%# estimated by 94Pa12 using theoretical β^+ half-life \approx 400 ms							**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{109}Zr	-47280#	500#		60#	ms (>300 ns)		99	97Be70 I	β^- ?
^{109}Nb	-58100#	500#		190	ms	30	99		β^- =100; β^- -n=31 5
^{109}Mo	-67250#	300#		530	ms	60	99		β^- =100
^{109}Tc	-74540	100		860	ms	40	99		β^- =100; β^- -n=0.08 2
^{109}Ru	-80850	70		34.5	s	1.0	99		β^- =100
^{109}Rh	-85011	12		80	s	2	99		β^- =100
^{109}Pd	-87607	3		13.7012	h	0.0024	99		β^- =100
$^{109}\text{Pd}^m$	-87418	3	188.990	0.010	4.696	m	0.003		IT=100
^{109}Ag	-88722.7	2.9		STABLE			99		IS=48.161 8
$^{109}\text{Ag}^m$	-88634.7	2.9	88.0341	0.0011	39.6	s	0.2		IT=100
^{109}Cd	-88508	4		461.4	d	1.2	99		ϵ =100
$^{109}\text{Cd}^m$	-88448	4	59.6	0.4	12	μs	2		IT=100
$^{109}\text{Cd}^n$	-88045	4	463.0	0.5	10.9	μs	0.5		IT=100
^{109}In	-86489	6		4.2	h	0.1	99		β^+ =100
$^{109}\text{In}^m$	-85839	6	650.1	0.3	1.34	m	0.07		IT=100
$^{109}\text{In}^n$	-84387	6	2101.8	0.2	209	ms	6	(19/2 ⁺)	IT=100
^{109}Sn	-82639	10		18.0	m	0.2	99		β^+ =100
^{109}Sb	-76259	19		17.0	s	0.7	99		β^+ =100
^{109}Te	-67610	60		4.6	s	0.3	99		β^+ =?; α =3.9 13; ... *
^{109}I	-57610	100		103	μs	5	(5/2 ⁺)	02 87Gi02 J	p=100
* ^{109}Te	D: ...; β^+ p=9.4 31; β^+ α <0.005 **								
^{110}Zr	-43900#	800#		30#	ms (>300 ns)		00	97Be70 I	β^- ?
^{110}Nb	-53620#	500#		170	ms	20	00		β^- =100; β^- -n=40 8
^{110}Mo	-65460#	400#		300	ms	40	00		β^- =100; β^- -n ?
^{110}Tc	-70960	80		920	ms	30	(2 ⁺)	00 96Me09 D	β^- =100; β^- -n=0.04 2
^{110}Ru	-79980	50		11.6	s	0.6	00		β^- =100
^{110}Rh	-82780	50		* 28.5	s	1.5	(> 3) ⁽⁺⁾	00	β^- =100
$^{110}\text{Rh}^m$	-82839	22	-60	50	BD *	3.2	s	0.2	1 ⁺ 00
^{110}Pd	-88349	11		STABLE			(>600 Py)	00	52Wi26 T
^{110}Ag	-87460.6	2.9		24.6	s	0.2	1 ⁺	00	IS=11.72 9; 2 β^- ?
$^{110}\text{Ag}^m$	-87343.0	2.9	117.59	0.05	249.950	d	0.024	6 ⁺	00 β^- \approx 100; ϵ =0.30 6
^{110}Cd	-90353.0	2.7		STABLE				00	β^- =98.64 6; IT=1.36 6
^{110}In	-86475	12		4.9	h	0.1	7 ⁺	00	IS=12.49 18
$^{110}\text{In}^m$	-86413	12	62.1	0.5	69.1	m	0.5	2 ⁺	00 β^+ =100
^{110}Sn	-85844	14		4.11	h	0.10	0 ⁺	00	β^+ =100
^{110}Sb	-77540#	200#		23.0	s	0.4	(4 ⁺)	00 97La13 J	β^+ =100
^{110}Te	-72280	50		18.6	s	0.8	0 ⁺	00	β^+ \approx 100; α =0.003#
^{110}I	-60320#	310#		650	ms	20	1 ⁺ #	00	β^+ =83 4; α =17 4; ... *
^{110}Xe	-51900	130		310	ms	190	0 ⁺	00 02Ma19 TD	α =64 35; β^+ ?
* ^{110}I	D: ...; β^+ p=11 3; β^+ α =1.1 3 **								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹¹¹ Nb	-50630# 500#		80# ms (>300 ns)	5/2 ⁺ #	97	97Be70 I	β^- ?	
¹¹¹ Mo	-61100# 400#		200# ms (>300 ns)		97	94Be24 I	β^- ? *	
¹¹¹ Tc	-69220 110		290 ms 20	3/2 ⁻ #	96	96Me09 TD	β^- =100; β^- n=0.85 20 *	
¹¹¹ Ru	-76670 70		2.12 s 0.07	(5/2 ⁺)	96	98Lh02 J	β^- =100	
¹¹¹ Rh	-82357 30		11 s 1	(7/2 ⁺)	96		β^- =100	
¹¹¹ Pd	-86004 11		23.4 m 0.2	5/2 ⁺	96		β^- =100	
¹¹¹ Pd ^m	-85832 11	172.18 0.08	5.5 h 0.1	11/2 ⁻	96		IT=73.3; β^- =27.3	
¹¹¹ Ag	-88221 3		7.45 d 0.01	1/2 ⁻	96		β^- =100	
¹¹¹ Ag ^m	-88161 3	59.82 0.04	64.8 s 0.8	7/2 ⁺	96		IT=99.3 2; β^- =0.7 2	
¹¹¹ Cd	-89257.5 2.7		STABLE	1/2 ⁺	00		IS=12.80 12	
¹¹¹ Cd ^m	-88861.3 2.7	396.214 0.021	48.50 m 0.09	11/2 ⁻	00		IT=100	
¹¹¹ In	-88396 5		2.8047 d 0.0004	9/2 ⁺	00		ϵ =100	
¹¹¹ In ^m	-87859 5	536.95 0.06	7.7 m 0.2	1/2 ⁻	00		IT=100	
¹¹¹ Sn	-85945 7		35.3 m 0.6	7/2 ⁺	96		β^+ =100	
¹¹¹ Sn ^m	-85690 7	254.72 0.08	12.5 μ s 1.0	1/2 ⁺				
¹¹¹ Sb	-80888 28		75 s 1	(5/2 ⁺)	96		β^+ =100	
¹¹¹ Te	-73480 70		19.3 s 0.4	5/2 ⁺ #	97		β^+ =100; β^+ p=?	
¹¹¹ I	-64950# 300#		2.5 s 0.2	5/2 ⁺ #	96		β^+ \approx 100; α =0.088	
¹¹¹ I ^m	-63550# 300#	1398 1	21 ns 2	(11/2 ⁻)				
¹¹¹ Xe	-54400# 300#		740 ms 200	5/2 ⁺ #	96	94Pa11 D	β^+ ?; α =10.7	
¹¹¹ Xe ^m		non existent	RN 900 ms 200			90Tu.A T		
* ¹¹¹ Mo	I : and T > 240 ns in 97So07							**
* ¹¹¹ Tc	T : supersedes 88Pe13=300(30) from same group							**
* ¹¹¹ Xe ^m	I : from assigning α decay to isomer in older version of ENSDF							**
¹¹² Nb	-45800# 700#		60# ms (>300 ns)	2 ⁺ #	97	97Be70 I	β^- ?	
¹¹² Mo	-58830# 600#		150# ms (>300 ns)	0 ⁺	97	94Be24 I	β^- ?	
¹¹² Tc	-66000 120		290 ms 20	2 ⁺ #	97	99Wa09 TD	β^- =100; β^- n=1.5 2	
¹¹² Ru	-75480 70		1.75 s 0.07	0 ⁺	97		β^- =100	
¹¹² Rh	-79740 50		3.4 s 0.4	1 ⁺	97	99Lh01 T	β^- =100 *	
¹¹² Rh ^m	-79410 60	330 70	BD 6.73 s 0.15	> 3	97	99Lh01 T	β^- =100 *	
¹¹² Pd	-86336 18		21.03 h 0.05	0 ⁺	97		β^- =100	
¹¹² Ag	-86624 17		3.130 h 0.009	2 ⁽⁻⁾	97		β^- =100	
¹¹² Cd	-90580.5 2.7		STABLE	0 ⁺	97		IS=24.13 21	
¹¹² In	-87996 5		14.97 m 0.10	1 ⁺	97		β^+ =56.3; β^- =44.3	
¹¹² In ^m	-87839 5	156.59 0.05	20.56 m 0.06	4 ⁺	97		IT=100	
¹¹² In ⁿ	-87645 5	350.76 0.09	690 ns 50	7 ⁺	97		IT=100	
¹¹² In ^p	-87382 5	613.69 0.14	2.81 μ s 0.03	8 ⁻	97	87Eb02 J	IT=100	
¹¹² Sn	-88661 4		STABLE	0 ⁺	97		IS=0.97 1; 2 β^+ ?	
¹¹² Sb	-81601 18		51.4 s 1.0	3 ⁺	97		β^+ =100	
¹¹² Te	-77300 170		2.0 m 0.2	0 ⁺	97		β^+ =100	
¹¹² I	-67100# 210#		3.42 s 0.11	1 ⁺ #	97	78Ro19 D	β^+ \approx 100; α =0.0012; ... *	
¹¹² Xe	-59970 100		2.7 s 0.8	0 ⁺	97	94Pa11 D	β^+ \approx 100; α =0.9 8 *	
¹¹² Cs	-46290# 300#		500 μ s 100	1 ⁺ #	02		p=100	
* ¹¹² Rh	T : supersedes 91Jo11=2.1(0.3) and 88Ay02=3.8(0.6) of same group							**
* ¹¹² Rh ^m	T : supersedes 88Ay02=6.8(0.2)							**
* ¹¹² I	D : ... ; β^+ p=0.88 10; β^+ α =0.104 12							**
* ¹¹² I	D : β^+ p and β^+ α are derived from β^+ p/ α =735(80) β^+ p/ β^+ α =8.5(2), in 85Ti02							**
* ¹¹² Xe	D : α intensity is estimated from 94Pa11=0.8(+1.1-0.5)% and 78Ro19=0.84%							**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{113}Nb	-42200# 800#			30# ms (>300 ns)	$5/2^+$ #	98	97Be70	I	β^- ?
^{113}Mo	-54140# 600#			100# ms (>300 ns)		98	94Be24	I	β^- ?
^{113}Tc	-63720# 300#			170 ms 20	$3/2^-$ #	98	99Wa09	TD	β^- =100; β^- n=2.1 3
^{113}Ru	-72200 70			800 ms 50	$(5/2^+)$	98	98Ku17	J	β^- =100
$^{113}\text{Ru}^m$	-72070 70	130	18	510 ms 30	$(11/2^-)$		98Ku17	ETJ	IT=?; β^- =?
^{113}Rh	-78680 50			2.80 s 0.12	$(7/2^+)$	98	93Pe11	J	β^- =100
^{113}Pd	-83690 40			93 s 5	$(5/2^+)$	98			β^- =100
$^{113}\text{Pd}^m$	-83610 40	81.1	0.3	300 ms 100	$(9/2^-)$	98			IT=100
$^{113}\text{Pd}^n$		non existent	RN	> 100 s			98	81Me17	I
^{113}Ag	-87033 17			5.37 h 0.05	$1/2^-$	98			β^- =100
$^{113}\text{Ag}^m$	-86990 17	43.50	0.10	68.7 s 1.6	$7/2^+$	98			IT=64 7; β^- =36 7
^{113}Cd	-89049.3 2.7			7.7 Py 0.3	$1/2^+$	98			IS=12.22 12; β^- =100
$^{113}\text{Cd}^m$	-88785.8 2.7	263.54	0.03	14.1 y 0.5	$11/2^-$	98			β^- ≈100; IT=0.14
^{113}In	-89370 3			STABLE	$9/2^+$	99			IS=4.29 5
$^{113}\text{In}^m$	-88978 3	391.699	0.003	1.6579 h 0.0004	$1/2^-$	99			IT=100
^{113}Sn	-88333 4			115.09 d 0.03	$1/2^+$	00			β^+ =100
$^{113}\text{Sn}^m$	-88256 4	77.386	0.019	21.4 m 0.4	$7/2^+$	00			IT=91.1 23; β^+ =8.9 23
^{113}Sb	-84420 18			6.67 m 0.07	$5/2^+$	98			β^+ =100
^{113}Te	-78347 28			1.7 m 0.2	$(7/2^+)$	98			β^+ =100
^{113}I	-71130 50			6.6 s 0.2	$5/2^+$ #	98			β^+ =100; α =3.31e-7; ...
^{113}Xe	-62090 80			2.74 s 0.08	$5/2^+$ #	98	85Ti02	D	β^+ ≈100; α =0.011 5; ...
^{113}Cs	-51700 100			16.7 μ s 0.7	$5/2^+$ #	02			p=100; α =0
* ^{113}Tc	T : 98Ku17=110(30) and 92Ay02=130(50) are from same authors								
* $^{113}\text{Ru}^m$	E : above the 99 keV level and below 160 keV								
* $^{113}\text{Pd}^n$	I : existence is not possible since discovery of $^{113}\text{Pd}^m$ by 93Pe11								
* ^{113}I	D : ... ; $\beta^+ \alpha$?								
* ^{113}Xe	D : ... ; $\beta^+ \text{p}=7 4$; $\beta^+ \alpha \approx 0.007 4$								
* ^{113}Xe	D : $\alpha=0.0024-0.0204\%$ from estimated limit for the reduced width, see 85Ti02								
* ^{113}Xe	D : $\beta^+ \text{p}$ and $\beta^+ \alpha$ derived from $\beta^+ \text{p}/\alpha=605(35)$ and $\beta^+ \text{p}/\beta^+ \alpha=500-1500$ in 85Ti02								
^{114}Mo	-51310# 700#			80# ms (>300 ns)	0^+	03	97Be70	I	β^- ?
^{114}Tc	-59730# 600#			150 ms 30	2^+ #	03			β^- =100; β^- n=?
^{114}Ru	-70530# 230#			530 ms 60	0^+	03			β^- =100; β^- n ?
^{114}Rh	-75630 110			1.85 s 0.05	1^+	03			β^- =100; β^- n ?
$^{114}\text{Rh}^m$	-75430# 190#	200#	150#	1.85 s 0.05	(4,5)	03			β^- =100
^{114}Pd	-83497 24			2.42 m 0.06	0^+	03			β^- =100
^{114}Ag	-84949 25			4.6 s 0.1	1^+	03			β^- =100
$^{114}\text{Ag}^m$	-84750 25	199	5	1.50 ms 0.05	($< 7^+$)	03			IT=100
^{114}Cd	-90020.9 2.7			STABLE	($> 92 \text{ Py}$)	03	95Ge14	T	IS=28.73 42; $2\beta^-$?
^{114}In	-88572 3			71.9 s 0.1	1^+	03			β^- =99.50 15; β^+ =0.50 15
$^{114}\text{In}^m$	-88382 3	190.29	0.03	49.51 d 0.01	5^+	03			IT=96.75 24; β^+ =3.25 24
$^{114}\text{In}^n$	-88070 3	501.94	0.03	43.1 ms 0.6	(8^-)	03			IT=100
$^{114}\text{In}^p$	-87930 3	641.72	0.03	4.3 μ s 0.4	(7^+)	03			IT=100
^{114}Sn	-90561 3			STABLE	0^+	03			IS=0.66 1
$^{114}\text{Sn}^m$	-87474 3	3087.37	0.07	733 ns 14	7^-	03			IT=100
^{114}Sb	-84515 28			3.49 m 0.03	(3^+)	03			β^+ =100
$^{114}\text{Sb}^m$	-84020 28	495.5	0.07	219 μ s 12	(8^-)	03			IT=100
^{114}Te	-81889 28			15.2 m 0.7	0^+	03			β^+ =100
^{114}I	-72800# 300#			2.1 s 0.2	1^+	03			β^+ =100; $\beta^+ \text{p}$?
$^{114}\text{I}^m$	-72530# 300#	265.9	0.5	6.2 s 0.5	(7)	03	ABBW96	D	β^+ =91 2; IT=9 2
^{114}Xe	-67086 11			10.0 s 0.4	0^+	03			β^+ =100
^{114}Cs	-54540# 310#			570 ms 20	(1^+)	03			β^+ ≈100; α =0.018 6; ...
^{114}Ba	-45950 140			530 ms 230	0^+	03	02Ma19	D	β^+ ≈100; $\beta^+ \text{p}=20 10$; ...
* $^{114}\text{I}^m$	D : evaluated for NUBASE by J. Blachot, based on ^{114}I IT decay								
* ^{114}Cs	D : ... ; $\beta^+ \text{p}=8.7 13$; $\beta^+ \alpha=0.19 3$								
* ^{114}Ba	D : ... ; $\alpha=0.9 3$; $^{12}\text{C}<0.038$								
* ^{114}Ba	D : ^{12}C intensity is from 95Gu10								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)			
^{115}Mo	-46310#	800#	60#	ms (>300 ns)		99	$\beta^- ?; \beta^- n ?$			
^{115}Tc	-57110#	700#	100#	ms (>300 ns)	3/2 ⁻	99	$\beta^- ?; \beta^- n ?$			
^{115}Ru	-66430	130	740	ms	80	99	$\beta^- =100; \beta^- n ?$			
^{115}Rh	-74210	80	990	ms	50	7/2 ⁺ #	$\beta^- =100$			
^{115}Pd	-80400	60	25	s	2	5/2 ⁺ #	$\beta^- =100$			
$^{115}\text{Pd}^m$	-80310	60	89.18	0.25	50	s 3	11/2 ⁻ #	$\beta^- =92.0$ 20; IT=8.0 20		
^{115}Ag	-84990	30			20.0	m 0.5	1/2 ⁻	$\beta^- =100$		
$^{115}\text{Ag}^m$	-84950	30	41.16	0.10	18.0	s 0.7	7/2 ⁺	$\beta^- =79.0$ 3; IT=21.0 3		
^{115}Cd	-88090.5	2.7			53.46	h 0.10	1/2 ⁺	$\beta^- =100$		
$^{115}\text{Cd}^m$	-87909.5	2.7	181.0	0.5	44.56	d 0.24	(11/2 ⁻)	$\beta^- \approx 100; IT < 0.003$		
^{115}In	-89537	4			441	Ty 25	9/2 ⁺	IS=95.71 5; $\beta^- =100$		
$^{115}\text{In}^m$	-89201	4	336.244	0.017	4.486	h 0.004	1/2 ⁻	IT=95.0 7; $\beta^- =5.0$ 7		
^{115}Sn	-90036.0	2.9			STABLE		1/2 ⁺	IS=0.34 1		
$^{115}\text{Sn}^m$	-89423.2	2.9	612.81	0.04	3.26	μs 0.08	7/2 ⁺	IT=100		
$^{115}\text{Sn}^n$	-89322.4	2.9	713.64	0.12	159	μs 1	11/2 ⁻	IT=100		
^{115}Sb	-87003	16			32.1	m 0.3	5/2 ⁺	$\beta^+ =100$		
^{115}Te	-82063	28			5.8	m 0.2	7/2 ⁺	$\beta^+ =100$		
$^{115}\text{Te}^m$	-82053	29	10	7	6.7	m 0.4	(1/2 ⁺)	$\beta^+ \approx 100; IT < 0.06$		
$^{115}\text{Te}^n$	-81783	28	280.05	0.20	7.5	μs 0.2	11/2 ⁻	IT=100		
^{115}I	-76338	29			1.3	m 0.2	5/2 ⁺ #	$\beta^+ =100$		
^{115}Xe	-68657	12			18	s 4	(5/2 ⁺)	$\beta^+ =100; \beta^+ p = 0.34$ 6; ...		
^{115}Cs	-59700#	300#			1.4	s 0.8	9/2 ⁺ #	$\beta^+ =100; \beta^+ p \approx 0.07$		
^{115}Ba	-49030#	600#			450	ms 50	5/2 ⁺ #	$\beta^+ =100; \beta^+ p > 15$		
* $^{115}\text{Pd}^m$	J : E3 transition to ground-state							**		
* $^{115}\text{Te}^m$	E : less than 20 keV, from ENSDF							**		
* ^{115}Xe	D : ... ; $\beta^+ \alpha = 0.0003$ 1							**		
^{116}Tc	-52750#	700#			90#	ms (>300 ns)	2 ⁺ #	01 97Be70 I	$\beta^- ?$	
^{116}Ru	-64450#	700#			400#	ms (>300 ns)	0 ⁺	01 94Be24 I	$\beta^- ?$	
^{116}Rh	-70740	140			680	ms 60	1 ⁺	01	$\beta^- =100; \beta^- n ?$	
$^{116}\text{Rh}^m$	-70540#	210#	200#	150#	570	ms 50	(6 ⁻)	01	$\beta^- =100$	
^{116}Pd	-79960	60			11.8	s 0.4	0 ⁺	01	$\beta^- =100$	
^{116}Ag	-82570	50			2.68	m 0.10	(2) ⁻	01	$\beta^- =100$	
$^{116}\text{Ag}^m$	-82490	50	81.90	0.20	8.6	s 0.3	(5 ⁺)	01	$\beta^- =94.0$ 15; IT=6.0 15	
^{116}Cd	-88719	3			30	Ey 4	0 ⁺	01	03Da09 T	IS=7.49 18; 2 $\beta^- =100$
^{116}In	-88250	4			14.10	s 0.03	1 ⁺	01	98Bh04 D	$\beta^- \approx 100; \epsilon = 0.23$ 6
$^{116}\text{In}^m$	-88123	4	127.267	0.006	54.29	m 0.17	5 ⁺	01		$\beta^- =100$
$^{116}\text{In}^n$	-87960	4	289.660	0.006	2.18	s 0.04	8 ⁻	01		IT=100
^{116}Sn	-91528.1	2.9			STABLE		0 ⁺	01		IS=14.54 9
^{116}Sb	-86821	6			15.8	m 0.8	3 ⁺	01		$\beta^+ =100$
$^{116}\text{Sb}^m$	-86440	40	380	40	BD	60.3	m 0.6	8 ⁻	01	$\beta^+ =100$
^{116}Te	-85269	28			2.49	h 0.04	0 ⁺	01		$\beta^+ =100$
^{116}I	-77490	100			2.91	s 0.15	1 ⁺	01		$\beta^+ =100$
$^{116}\text{I}^m$	-77090#	110#	400#	50#	3.27	μs 0.16	(7 ⁻)	01		IT=100
^{116}Xe	-73047	13			59	s 2	0 ⁺	01		$\beta^+ =100$
^{116}Cs	-62070#	100#			700	ms 40	(1 ⁺)	01		$\beta^+ =100; \beta^+ p = 0.28$ 7; ...
$^{116}\text{Cs}^m$	-61970#	120#	100#	60#	3.85	s 0.13	4 ⁺ , 5, 6	01		$\beta^+ =100; \beta^+ p = 0.51$ 15; ...
^{116}Ba	-54600#	400#			1.3	s 0.2	0 ⁺	01		$\beta^+ =100; \beta^+ p = 3$ 1
* ^{116}Ru	I : and $T > 240$ ns in 97So07							**		
* ^{116}Cd	T : from 29(1 statistics +4-3 systematics); supersedes 00Da27=26(1 statistics +7-4 systematics)							**		
* ^{116}Cs	D : ... ; $\beta^+ \alpha = 0.049$ 25							**		
* $^{116}\text{Cs}^m$	D : ... ; $\beta^+ \alpha = 0.008$ 2							**		

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{117}Tc	-49850#	700#	40# ms (>300 ns)	$3/2^-$	#	02 97Be70 I	β^- ?	
^{117}Ru	-60010#	700#	300# ms (>300 ns)			02 94Be24 I	β^- ? *	
^{117}Rh	-68950#	500#	440 ms	40		02	β^- =100	
^{117}Pd	-76530	60	4.3 s	0.3		02	β^- =100	
$^{117}\text{Pd}^m$	-76330	60	203.2	0.3		02	IT=100	
^{117}Ag	-82270	50	73.6 s	1.4		02	β^- =100	
$^{117}\text{Ag}^m$	-82240	50	28.6	0.2		02	β^- =94.0 15; IT=6.0 15	
^{117}Cd	-86425	3	2.49 h	0.04		02	β^- =100	
$^{117}\text{Cd}^m$	-86289	3	136.4	0.2		02	β^- ≈100; IT≈0	
^{117}In	-88945	6	43.2 m	0.3		02	β^- =100	
$^{117}\text{In}^m$	-88630	6	315.302	0.012		02	β^- =52.9 15; IT=47.1 15	
^{117}Sn	-90400.0	2.9	STABLE			02	IS=7.68 7	
$^{117}\text{Sn}^m$	-90085.4	2.9	314.58	0.04		02	IT=100	
^{117}Sb	-88645	9	2.80 h	0.01		02	β^+ =100	
^{117}Te	-85097	13	62 m	2		02	β^+ =100; e^+ =25 1	
$^{117}\text{Te}^m$	-84801	13	296.1	0.5		02	IT ?	
$^{117}\text{Te}^n$	-84823	13	274.4	0.1		02	IT=100	
^{117}I	-80435	28	2.22 m	0.04		02	β^+ =100; e^+ ≈77	
^{117}Xe	-74185	10	61 s	2		02	β^+ =100; β^+ p=0.0029 6	
^{117}Cs	-66440	60	* 8.4 s	0.6		02	β^+ =100	
$^{117}\text{Cs}^m$	-66290#	100#	150#	80#	*	02	β^+ =100	
$^{117}\text{Cs}^x$	-66390	80	50	50			$R=?$	
^{117}Ba	-57290#	300#	1.75 s	0.07		02 97Ja12 D	β^+ =100; β^+ p=13 3; ... *	
^{117}La	-46510#	400#	23.5 ms	2.6		02	$p=?$; β^+ =6#	
$^{117}\text{La}^m$	-46370#	400#	138	15	p	02	$p=?$; β^+ =3#	
* ^{117}Ru	I : and $T > 240$ ns in 97So07							**
* ^{117}Ba	D : ... ; β^+ α =0.024 8							**
* ^{117}Ba	D : β^+ p from 97Ja12. β^+ p/ β^+ α =350-1200 from 85Ti02 yields β^+ α =0.011-0.037							**
^{118}Tc	-45200#	900#	30# ms (>300 ns)	2^+	#	97 95Cz.A I	β^- ?	
^{118}Ru	-57920#	800#	200# ms (>300 ns)	0^+		94Be24 I	β^- ?	
^{118}Rh	-65140#	500#	310 ms	30		97 00Jo18 TJD	β^- =100	
^{118}Pd	-75470	210	1.9 s	0.1		95	β^- =100	
^{118}Ag	-79570	60	3.76 s	0.15		95 93Ja03 J	β^- =100	
$^{118}\text{Ag}^m$	-79440	60	127.49	0.05		95 95Ap.A E	β^- =59; IT=41	
^{118}Cd	-86709	20	50.3 m	0.2		95	β^- =100	
^{118}In	-87230	8	* 5.0 s	0.5		95	β^- =100	
$^{118}\text{In}^m$	-87130#	50#	100#	50#	*	95 94It.A T	β^- =100	
$^{118}\text{In}^n$	-86990#	50#	240#	50#	*	95	IT=98.6 3; β^- =1.4 3 *	
^{118}Sn	-91656.1	2.9	STABLE			95	IS=24.22 9	
^{118}Sb	-87999	4	3.6 m	0.1		95	β^+ =100	
$^{118}\text{Sb}^m$	-87749	6	250	6	BD	95	β^+ =100	
$^{118}\text{Sb}^n$	-87948	4	50.814	0.021		95	β^+ =100	
^{118}Te	-87721	15	6.00 d	0.02		95	ϵ =100	
^{118}I	-80971	20	13.7 m	0.5		95	β^+ =100	
$^{118}\text{I}^m$	-80781	20	190.1	1.0		95 94Ka39 E	β^+ ≈100; IT=?	
^{118}Xe	-78079	10	3.8 m	0.9		95	β^+ =100	
^{118}Cs	-68409	13	* 14 s	2		95	β^+ =100; β^+ p=0.021 14;... *	
$^{118}\text{Cs}^m$	-68310#	60#	100#	60#	*	95 93Be46 J	β^+ =100; β^+ p=0.021 14;... *	
$^{118}\text{Cs}^x$	-68404	12	5	4			$R < 0.1$	
^{118}Ba	-62370#	200#	5.2 s	0.2		97 97Ja12 TD	β^+ =100; β^+ p ?	
^{118}La	-49620#	300#	200# ms				β^+ ?	
* $^{118}\text{In}^n$	E : 138.2(0.5) keV above $^{118}\text{In}^m$, from ENSDF							**
* ^{118}Cs	D : ... ; β^+ α =0.0012 5							**
* ^{118}Cs	D : derived from β^+ p=0.042(6)%, β^+ α =0.0024(4)% for mixture of ground-state and isomer.							**
* ^{118}Cs	D : Replaced by uniform distributions from zero to values for each isomer							**
* $^{118}\text{Cs}^m$	D : ... ; β^+ α =0.0012 5							**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹¹⁹ Ru	-53240#	700#		170#	ms (>300 ns)		97Be70 I	β^- ?
¹¹⁹ Rh	-63240#	600#		300#	ms (>300 ns)	7/2 ⁺ #	94Be24 I	β^- ?
¹¹⁹ Pd	-71620#	300#		920	ms	130	00	β^- =100
¹¹⁹ Ag	-78560	90		6.0	s	0.5	1/2 ⁻ #	β^- =100
¹¹⁹ Ag ^m	-78540#	90#	20#	2.1	s	0.1	7/2 ⁺ #	β^- =100
¹¹⁹ Cd	-83910	80		2.69	m	0.02	(3/2 ⁺)	β^- =100
¹¹⁹ Cd ^m	-83760	80	146.54	2.20	m	0.02	11/2 ⁻ #	β^- =100
¹¹⁹ In	-87704	8		2.4	m	0.1	9/2 ⁺	β^- =100
¹¹⁹ In ^m	-87393	8	311.37	18.0	m	0.3	1/2 ⁻	β^- =94.4 15; IT=5.6 15
¹¹⁹ Sn	-90068.4	2.9		STABLE			1/2 ⁺	IS=8.59 4
¹¹⁹ Sn ^m	-89978.9	2.9	89.531	293.1	d	0.7	11/2 ⁻	IT=100
¹¹⁹ Sb	-89477	8		38.19	h	0.22	5/2 ⁺	ϵ =100
¹¹⁹ Sb ^m	-86625	11	2852	850	ms	90	27/2 ⁺ #	IT=100
¹¹⁹ Te	-87184	8		16.05	h	0.05	1/2 ⁺	β^+ =100
¹¹⁹ Te ^m	-86923	8	260.96	4.70	d	0.04	11/2 ⁻	ϵ =99.59 4; e^+ =0.41 4; ...
¹¹⁹ I	-83766	28		19.1	m	0.4	5/2 ⁺	β^+ =100
¹¹⁹ Xe	-78794	10		5.8	m	0.3	5/2 ⁽⁺⁾	90Ne.A J e^+ =79 5; ϵ =21 5
¹¹⁹ Cs	-72305	14		43.0	s	0.2	9/2 ⁺	β^+ =100; $\beta^+\alpha < 2e-6$
¹¹⁹ Cs ^m	-72260#	30#	50#	30.4	s	0.1	3/2 ⁽⁺⁾	β^+ =100
¹¹⁹ Cs ^s	-72289	9	16	R = .5 .25		spmix		
¹¹⁹ Ba	-64590	200		5.4	s	0.3	(5/2 ⁺)	β^+ =100; $\beta^+p < 25$
¹¹⁹ La	-54970#	400#		1#	s		11/2 ⁻ #	β^+ ?
¹¹⁹ Ce	-44000#	600#		200#	ms		5/2 ⁺ #	β^+ ?
* ¹¹⁹ Ag ^m	E : estimated from 7/2 ⁺ level in isotopes ¹¹³ Ag=43 ¹¹⁵ Ag=41 ¹¹⁷ Ag=28							
* ¹¹⁹ Sb ^m	E : estimated less than 20 keV above 2841.7 level							
* ¹¹⁹ Te ^m	D : ... ; IT<0.008							
¹²⁰ Ru	-50940#	800#		80#	ms (>300 ns)	0 ⁺	02 95Cz.A I	β^- ?
¹²⁰ Rh	-59230#	600#		200#	ms (>300 ns)		94Be24 I	β^- ?
¹²⁰ Pd	-70150	120		500	ms	100	0 ⁺	β^- =100
¹²⁰ Ag	-75650	70		1.23	s	0.04	3 ⁽⁺⁾ #	02 93Ru01 D β^- =100; $\beta^-n < 0.003$
¹²⁰ Ag ^m	-75450	70	203.0	371	ms	24	6 ⁽⁻⁾	02 03Wa13 T $\beta^- \approx 63$; IT ≈ 37
¹²⁰ Cd	-83974	19		50.80	s	0.21	0 ⁺	β^- =100
¹²⁰ In	-85740	40		3.08	s	0.08	1 ⁺	β^- =100
¹²⁰ In ^m	-85690#	50#	50#	46.2	s	0.8	5 ⁺	02 87Eb02 J β^- =100
¹²⁰ In ⁿ	-85440#	200#	300#	47.3	s	0.5	8 ⁽⁻⁾	02 79Fo10 J β^- =100
¹²⁰ Sn	-91105.1	2.5		STABLE			0 ⁺	IS=32.58 9
¹²⁰ Sn ^m	-88623.5	2.5	2481.63	11.8	μ s	0.5	(7 ⁻)	02 IT=100
¹²⁰ Sn ⁿ	-88202.9	2.5	2902.22	6.26	μ s	0.11	10 ⁺ #	02 IT=100
¹²⁰ Sb	-88424	8		15.89	m	0.04	1 ⁺	β^+ =100
¹²⁰ Sb ^m	-88420#	100#	0#	5.76	d	0.02	8 ⁻	β^+ =100
¹²⁰ Sb ⁿ	-88346	8	78.16	246	ns	2	(3 ⁺)	02 IT=100
¹²⁰ Sb ^p	-86096	8	2328.3	400	ns	8	(6)	02 IT=100
¹²⁰ Te	-89405	10		STABLE			0 ⁺	IS=0.09 1; 2 β^+ ?
¹²⁰ I	-83790	18		81.6	m	0.2	2 ⁻	β^+ =100
¹²⁰ I ^m	-83717	18	72.61	228	ns	15	(1 ⁺ , 2 ⁺ , 3 ⁺)	02 IT=100
¹²⁰ I ⁿ	-83470	23	320	53	m	4	(7 ⁻)	02 β^+ =100
¹²⁰ Xe	-82172	12		40	m	1	0 ⁺	β^+ =100
¹²⁰ Cs	-73889	10		61.2	s	1.8	2 ⁽⁻⁾ #	02 β^+ =100; $\beta^+\alpha < 2.0e-5$ 4; ...
¹²⁰ Cs ^m	-73790#	60#	100#	57	s	6	(7 ⁻)	02 75Ho09 D β^+ =100; $\beta^+\alpha < 2.0e-5$ 4; ...
¹²⁰ Cs ^s	-73884	9	5	R < 0.1		spmix		
¹²⁰ Ba	-68890	300		24	s	2	0 ⁺	02 92Xu04 T β^+ =100
¹²⁰ La	-57690#	500#		2.8	s	0.2	02	β^+ =100; β^+p =?
¹²⁰ Ce	-49710#	700#		250#	ms		0 ⁺	β^+ ?
* ¹²⁰ Ag ^m	T : average 03Wa13=400(30) 71Fo22=320(40)							
* ¹²⁰ Cs	D : ... ; $\beta^+p < 7e-6$ 3							
* ¹²⁰ Cs	D : isomers not distinguished by 75Ho09 in $\beta^+\alpha$ and β^+p . Values replaced							
* ¹²⁰ Cs	D : ... by upper limits for both (cf. ENSDF evaluation of ¹¹⁸ Cs)							
* ¹²⁰ Cs ^m	D : ... ; $\beta^+p < 7e-6$ 3							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)		
¹²¹ Rh	-57080#	900#	100# ms (>300 ns)	7/2 ⁺ #		94Be24 I	β^- ?		
¹²¹ Pd	-66260#	500#	400# ms (>300 ns)		00	94Be24 I	β^- ? *		
¹²¹ Ag	-74660	150	790 ms	20	7/2 ⁺ #	00	β^- =100; β^- n=0.080 13		
¹²¹ Cd	-81060	80	13.5 s	0.3	(3/2 ⁺)	00	β^- =100		
¹²¹ Cd ^m	-80850	80	214.86	0.15	8.3 s	0.8	(11/2 ⁻) 00	β^- =100	
¹²¹ In	-85841	27	23.1 s	0.6	9/2 ⁺	00	β^- =100		
¹²¹ In ^m	-85528	27	312.98	0.08	3.88 m	0.10	1/2 ⁻ 00	β^- =98.8 2; IT=1.2 2	
¹²¹ Sn	-89204.1	2.5	27.03 h	0.04	3/2 ⁺	00	β^- =100		
¹²¹ Sn ^m	-89197.8	2.5	6.30	0.06	43.9 y	0.5	11/2 ⁻ 00	IT=77.6 20; β^- =22.4 20	
¹²¹ Sn ⁿ	-87205.3	2.7	1998.8	0.9	5.3 μ s	0.5	19/2 ⁺ # 00	IT=100	
¹²¹ Sb	-89595.1	2.2			STABLE		5/2 ⁺ 00	IS=57.21 5	
¹²¹ Te	-88551	26	19.16 d	0.05	1/2 ⁺	00	β^+ =100		
¹²¹ Te ^m	-88257	26	293.991	0.022	154 d	7	11/2 ⁻ 00	IT=88.6 11; β^+ =11.4 11	
¹²¹ I	-86287	10	2.12 h	0.01	5/2 ⁺	00	β^+ =100		
¹²¹ I ^m	-83910	10	2376.9	0.4	9.0 μ s	1.5	00	IT=100	
¹²¹ Xe	-82473	11	40.1 m	2.0	(5/2 ⁺)	00	β^+ =100		
¹²¹ Cs	-77100	14	155 s	4	3/2 ⁽⁺⁾	00	β^+ =100		
¹²¹ Cs ^m	-77032	14	68.5	0.3	122 s	3	9/2 ⁽⁺⁾ 00	β^+ =83; IT=17	
¹²¹ Ba	-70740	140	29.7 s	1.5	5/2 ⁽⁺⁾	00	β^+ =100; β^+ p=0.02 1		
¹²¹ La	-62400#	500#	5.3 s	0.2	11/2 ⁻ #	00	β^+ =100; β^+ p ?		
¹²¹ Ce	-52700#	500#	1.1 s	0.1	(5/2) ⁽⁺⁾ #	00	99Li46 J β^+ =100; β^+ p \approx 1		
¹²¹ Pr	-41580#	700#	600 ms	300	(3/2 ⁻)	00	90Bo39 TJD p=?; β^+ ?; β^+ p ? *		
* ¹²¹ Pd	I : and T>240 ns in 97So07 **								
* ¹²¹ Pr	T : T=1.4(0.8) s in ENSDF: not trusted to belong to this nuclide **								
¹²² Rh	-52900#	700#	50# ms (>300 ns)			97Be70 I	β^- ?		
¹²² Pd	-64690#	400#	300# ms (>300 ns)		0 ⁺	98 94Be24 I	β^- ? *		
¹²² Ag	-71230#	210#	* 520 ms	14	(3 ⁺)	94 95Fe12 T	β^- =100; β^- n=0.186 10 *		
¹²² Ag ^m	-71150#	220#	* 1.5 s	0.5	8 ⁻ #	94	β^- =100; β^- n ?		
¹²² Cd	-80730	40	* 5.24 s	0.03	0 ⁺	94	β^- =100		
¹²² In	-83580	50	* 1.5 s	0.3	1 ⁺	94	β^- =100		
¹²² In ^m	-83540#	80#	* 10.3 s	0.6	5 ⁺	94	β^- =100		
¹²² In ⁿ	-83290	130	290	140	BD	10.8 s	0.4	8 ⁻ 94	β^- =100
¹²² Sn	-89945.9	2.7			STABLE		0 ⁺ 94	IS=4.63 3; 2 β^- ?	
¹²² Sb	-88330.2	2.2	2.7238 d	0.0002	2 ⁻	94	β^- =97.59 12; ... *		
¹²² Sb ^m	-88166.6	2.2	163.5591	0.0017	4.191 m	0.003	(8) ⁻ 94	IT=100	
¹²² Sb ⁿ	-88192.7	2.2	137.472	0.001	530 μ s		5 ⁺		
¹²² Te	-90314.0	1.5			STABLE		0 ⁺ 94	IS=2.55 12	
¹²² I	-86080	5	3.63 m	0.06	1 ⁺	94	β^+ =100		
¹²² Xe	-85355	11	20.1 h	0.1	0 ⁺	94	ϵ =100		
¹²² Cs	-78140	30	21.18 s	0.19	1 ⁺	96 93Al03 T	β^+ =100; β^+ α <2e-7 *		
¹²² Cs ^m	-78005	9	140	30	MD	3.70 m	0.11	8 ⁻ 96	β^+ =100
¹²² Cs ⁿ	-78010	30	127.0	0.5	360 ms	20	(5) ⁻ 96	IT=100	
¹²² Ba	-74609	28	1.95 m	0.15	0 ⁺	94	β^+ =100		
¹²² La	-64540#	300#	8.7 s	0.7		94	β^+ =100; β^+ p=?		
¹²² Ce	-57840#	400#	2# s		0 ⁺	94	β^+ ?; β^+ p ? *		
¹²² Pr	-44890#	500#	500# ms				β^+ ?		
* ¹²² Pd	I : and T>240 ns in 97So07 **								
* ¹²² Ag	D : β^- n intensity is from 93Ru01 **								
* ¹²² Sb	D : ... ; β^+ =2.41 12 **								
* ¹²² Cs	T : average 93Al03=21.2(0.2) 69Ch18=21.0(0.7) **								
* ¹²² Cs	D : β^+ α intensity upper limit is from 75Ho09 **								
* ¹²² Ce	I : T=8.7(0.7) s in NDS 71 (1994) was misprint for ¹²² La; corrected in ENSDF **								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹²³ Pd	-60610# 600#		200# ms (>300 ns)			94Be24 I	β^- ?
¹²³ Ag	-69960# 210#		296 ms	6	(7/2 ⁺)	94 95Fe12 T	β^- =100; β^- -n=0.55 5 *
¹²³ Cd	-77310 40		2.10 s	0.02	(3/2 ⁺)	94	β^- =100
¹²³ Cd ^m	-76990 40	316.52 0.23	1.82 s	0.03	(11/2 ⁻)	94	β^- =?; IT=?
¹²³ In	-83426 24		5.98 s	0.06	9/2 ⁺	94	β^- =100
¹²³ In ^m	-83099 24	327.21 0.04	47.8 s	0.5	1/2 ⁻	94	β^- =100
¹²³ Sn	-87820.5 2.7		129.2 d	0.4	11/2 ⁻	94	β^- =100
¹²³ Sn ^m	-87795.9 2.7	24.6 0.4	40.06 m	0.01	3/2 ⁺	94	β^- =100
¹²³ Sb	-89224.1 2.1		STABLE		7/2 ⁺	94	IS=42.79 5
¹²³ Te	-89171.9 1.5		> 600 Ty		1/2 ⁺	94 96Al30 T	IS=0.89 3; ϵ =100 *
¹²³ Te ^m	-88924.3 1.5	247.55 0.04	119.25 d	0.15	11/2 ⁻	94	IT=100
¹²³ I	-87943 4		13.2235 h	0.0019	5/2 ⁺	94 02Un02 T	β^+ =100
¹²³ Xe	-85249 10		2.08 h	0.02	1/2 ⁺	94 90Ne.A J	β^+ =100
¹²³ Xe ^m	-85064 10	185.18 0.22	5.49 μ s	0.26	7/2 ⁽⁻⁾		
¹²³ Cs	-81044 12		5.87 m	0.04	1/2 ⁺	94 93Al03 T	β^+ =100 *
¹²³ Cs ^m	-80887 12	156.74 0.21	1.64 s	0.12	(11/2 ⁻)	94	IT=100
¹²³ Cs ^x	-81037 13	7 4	$R < 0.1$		spmix		
¹²³ Ba	-75655 12		2.7 m	0.4	5/2 ⁺	94	β^+ =100
¹²³ La	-68710# 200#		17 s	3	11/2 ⁻ #	94	β^+ =100
¹²³ Ce	-60180# 300#		3.8 s	0.2	(5/2) ⁽⁺⁾ #	94	β^+ =100; β^+ p=?
¹²³ Pr	-50340# 600#		800# ms		3/2 ⁺ #		β^+ ?
* ¹²³ Ag	T : average 95Fe12=293(7) 86Ma42=300(20) 83Re05=300(10)				D : from 93Ru01		**
* ¹²³ Te	T : and T=24(9) Ey for ϵ (K), same authors						
* ¹²³ Te	I : this nuclide is not considered 'stable' since K ϵ has been observed						
* ¹²³ Cs	T : average 93Al03=5.87(0.05) 68Ch18=5.87(0.05)						
¹²⁴ Pd	-58800# 500#		100# ms (>300 ns)	0 ⁺		97Be70 I	β^- ?
¹²⁴ Ag	-66470# 200#		* 172 ms	5	3 ⁺ #	97	β^- =100; β^- -n>0.1
¹²⁴ Ag ^m	-66470# 220#	0# 100#	* 200# ms		8 ⁻ #	95Kr.A I	β^- ?; IT ? *
¹²⁴ Cd	-76710 60		1.25 s	0.02	0 ⁺	97	β^- =100
¹²⁴ In	-80880 50		* 3.11 s	0.10	3 ⁺	97	β^- =100
¹²⁴ In ^m	-80900 50	-20 70	BD * 3.7 s	0.2	(8) ⁽⁻⁾ #	97	β^- \approx 100; IT ?
¹²⁴ Sn	-88236.8 1.4		STABLE		0 ⁺	97 52Ka41 T	IS=5.79 5; 2 β^- ?
¹²⁴ Sn ^m	-85911.8 1.4	2325.01 0.04	3.1 μ s	0.5	7 ⁻	97	IT=100
¹²⁴ Sn ⁿ	-85580.2 1.5	2656.6 0.5	45 μ s	5	10 ⁺ #	97	IT=100
¹²⁴ Sb	-87620.3 2.1		60.20 d	0.03	3 ⁻	98	β^- =100
¹²⁴ Sb ^m	-87609.4 2.1	10.8627 0.0008	93 s	5	5 ⁺	97	IT=75 5; β^- =25 5
¹²⁴ Sb ⁿ	-87583.5 2.1	36.8440 0.0014	20.2 m	0.2	(8) ⁻	97	IT=100
¹²⁴ Sb ^p	-87579.5 2.1	40.8038 0.0007	3.2 μ s	0.3	(3 ⁺ , 4 ⁺)	97	IT=100
¹²⁴ Te	-90524.5 1.5		STABLE		0 ⁺	97	IS=4.74 14
¹²⁴ I	-87365.0 2.4		4.1760 d	0.0003	2 ⁻	97	β^+ =100
¹²⁴ Xe	-87660.1 1.8		STABLE		0 ⁺	97 89Ba22 T	IS=0.09 1; 2 β^+ ?
¹²⁴ Cs	-81731 8		30.9 s	0.4	1 ⁺	97 93Al03 T	β^+ =100 *
¹²⁴ Cs ^m	-81268 8	462.55 0.17	6.3 s	0.2	(7) ⁺	97	IT=100
¹²⁴ Cs ^x	-81701 22	30 20	$R=?$		spmix		
¹²⁴ Ba	-79090 12		11.0 m	0.5	0 ⁺	97	β^+ =100
¹²⁴ La	-70260 60		* 29.21 s	0.17	(7 ⁻ , 8 ⁻)	97 97As05 T	β^+ =100 *
¹²⁴ La ^m	-70160# 120#	100# 100#	* 21 s	4	low ⁽⁺⁾ #	97 97As05 T	β^+ =100
¹²⁴ Ce	-64820# 300#		9.1 s	1.2	0 ⁺	98 97As05 T	β^+ =100 *
¹²⁴ Pr	-53130# 600#		1.2 s	0.2		97	β^+ =100; β^+ p=?
¹²⁴ Nd	-44500# 600#		500# ms		0 ⁺		β^+ ?
* ¹²⁴ Ag ^m	I : "There is some evidence for a low-spin and a high-spin isomer in ¹²⁴ Ag"						
* ¹²⁴ Cs	T : average 93Al03=30.9(0.5) 78Ek05=30.8(0.5)						
* ¹²⁴ La	J : for ¹²⁴ La and ¹²⁴ La ^m are from 92Id01						
* ¹²⁴ Ce	T : average 97As05=10.8(1.5) 78Bo32=6(2)						

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		J^π	Ens	Reference	Decay modes and intensities (%)	
¹²⁵ Ag	-64800#	300#			166	ms	7	7/2 ⁺ #	99	β^- =100; β^- n=?	
¹²⁵ Cd	-73360	70			* 650	ms	20	3/2 ⁺ #	99	β^- =100	
¹²⁵ Cd ^m	-73310	50	50	70	BD *	570	ms	90	11/2 ⁻ #	99 89Hu03 T	β^- =100
¹²⁵ In	-80480	30			2.36	s	0.04	9/2 ⁺	99	β^- =100	
¹²⁵ In ^m	-80120	30	360.12	0.09	12.2	s	0.2	1/2 ⁽⁻⁾	99	β^- =100	
¹²⁵ Sn	-85898.5	1.5			9.64	d	0.03	11/2 ⁻	99	β^- =100	
¹²⁵ Sn ^m	-85871.0	1.5	27.50	0.14	9.52	m	0.05	3/2 ⁺	99	β^- =100	
¹²⁵ Sb	-88255.5	2.6			2.75856	y	0.00025	7/2 ⁺	99	β^- =100	
¹²⁵ Te	-89022.2	1.5			STABLE			1/2 ⁺	99	IS=7.07 15	
¹²⁵ Te ^m	-88877.4	1.5	144.772	0.009	57.40	d	0.15	11/2 ⁻	99	IT=100	
¹²⁵ I	-88836.4	1.5			59.400	d	0.010	5/2 ⁺	99	ϵ =100	
¹²⁵ Xe	-87192.1	1.9			16.9	h	0.2	1/2 ⁽⁺⁾	99	β^+ =100	
¹²⁵ Xe ^m	-86939.5	1.9	252.60	0.14	56.9	s	0.9	9/2 ⁽⁻⁾	99	IT=100	
¹²⁵ Cs	-84088	8			45	m	1	1/2 ⁽⁺⁾	99	β^+ =100	
¹²⁵ Cs ^m	-83821	8	266.6	1.1	900	ms	30	(11/2 ⁻)	99 98Su16 TJ	IT=100	
¹²⁵ Ba	-79668	11			3.5	m	0.4	1/2 ⁽⁺⁾ #	99	β^+ =100	
¹²⁵ La	-73759	26			64.8	s	1.2	(11/2 ⁻)	99	β^+ =100	
¹²⁵ La ^m	-73652	26	107.0	0.1	390	ms	40	(3/2 ⁺)	99 99Ca21 ETJ	IT=100	
¹²⁵ Ce	-66660#	200#			9.3	s	0.3	(7/2 ⁻)	99 02Pe15 J	β^+ =100; β^+ p=?	
¹²⁵ Pr	-57910#	400#			3.3	s	0.7	3/2 ⁺ #	02	β^+ =100; β^+ p?	
¹²⁵ Nd	-47620#	400#			600	ms	150	5/2 ⁽⁺⁾ #	02	β^+ =100	
* ¹²⁵ Cd ^m	T : unweighed average 89Hu03=480(30) 86Ma42=660(30) (Birge ratio $B=4.24$)										
* ¹²⁵ La	J : ENSDF'99 says ground-state spin unknown; a (11/2 ⁻) level lies at 8-9 keV above ground-state										
* ¹²⁵ La ^m	J : 3/2 ⁺ # from systematics; low spin and even-parity from 99Ca21										
* ¹²⁵ Ce	T : average 99Ca21=9.6(0.4) 86Wi15=9.2(1.0) 83Ni05=8.9(0.5)										
¹²⁶ Ag	-61010#	300#			107	ms	12	3 ⁺ #	03	β^- =100; β^- n=?	
¹²⁶ Cd	-72330	50			515	ms	17	0 ⁺	03	β^- =100	
¹²⁶ In	-77810	40			* 1.53	s	0.01	3 ⁽⁺⁾ #	03	β^- =100	
¹²⁶ In ^m	-77710	50	100	60	BD *	1.64	s	0.05	8 ⁽⁻⁾ #	03 79Fo10 J	β^- =100
¹²⁶ Sn	-86020	11			230	ky	14	0 ⁺	03	β^- =100	
¹²⁶ Sn ^m	-83801	11	2218.99	0.08	6.6	μ s	1.4	7 ⁻	03	IT=100	
¹²⁶ Sn ⁿ	-83456	11	2564.5	0.5	7.7	μ s	0.5	10 ⁺ #	03	IT=100	
¹²⁶ Sb	-86400	30			12.35	d	0.06	(8 ⁻)	03	β^- =100	
¹²⁶ Sb ^m	-86380	30	17.7	0.3	19.15	m	0.08	(5 ⁺)	03	β^- =86 4; IT=14 4	
¹²⁶ Sb ⁿ	-86360	30	40.4	0.3	11	s		(3 ⁻)	03	IT=100	
¹²⁶ Sb ^p	-86300	30	104.6	0.3	553	ns	5	(3 ⁺)	03	IT=100	
¹²⁶ Te	-90064.6	1.5			STABLE			0 ⁺	03	IS=18.84 25	
¹²⁶ I	-87911	4			12.93	d	0.05	2 ⁻	03	β^+ =52.7 5; β^- =47.3 5	
¹²⁶ Xe	-89169	6			STABLE			0 ⁺	03	IS=0.09 1; 2 β^+ ?	
¹²⁶ Cs	-84345	12			1.64	m	0.02	1 ⁺	03	β^+ =100	
¹²⁶ Cs ^m	-84072	12	273.0	0.7	> 1	μ s			03	IT=100	
¹²⁶ Cs ⁿ	-83749	12	596.1	1.1	171	μ s	14		03	IT=100	
¹²⁶ Ba	-82670	12			100	m	2	0 ⁺	03	β^+ =100	
¹²⁶ La	-74970	90			* 54	s	2	(5 ⁽⁺⁾)	03	β^+ =100	
¹²⁶ La ^m	-74760	400	210	410	BD *	20	s	20	(0 ⁻ , 1 ⁻ , 2 ⁻)	03	β^+ =100
¹²⁶ Ce	-70821	28			51.0	s	0.3	0 ⁺	03	β^+ =100	
¹²⁶ Pr	-60260#	200#			3.12	s	0.18	(4, 5, 6)	03 88Ba42 T	β^+ =100; β^+ p=?	
¹²⁶ Nd	-52890#	400#			1#	s	(>200 ns)	0 ⁺	03 00So11 I	β^+ ?	
¹²⁶ Pm	-39570#	500#			500#	ms				β^+ ?	
* ¹²⁶ La ^m	T : 97As05: "by far shorter than 50 s"										
* ¹²⁶ Pr	T : average 95Os03=3.14(0.22) 88Ba42=3.0(0.4) and 83Ni05=3.2(0.6)										

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹²⁷ Ag	-58900# 300#				79 ms	3	7/2 ⁺ #	98	96Wo.A TD	β^- =100; β^- -n=? *
¹²⁷ Cd	-68520 70				370 ms	70	(3/2 ⁺)	96		β^- =100
¹²⁷ In	-76990 40				1.09 s	0.01	9/2 ⁽⁺⁾	96	87Eb02 J	β^- =100; β^- -n≤0.03
¹²⁷ In ^m	-76520 70	460	70	BD	3.67 s	0.04	(1/2 ⁻)	96		β^- =100; β^- -n=0.69 4
¹²⁷ Sn	-83499 25				2.10 h	0.04	(11/2 ⁻)	96		β^- =100
¹²⁷ Sn ^m	-83494 25	4.7	0.3		4.13 m	0.03	(3/2 ⁺)	96		β^- =100
¹²⁷ Sb	-86700 5				3.85 d	0.05	7/2 ⁺	96		β^- =100
¹²⁷ Te	-88281.1 1.5				9.35 h	0.07	3/2 ⁺	96		β^- =100
¹²⁷ Te ^m	-88192.8 1.5	88.26	0.08		109 d	2	11/2 ⁻	96		IT=97.6 2; β^- =2.4 2
¹²⁷ I	-88983 4				STABLE		5/2 ⁺	96		IS=100.
¹²⁷ Xe	-88321 4				36.345 d	0.003	1/2 ⁺	96	02Un02 T	ϵ =100
¹²⁷ Xe ^m	-88024 4	297.10	0.08		69.2 s	0.9	9/2 ⁻	96		IT=100
¹²⁷ Cs	-86240 6				6.25 h	0.10	1/2 ⁺	96		β^+ =100
¹²⁷ Cs ^m	-85788 6	452.23	0.21		55 μ s	3	(11/2 ⁻)	96		IT=100
¹²⁷ Ba	-82816 11				12.7 m	0.4	1/2 ⁺	96		β^+ =100
¹²⁷ Ba ^m	-82736 11	80.33	0.12		1.9 s	0.2	7/2 ⁻	96		IT=100
¹²⁷ La	-77896 26				5.1 m	0.1	(11/2 ⁻)	96		β^+ =100
¹²⁷ La ^m	-77881 26	14.8	1.2		3.7 m	0.4	(3/2 ⁺)	96		β^+ ≈100; IT ?
¹²⁷ Ce	-71980 60			*	29 s	2	5/2 ⁺ #	98	96Ge07 T	β^+ =100
¹²⁷ Ce ^m	-71980# 120#	0#	100#	*	34 s	2	(1/2 ⁺)	98	96Ge07 TJD	β^+ =100
¹²⁷ Pr	-64430# 200#				4.2 s	0.3	3/2 ⁺ #	98		β^+ =100
¹²⁷ Pr ^m	-63830# 280#	600#	200#		50# ms		11/2 ⁻	96	98Mo30 J	β^+ ?; IT ?
¹²⁷ Nd	-55420# 400#				1.8 s	0.4	5/2 ⁺ #	96		β^+ =100; β^+ p=?
¹²⁷ Pm	-45060# 600#				1# s		5/2 ⁺ #			β^+ ?; p ?
* ¹²⁷ Ag	T : supersedes 95Fe12=109(25) from same group									**
¹²⁸ Ag	-54800# 300#				58 ms	5		01		β^- =100; β^- -n=?
¹²⁸ Cd	-67290 290				280 ms	40	0 ⁺	01		β^- =100
¹²⁸ In	-74360 50				840 ms	60	(3) ⁺	01	93Ru01 D	β^- =100; β^- -n=0.038 3
¹²⁸ In ^m	-74110 50	247.87	0.10		10 ms	7	(1) ⁻	01		IT=100 *
¹²⁸ In ⁿ	-74040 50	320	60	BD	720 ms	100	(8 ⁻)	01		β^- =100
¹²⁸ Sn	-83335 27				59.07 m	0.14	0 ⁺	01		β^- =100
¹²⁸ Sn ^m	-81244 27	2091.50	0.11		6.5 s	0.5	(7 ⁻)	01		IT=100
¹²⁸ Sb	-84609 25			*	9.01 h	0.04	8 ⁻	01		β^- =100
¹²⁸ Sb ^m	-84599 24	10	7	*	10.4 m	0.2	5 ⁺	01		β^- =96.4 10; IT=3.6 10 *
¹²⁸ Te	-88992.1 1.7				2.2 Yy	0.3	0 ⁺	01	96Ta04 T	IS=31.74 8; 2 β^- =100 *
¹²⁸ Te ^m	-86201.4 1.7	2790.7	0.4		370 ns	30	10 ⁺	01		IT=100
¹²⁸ I	-87738 4				24.99 m	0.02	1 ⁺	01		β^- =93.1 8; β^+ =6.9 8
¹²⁸ I ^m	-87600 4	137.850	0.004		845 ns	20	4 ⁻	01		IT=100
¹²⁸ I ⁿ	-87571 4	167.367	0.005		175 ns	15	(6) ⁻	01		IT=100
¹²⁸ Xe	-89860.0 1.4				STABLE		0 ⁺	01		IS=1.92 3
¹²⁸ Xe ^m	-87072.7 1.5	2787.3	0.4		83 ns	2	8 ⁻	01		IT=100
¹²⁸ Cs	-85931 5				3.640 m	0.014	1 ⁺	01	93Al03 T	β^+ =100 *
¹²⁸ Ba	-85402 10				2.43 d	0.05	0 ⁺	01		ϵ =100
¹²⁸ La	-78630 50			*	5.18 m	0.14	(5 ⁺)	01		β^+ =100
¹²⁸ La ^m	-78530# 110#	100#	100#	*	< 1.4 m		(1 ⁺ , 2 ⁻)	01		β^+ =100
¹²⁸ Ce	-75534 28				3.93 m	0.02	0 ⁺	01		β^+ =100
¹²⁸ Pr	-66331 30				2.84 s	0.09	(3 ⁺)	01	99Xi03 J	β^+ =100; β^+ p=? *
¹²⁸ Nd	-60180# 200#				5# s		0 ⁺	01		β^+ ?; β^+ p ? *
¹²⁸ Pm	-48050# 400#				1.0 s	0.3	6 ⁺ #	01	93Li40 D	β^+ ≈100; β^+ p ?; p=0 *
¹²⁸ Sm	-39050# 500#				500# ms		0 ⁺			β^+ ?; p ? *
* ¹²⁸ In ^m	T : 10 μ s < half-life < 20 ms, cf. ENSDF									**
* ¹²⁸ Sb ^m	E : less than 20 keV above ground state, cf. ENSDF									**
* ¹²⁸ Te	T : see also 92Be30=7.7(0.4) not used for consistency with ¹³⁰ Te (see below)									**
* ¹²⁸ Cs	T : average 93Al03=3.66(0.02) 76He04=3.62(0.02)									**
* ¹²⁸ Pr	D : from 85Wi07									**
* ¹²⁸ Nd	T : 83Ni05 gave 4(2) s. Proved, by 85Wi07, to be due to ¹²⁸ Pr, not to ¹²⁸ Nd									**
* ¹²⁸ Pm	D : p=0 from 93Li40 J : as calculated by 02Xu11									**

Nuclide	Mass excess (keV)		Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹²⁹ Ag	-52450# 400#			*	44 ms	7	7/2 ⁺ #	03	β^- =100; β^- n=?
¹²⁹ Ag ^m	-52450# 450#	0#	200#	EU *	160 ms		1/2 ⁻ #	03	β^- ?; β^- n? *
¹²⁹ Cd	-63200# 300#			*	242 ms	8	3/2 ⁺ #	96	03Pf.A TD β^- =100; β^- n=?
¹²⁹ Cd ^m	-63200# 360#	0#	200#	*	104 ms	6	11/2 ⁻ #	96	03Pf.A TD β^- =100; β^- n=?
¹²⁹ In	-72940 40				611 ms	4	9/2 ⁺ #	96	93Ru01 T β^- =100; β^- n=0.25 5 *
¹²⁹ In ^m	-72560 70	380	70	BD	1.23 s	0.03	1/2 ⁻ #	96	β^- ≈100; IT<0.3; ... *
¹²⁹ In ⁿ	-71250 40	1688.0	0.5		8.5 μs	0.5	17/2 ⁻		03Ge04 ETJ IT=100
¹²⁹ Sn	-80594 29				2.23 m	0.04	3/2 ⁺ #	96	β^- =100
¹²⁹ Sn ^m	-80559 29	35.2	0.3		6.9 m	0.1	11/2 ⁻ #	96	β^- ≈100; IT≈0.002
¹²⁹ Sb	-84628 21				4.40 h	0.01	7/2 ⁺	96	β^- =100
¹²⁹ Sb ^m	-82777 21	1851.05	0.10		17.7 m	0.1	(19/2 ⁻)	96	β^- =85; IT=15
¹²⁹ Sb ⁿ	-82767 21	1860.90	0.10		> 2 μs		(15/2 ⁻)	96	IT=100
¹²⁹ Sb ^p	-82489 21	2138.9	0.5		1.1 μs	0.1	(23/2 ⁺)		03Ge04 ETJ IT=100
¹²⁹ Te	-87003.2 1.8				69.6 m	0.3	3/2 ⁺	96	β^- =100
¹²⁹ Te ^m	-86897.7 1.8	105.50	0.05		33.6 d	0.1	11/2 ⁻	96	IT=63 17; β^- =37 17
¹²⁹ I	-88503 3				15.7 My	0.4	7/2 ⁺	96	β^- =100
¹²⁹ Xe	-88697.4 0.7				STABLE		1/2 ⁺	96	IS=26.44 24
¹²⁹ Xe ^m	-88461.3 0.7	236.14	0.05		8.88 d	0.02	11/2 ⁻	96	IT=100
¹²⁹ Cs	-87500 5				32.06 h	0.06	1/2 ⁺	96	β^+ =100
¹²⁹ Ba	-85065 11				2.23 h	0.11	1/2 ⁺	96	β^+ =100
¹²⁹ Ba ^m	-85057 11	8.42	0.06		2.16 h	0.02	7/2 ⁺ #	96	β^+ ≈100; IT=?
¹²⁹ La	-81326 21				11.6 m	0.2	3/2 ⁺	96	β^+ =100
¹²⁹ La ^m	-81154 21	172.1	0.4		560 ms	50	11/2 ⁻	96	IT=100
¹²⁹ Ce	-76287 28				3.5 m	0.3	(5/2 ⁺)	97	93A103 T β^+ =100 *
¹²⁹ Ce ^m	-76179 28	107.6	0.1		62 ns	5	(7/2 ⁻)	96	IT=100
¹²⁹ Pr	-69774 30			&	30 s	4	(3/2 ⁺)	96	96Gi08 J β^+ =100
¹²⁹ Pr ^m	-69390 30	382.7	0.5	&	1# ms		(11/2 ⁻)		97Gi07 EJD IT=100
¹²⁹ Nd	-62240# 200#				4.9 s	0.2	5/2 ⁺ #	96	β^+ =100; β^+ p=?
¹²⁹ Pm	-52950# 400#				3# s	(>200 ns)	5/2 ⁺ #		00So11 I β^+ ? *
¹²⁹ Sm	-42250# 500#				550 ms	100	5/2 ⁺ #		99Xu05 TD β^+ =100
* ¹²⁹ Ag	I : the evaluators are not convinced by the identification arguments **								
* ¹²⁹ In	T : average 93Ru01=611(5) 86Wa17=610(10) **								
* ¹²⁹ In ^m	D : ... ; β^- n=2.5 5 **								
* ¹²⁹ Ce	J : from 96Gi08 (5/2 ⁺ in ENSDF was from theory) **								
¹³⁰ Ag	-46160# 330#				50 ms		0 ⁺	01	β^- =100; β^- n?
¹³⁰ Cd	-61570 280				162 ms	7	0 ⁺	01	01Ha39 TD β^- =100; β^- n=3.5 10
¹³⁰ In	-69890 40			*	290 ms	20	(1 ⁻)	01	β^- =100; β^- n=0.93 13
¹³⁰ In ^m	-69840 40	50	50	BD *	538 ms	5	10 ⁻ #	01	93Ru01 T β^- =100; β^- n=1.65 15 *
¹³⁰ In ⁿ	-69490 50	400	60	BD	540 ms	10	(5 ⁺)	01	β^- =100; β^- n=1.65 15
¹³⁰ Sn	-80139 11				3.72 m	0.07	0 ⁺	01	β^- =100
¹³⁰ Sn ^m	-78192 11	1946.88	0.10		1.7 m	0.1	7 ⁻ #	01	β^- =100
¹³⁰ Sb	-82292 17				39.5 m	0.8	8 ⁻ #	01	β^- =100
¹³⁰ Sb ^m	-82287 17	4.80	0.20		6.3 m	0.2	(4,5 ⁺)	01	β^- =100
¹³⁰ Te	-87351.4 1.9				790 Ey	100	0 ⁺	01	96Ta04 TD IS=34.08 62; 2 β^- =100 *
¹³⁰ Te ^m	-85205.0 1.9	2146.41	0.04		115 ns	8	(7 ⁻)	01	IT=100
¹³⁰ Te ⁿ	-84690 7	2661	7		1.90 μs	0.08	(10 ⁺)	01	IT=100 *
¹³⁰ Te ^p	-82976.0 2.6	4375.4	1.8		261 ns	33		01	IT=100
¹³⁰ I	-86932 3				12.36 h	0.01	5 ⁺	01	β^- =100
¹³⁰ I ^m	-86892 3	39.9525	0.0013		8.84 m	0.06	2 ⁺	01	IT=84 2; β^- =16 2
¹³⁰ Xe	-89881.7 0.7				STABLE		0 ⁺	01	IS=4.08 2
¹³⁰ Cs	-86900 8				29.21 m	0.04	1 ⁺	01	β^+ =98.4; β^- =1.6
¹³⁰ Cs ^m	-86737 8	163.25	0.11		3.46 m	0.06	5 ⁻	01	IT≈100; β^+ =0.16 2
¹³⁰ Cs ^x	-86873 17	27	15		R = .2 .1		fsmix		
¹³⁰ Ba	-87261.6 2.8				STABLE	(>4.0 Zy)	0 ⁺	01	96Ba24 T IS=0.106 1; 2 β^+ ? *
¹³⁰ Ba ^m	-84786.5 2.8	2475.12	0.18		9.54 ms	0.14	8 ⁻	01	02Mo31 T IT=100 *

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...										
¹³⁰ La	-81628	26			8.7	m	0.1	3(+)	01	$\beta^+=100$
¹³⁰ Ce	-79423	28			22.9	m	0.5	0+	01	$\beta^+=100$
¹³⁰ Ce ^m	-76969	28	2453.6	0.3	100	ns	8	(7-)	01	IT=100
¹³⁰ Pr	-71180	60			40.0	s	0.4	(6,7)(+)	01	$\beta^+=100$
¹³⁰ Pr ^m	-71080#	120#	100#	100#	10#	s		2+#	01	$\beta^+?$ *
¹³⁰ Nd	-66596	28			21	s	3	0+	01	$\beta^+=100$ *
¹³⁰ Pm	-55470#	300#			2.6	s	0.2	(5+,6+,4+)	01	$\beta^+=100; \beta^+p=?$
¹³⁰ Sm	-47580#	400#			1#	s		0+	01	$\beta^+?$
¹³⁰ Eu	-33940#	500#			1.1	ms	0.5	2+#	02Ma61	TD p=?; $\beta^+=1\#$
* ¹³⁰ In ^m	T : average 93Ru01=542(9) 85Re.A=532(6) and 86Wa17=550(10)									
* ¹³⁰ In ^m	T : ⁷⁶ Lu02=580(10) at variance, not used									
* ¹³⁰ Te	T : see also numerous (not used) results in 95Tr07									
* ¹³⁰ Te	T : treated by ENSDF'01 as a lower limit (not accepted by NUBASE)									
* ¹³⁰ Te ⁿ	E : less than 25 keV above 2648.57(0.22) (8+) level, see ENSDF'01									
* ¹³⁰ Ba ^m	T : others 66Br14=8.8(0.2) 69Wa.A=13.5(1.0) not used									
* ¹³⁰ Pr ^m	J : 88Ba42: there is also a low-spin component in ¹³⁰ Pr activity									
* ¹³⁰ Pr ^m	J : see also the discussion in 01Gi17 on three isomeric states in ¹³⁰ Pr									
* ¹³⁰ Nd	T : other conflicting data, not used: 00Xu08=13(3) 77Bo02=28(3)									
¹³¹ Cd	-55270#	300#			68	ms	3	7/2-#	00Ha55	TD $\beta^-=100; \beta^-n=3.5$ 10
¹³¹ In	-68137	28			280	ms	30	(9/2+)	94 93Ru01	D $\beta^-=100; \beta^-n=2.2$ 3
¹³¹ In ^m	-67790	40	350	40	350	ms	50	(1/2-)	94	$\beta^- \approx 100; \dots$ *
¹³¹ In ⁿ	-64040	70	4100	70	320	ms	60	(19..23/2+)	94	$\beta^- > 99; \dots$ *
¹³¹ Sn	-77314	21			56.0	s	0.5	(3/2+)	94	$\beta^-=100$
¹³¹ Sn ^m	-77230#	40#	80#	30#	58.4	s	0.5	(11/2-)	94 01Si.A	E $\beta^-=100; IT < 0.0004\#$ *
¹³¹ Sb	-81988	21			23.03	m	0.04	(7/2+)	94	$\beta^-=100$
¹³¹ Te	-85209.5	1.9			25.0	m	0.1	3/2+	94	$\beta^-=100$
¹³¹ Te ^m	-85027.3	1.9	182.250	0.020	30	h	2	11/2-	94	$\beta^-=77.8$ 16; IT=22.2 16
¹³¹ I	-87444.4	1.1			8.02070	d	0.00011	7/2+	94	$\beta^-=100$
¹³¹ Xe	-88415.2	1.0			STABLE			3/2+	94	IS=21.18 3
¹³¹ Xe ^m	-88251.3	1.0	163.930	0.008	11.84	d	0.07	11/2-	94	IT=100
¹³¹ Cs	-88060	5			9.689	d	0.016	5/2+	94	$\epsilon=100$
¹³¹ Ba	-86683.8	2.8			11.50	d	0.06	1/2+	94	$\beta^+=100$
¹³¹ Ba ^m	-86496.7	2.8	187.14	0.12	14.6	m	0.2	9/2-	94	IT=100
¹³¹ La	-83769	28			59	m	2	3/2+	94	$\beta^+=100$
¹³¹ La ^m	-83464	28	304.52	0.24	170	μ s	10	11/2-	94	IT=100
¹³¹ Ce	-79720	30			10.2	m	0.3	(7/2+)	99	$\beta^+=100$
¹³¹ Ce ^m	-79660	30	61.8	0.1	5.0	m	1.0	(1/2+)	99 96Gi08	E $\beta^+=100$
¹³¹ Ce ⁿ	-79560	30	162.00	0.09	70	ns	5	(9/2-)		
¹³¹ Pr	-74280	50			1.50	m	0.03	(3/2+)	94 96Gi08	T $\beta^+=100$ *
¹³¹ Pr ^m	-74130	50	152.4	0.2	5.7	s	0.2	(11/2-)	94 96Ge12	ED IT=96.4 12; $\beta^+=3.6$ 12
¹³¹ Nd	-67769	28			33	s	3	(5/2)(+)	94 96Ge12	T $\beta^+=100; \beta^+p=?$
¹³¹ Nd ^m	-67412	28	357	3	50	ns		(7/2-)	94 96Ge12	J IT=100
¹³¹ Pm	-59740#	200#			6.3	s	0.8	5/2+#	94 99Ga41	T $\beta^+=100; \beta^+p?$
¹³¹ Sm	-50200#	300#			1.2	s	0.2	5/2+#	94	$\beta^+=100; \beta^+p=?$
¹³¹ Eu	-39350#	400#			17.8	ms	1.9	3/2+	02	p=?; $\beta^+=12\#$
* ¹³¹ In ^m	D : ... ; $\beta^-n \leq 2.0$ 4; IT ≤ 0.018									
* ¹³¹ In ⁿ	D : ... ; $\beta^-n = 0.028$ 5; IT < 1									
* ¹³¹ Sn ^m	E : ENSDF'94=241.8(0.8) questioned from theoretical and exp. considerations									
* ¹³¹ Pr	T : average 96Gi08=1.57(0.07) 93Al03=1.48(0.02) and 83Ga.A=1.58(0.05)									

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)			
^{132}Cd	-50720#	500#	97 ms	10	0 ⁺	00Ha55	TD $\beta^- = 100; \beta^- n = 60$ 15			
^{132}In	-62420	60	206 ms	4	(7 ⁻)	02	$\beta^- = 100; \beta^- n = 6.2$ 11			
^{132}Sn	-76554	14	39.7 s	0.5	0 ⁺	92	$\beta^- = 100$			
^{132}Sb	-79674	14	2.79 m	0.05	(4 ⁺)	92	$\beta^- = 100$			
$^{132}\text{Sb}^m$	-79470	30	4.15 m	0.05	(8 ⁻)	92	$\beta^- = 100$			
^{132}Te	-85182	7	3.204 d	0.013	0 ⁺	92	$\beta^- = 100$			
^{132}I	-85700	6	2.295 h	0.013	4 ⁺	92	$\beta^- = 100$			
$^{132}\text{I}^m$	-85595	10	1.387 h	0.015	(8 ⁻)	92	IT=86.2; $\beta^- = 14$ 2			
^{132}Xe	-89280.5	1.0	STABLE		0 ⁺	92	IS=26.89 6			
$^{132}\text{Xe}^m$	-86528.2	1.0	2752.27	0.17	8.39 ms	0.11	(10 ⁺) 92	IT=100		
^{132}Cs	-87155.9	1.9	6.479 d	0.007	2 ⁺	92	$\beta^+ = 98.13$ 9; $\beta^- = 1.87$ 9			
^{132}Ba	-88434.8	1.1	STABLE	(>300 Ey)	0 ⁺	94	96Ba24 T IS=0.101 1; $2\beta^+$?			
^{132}La	-83740	40	4.8 h	0.2	2 ⁻	94	$\beta^+ = 100$			
$^{132}\text{La}^m$	-83550	40	188.18	0.11	24.3 m	0.5	6 ⁻ 94	IT=76; $\beta^+ = 24$		
^{132}Ce	-82474	21	3.51 h	0.11	0 ⁺	99	$\beta^+ = 100$			
$^{132}\text{Ce}^m$	-80133	21	2340.8	0.5	9.4 ms	0.3	(8 ⁻) 99	01Mo05 TJ IT=100		
^{132}Pr	-75210	60	1.49 m	0.11	(2 ⁺)	01	94Bu18 TJ $\beta^+ = 100$	*		
$^{132}\text{Pr}^m$	-75210#	120#	0#	100#	20#	s	(5 ⁺)	90Ko25 J $\beta^+ ?$	*	
^{132}Nd	-71426	24	1.56 m	0.10	0 ⁺	97	95Bu11 T $\beta^+ = 100$	*		
^{132}Pm	-61710#	200#	6.3 s	0.7	(3 ⁺)	92	$\beta^+ = 100; \beta^+ p \approx 5e-5$			
^{132}Sm	-55250#	300#	4.0 s	0.3	0 ⁺	92	$\beta^+ = 100; \beta^+ p ?$			
^{132}Eu	-42500#	400#	100#	ms			93Li40 D $\beta^+ ?; p=0$			
* ^{132}Pr	T : average	94Bu18=1.47(0.12)	74Ar27=1.6(0.3)					**		
* ^{132}Nd	T : average	95Bu11=1.47(0.12)	77Bo02=1.75(0.17)					**		
^{133}In	-57930#	300#	165 ms	3	(9/2 ⁺)	02	96Ho16 J $\beta^- = 100; \beta^- n = 85$ 10	*		
$^{133}\text{In}^m$	-57600#	300#	180#	ms	(1/2 ⁻)	96Ho16 J	IT ?			
^{133}Sn	-70950	40	1.45 s	0.03	7/2 ⁻	# 98	93Ru01 D $\beta^- = 100; \beta^- n = 0.0294$ 24			
^{133}Sb	-78943	25	2.5 m	0.1	(7/2 ⁺)	95	$\beta^- = 100$			
^{133}Te	-82945	24	12.5 m	0.3	(3/2 ⁺)	95	$\beta^- = 100$			
$^{133}\text{Te}^m$	-82611	24	334.26	0.04	55.4 m	0.4	(11/2 ⁻) 95	$\beta^- = 82.5$ 30; IT=17.5 30		
^{133}I	-85887	5	20.8 h	0.1	7/2 ⁺	95	$\beta^- = 100$			
$^{133}\text{I}^m$	-84253	5	1634.174	0.017	9 s	2	(19/2 ⁻) 95	IT=100		
^{133}Xe	-87643.6	2.4	5.2475 d	0.0005	3/2 ⁺	95	02Un02 T $\beta^- = 100$			
$^{133}\text{Xe}^m$	-87410.4	2.4	233.221	0.018	2.19 d	0.01	11/2 ⁻ 95	IT=100		
^{133}Cs	-88070.958	0.022	STABLE		7/2 ⁺	95	IS=100.			
^{133}Ba	-87553.5	1.0	10.51 y	0.05	1/2 ⁺	95	$\epsilon = 100$			
$^{133}\text{Ba}^m$	-87265.3	1.0	288.247	0.009	38.9 h	0.1	11/2 ⁻ 95	IT \approx 100; $\epsilon = 0.0096$ 11		
^{133}La	-85494	28	3.912 h	0.008	5/2 ⁺	95	$\beta^+ = 100$			
$^{133}\text{La}^m$	-84958	28	535.60	0.02	62 ns	3	11/2 ⁻			
^{133}Ce	-82423	16	97 m	4	1/2 ⁺	97	$\beta^+ = 100$			
$^{133}\text{Ce}^m$	-82386	16	37.1	0.8	4.9 h	0.4	9/2 ⁻ 97	$\beta^+ = 100$		
^{133}Pr	-77938	12	6.5 m	0.3	(3/2 ⁺)	97	$\beta^+ = 100$			
$^{133}\text{Pr}^m$	-77746	12	192.05	0.14	1.1 μ s	0.2	(11/2 ⁻) 97	01Xu04 T IT=100		
^{133}Nd	-72330	50	70 s	10	(7/2 ⁺)	97	$\beta^+ = 100$			
$^{133}\text{Nd}^m$	-72200	50	127.97	0.11	70 s		(1/2 ⁺) 97	95Br24 D $\beta^+ \approx 100$; IT=?		
$^{133}\text{Nd}^n$	-72150	50	176.10	0.10	300 ns		(9/2 ⁻) 97	IT=100		
^{133}Pm	-65410	50	&	15 s	3	(3/2 ⁺)	95	96Ga17 J $\beta^+ = 100$		
$^{133}\text{Pm}^m$	-65280	50	130.4	1.0	&	10#	s	(11/2 ⁻)	96Ga17 EJ $\beta^+ ?$; IT ?	*
^{133}Sm	-57130#	200#	2.90 s	0.17	(5/2 ⁺)	01	01Xu04 T $\beta^+ = 100; \beta^+ p = ?$	*		
^{133}Eu	-47280#	300#	200#	ms			11/2 ⁻	$\beta^+ ?$		
* ^{133}In	D : $\beta^- n$ intensity is from	93Ru01						**		
* $^{133}\text{Pm}^m$	E : combining γ s from Table 1:	214.7 + 357.7 + 453.8 - 252.8 - 643(1)						**		
* ^{133}Sm	T : average	01Xu04=3.1(0.5)	85Wi07=2.8(0.2)	77Bo02=3.2(0.4)				**		

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens Reference	Decay modes and intensities (%)
^{134}In	-52020#	400#			140 ms	4	high	02 96Ho16 J $\beta^- = 100; \beta^- n = 65; \dots$ *
^{134}Sn	-66800	100			1.12 s	0.08	0^+	94 $\beta^- = 100; \beta^- n = 17$ 13
^{134}Sb	-74170	40		*	780 ms	60	(0^-)	95 $\beta^- = 100$
$^{134}\text{Sb}^m$	-74090	100	80 110	BD*	10.22 s	0.09	(7^-)	95 $\beta^- = 100; \beta^- n = 0.091$ 8
^{134}Te	-82559	11			41.8 m	0.8	0^+	98 $\beta^- = 100$
$^{134}\text{Te}^m$	-80868	11	1691.24	0.17	164 ns	1	6^+	98 IT=100
^{134}I	-84072	8			52.5 m	0.2	$(4)^+$	94 $\beta^- = 100$
$^{134}\text{I}^m$	-83756	8	316.49	0.22	3.60 m	0.10	$(8)^-$	94 IT=97.7 10; $\beta^- = 2.3$ 10
^{134}Xe	-88124.5	0.8			STABLE	(>11 Py)	0^+	94 89Ba22 T IS=10.44 10; $2\beta^-$?
$^{134}\text{Xe}^m$	-86159.0	0.9	1965.5	0.5	290 ms	17	7^-	94 IT=100
^{134}Cs	-86891.181	0.026			2.0648 y	0.0010	4^+	94 $\beta^- = 100; \epsilon = 0.0003$ 1
$^{134}\text{Cs}^m$	-86752.437	0.026	138.7441	0.0026	2.903 h	0.008	8^-	94 IT=100
^{134}Ba	-88949.9	0.4			STABLE		0^+	95 IS=2.417 18
^{134}La	-85219	20			6.45 m	0.16	1^+	94 $\beta^+ = 100$
^{134}Ce	-84836	20			3.16 d	0.04	0^+	94 $\epsilon = 100$
^{134}Pr	-78510	40			& 11 m		(5^-)	94 $\beta^+ = 100$
$^{134}\text{Pr}^m$	-78510#	110#	0# 100#		& 17 m	2	2^-	94 $\beta^+ = 100$
^{134}Nd	-75646	12			8.5 m	1.5	0^+	99 $\beta^+ = 100$
$^{134}\text{Nd}^m$	-73353	12	2293.1	0.4	410 μs	30	$(8)^-$	99 IT=100
^{134}Pm	-66740	60			* 22 s	1	(5^+)	94 $\beta^+ = 100$
$^{134}\text{Pm}^m$	-66740#	120#	0# 100#		* 5 s		(2^+)	94 $\beta^+ = 100$
^{134}Sm	-61510#	200#			10 s	1	0^+	94 $\beta^+ = 100$
^{134}Eu	-49830#	200#			500 ms	200		94 $\beta^+ = 100; \beta^+ p = ?$
^{134}Gd	-41570#	400#			400# ms		0^+	$\beta^+ ?$
* ^{134}In	D : ... ; $\beta^- 2n < 4$							**
* ^{134}In	D : $\beta^- 2n$ intensity limits is from 95Jo.A							**
^{135}In	-47200#	500#			92 ms	10	$9/2^+ \#$	02 $\beta^- ?; \beta^- n ?$
^{135}Sn	-60800#	400#			530 ms	20	$(7/2^-)$	02 $\beta^- = 100; \beta^- n = 21$ 3
^{135}Sb	-69710	100			1.68 s	0.02	$(7/2^+)$	02 02Sh08 J $\beta^- = 100; \beta^- n = 22$ 3
^{135}Te	-77830	90			19.0 s	0.2	$(7/2^-)$	98 $\beta^- = 100$
$^{135}\text{Te}^m$	-76280	90	1554.88	0.17	510 ns	20	$(19/2^-)$	98 IT=100
^{135}I	-83790	7			6.57 h	0.02	$7/2^+$	98 $\beta^- = 100$
^{135}Xe	-86417	5			9.14 h	0.02	$3/2^+$	98 $\beta^- = 100$
$^{135}\text{Xe}^m$	-85890	5	526.551	0.013	15.29 m	0.05	$11/2^-$	98 IT \approx 100; $\beta^- = 0.30$ 17 *
^{135}Cs	-87581.9	1.0			2.3 My	0.3	$7/2^+$	98 $\beta^- = 100$
$^{135}\text{Cs}^m$	-85949.0	1.8	1632.9	1.5	53 m	2	$19/2^-$	98 IT=100
^{135}Ba	-87850.5	0.4			STABLE		$3/2^+$	98 IS=6.592 12
$^{135}\text{Ba}^m$	-87582.3	0.4	268.22	0.02	28.7 h	0.2	$11/2^-$	98 IT=100
^{135}La	-86651	10			19.5 h	0.2	$5/2^+$	98 $\beta^+ = 100$
^{135}Ce	-84625	11			17.7 h	0.3	$1/2^{(+)}$	98 $\beta^+ = 100$
$^{135}\text{Ce}^m$	-84179	11	445.8	0.2	20 s	1	$(11/2^-)$	98 IT=100
^{135}Pr	-80936	12			24 m	2	$3/2^{(+)}$	98 $\beta^+ = 100$
$^{135}\text{Pr}^m$	-80578	12	358.06	0.06	105 μs	10	$(11/2^-)$	98 IT=100
^{135}Nd	-76214	19			12.4 m	0.6	$9/2^{(-)}$	98 $\beta^+ = 100$
$^{135}\text{Nd}^m$	-76149	19	65.0	0.2	5.5 m	0.5	$(1/2^+)$	98 $\beta^+ > 99.97; IT < 0.03$
^{135}Pm	-69980	60			* & 49 s	3	$(5/2^+, 3/2^+)$	98 $\beta^+ = 100$
$^{135}\text{Pm}^m$	-69930#	120#	50# 100#		* & 40 s	3	$(11/2^-)$	98 89Ko07 TJ $\beta^+ = 100$
^{135}Sm	-62860	150			* 10.3 s	0.5	$(7/2^+)$	98 77Bo02 J $\beta^+ = 100; \beta^+ p = 0.02$ 1
$^{135}\text{Sm}^m$	-62860#	340#	0# 300#		* 2.4 s	0.9	$(3/2^+, 5/2^+)$	98 89Vi04 TJD $\beta^+ = 100$ *
^{135}Eu	-54190#	300#			1.5 s	0.2	$11/2^- \#$	98 $\beta^+ = 100; \beta^+ p ?$
^{135}Gd	-44180#	500#			1.1 s	0.2	$3/2^-$	98 98St28 J $\beta^+ = 100; \beta^+ p \approx 2$
* $^{135}\text{Xe}^m$	D : β^- ranging 0.004 to 0.6%							**
* $^{135}\text{Sm}^m$	I : existence of $^{135}\text{Sm}^m$ and spins of both states are discussed in ENSDF							**

Nuclide	Mass excess (keV)	Excitation energy (keV)				Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{136}Sn	−56500# 500#					250 ms	30	0^+	02		β^- =100; β^- n=30 5
^{136}Sb	−64880# 300#					923 ms	14	1^- #	02		β^- =100; β^- n=16.3 32... *
$^{136}\text{Sb}^m$	−64710# 300#	173	3			570 ns	50	6^- #	02	01Mi22 E	IT=100
^{136}Te	−74430 50					17.63 s	0.08	0^+	02		β^- =100; β^- n=1.31 5
^{136}I	−79500 50					83.4 s	1.0	(1^-)	02		β^- =100
$^{136}\text{I}^m$	−78850 110	650	120	BD		46.9 s	1.0	(6^-)	02		β^- =100; IT=0
^{136}Xe	−86425 7					STABLE	(>10 Zy)	0^+	02	02Be74 T	IS=8.87 16; $2\beta^-$?
$^{136}\text{Xe}^m$	−84533 7	1891.703	0.014			2.95 μ s	0.09	6^+	02		IT=100
^{136}Cs	−86338.7 1.9				*	13.16 d	0.03	5^+	02		β^- =100
$^{136}\text{Cs}^m$	−85821 5	518	5	*		19 s	2	8^-	02	83We07 E	IT=?; β^- ?
^{136}Ba	−88886.9 0.4					STABLE		0^+	02		IS=7.854 24
$^{136}\text{Ba}^m$	−86856.4 0.4	2030.466	0.018			308.4 ms	1.9	7^-	02		IT=100
^{136}La	−86040 50					9.87 m	0.03	1^+	02		β^+ =100
$^{136}\text{La}^m$	−85790 50	255	9			114 ms	3	$(8)^{-\#}$	02	ABBW E	IT=100
^{136}Ce	−86468 13					STABLE	(>38 Py)	0^+	02	01Da22 T	IS=0.185 2; $2\beta^+$?
$^{136}\text{Ce}^m$	−83373 13	3095.5	0.4			2.2 μ s	0.2	10^+	02		IT=100
^{136}Pr	−81327 12					13.1 m	0.1	2^+	02		β^+ =100
$^{136}\text{Pr}^m$	−80732 12	594.62	0.22			91.7 ns	0.9	$(6)^+$	02		IT=100
^{136}Nd	−79199 12					50.7 m	0.3	0^+	02		β^+ =100
^{136}Pm	−71200 80				* &	107 s	6	(5^-)	02		β^+ =100
$^{136}\text{Pm}^m$	−71070 90	130	120	BD * &		47 s	2	(2^+)	02		β^+ =100
^{136}Sm	−66811 12					47 s	2	0^+	02		β^+ =100
$^{136}\text{Sm}^m$	−64546 12	2264.7	1.1			15 μ s	1	(8^-)	02		IT=100
^{136}Eu	−56260# 200#				*	3.3 s	0.3	(7^+)	02	89Vi04 D	β^+ =100; β^+ p=0.09 3
$^{136}\text{Eu}^m$	−56260# 540#	0#	500#	*		3.8 s	0.3	(3^+)	02	89Vi04 D	β^+ =100; β^+ p=0.09 3
^{136}Gd	−49050# 400#					1# s	(>200 ns)	0^+	02	00So11 I	β^+ ?
^{136}Tb	−35970# 600#					200# ms			02		β^+ ?
* ^{136}Sb	D : ... ; β^- 2n=0.28#										**
* $^{136}\text{La}^m$	E : approx. 10–40 keV above 230.1 level, from ENSDF'02, thus 230.1 + 25(9)										**
^{137}Sn	−50310# 600#					190 ms	60	$5/2^-$ #	02		β^- =100; β^- n=58 15
^{137}Sb	−60260# 400#					450 ms	50	$7/2^+$ #	94	02Sh08 TD	β^- =100; β^- n=49 10
^{137}Te	−69560 120					2.49 s	0.05	$3/2^-$ #	94	93Ru01 D	β^- =100; β^- n=2.99 16
^{137}I	−76503 28					24.13 s	0.12	$(7/2^+)$	94	93Ru01 TD	β^- =100; β^- n=7.14 23 *
^{137}Xe	−82379 7					3.818 m	0.013	$7/2^-$	94		β^- =100
^{137}Cs	−86545.6 0.5					30.1671 y	0.0013	$7/2^+$	01	02Un02 T	β^- =100
^{137}Ba	−87721.2 0.4					STABLE		$3/2^+$	97		IS=11.232 24
$^{137}\text{Ba}^m$	−87059.5 0.4	661.659	0.003			2.552 m	0.001	$11/2^-$	97		IT=100
^{137}La	−87101 13					60 ky	20	$7/2^+$	94		ϵ =100
^{137}Ce	−85879 13					9.0 h	0.3	$3/2^+$	94		β^+ =100
$^{137}\text{Ce}^m$	−85625 13	254.29	0.05			34.4 h	0.3	$11/2^-$	94		IT=99.22 3; β^+ =0.78 3
^{137}Pr	−83177 12					1.28 h	0.03	$5/2^+$	94		β^+ =100
$^{137}\text{Pr}^m$	−82616 12	561.22	0.23			2.66 μ s		$11/2^-$			
^{137}Nd	−79580 11					38.5 m	1.5	$1/2^+$	01		β^+ =100
$^{137}\text{Nd}^m$	−79061 11	519.43	0.17			1.60 s	0.15	$(11/2^-)$	01		IT=100
^{137}Pm	−74073 13				&	2# m		$5/2^+$ #			β^+ ?
$^{137}\text{Pm}^m$	−73920 50	150	50	BD &		2.4 m	0.1	$11/2^-$	94		β^+ =100
^{137}Sm	−68030 40					45 s	1	$(9/2^-)$	94		β^+ =100
$^{137}\text{Sm}^m$	−67850# 60#	180#	50#			20# s		$1/2^+$ #			β^+ ?
^{137}Eu	−60020# 200#					8.4 s	0.5	$11/2^-$ #	94	88Be.A T	β^+ =100
^{137}Gd	−51210# 400#					2.2 s	0.2	$7/2^+$ #	94	99Xu05 T	β^+ =100; β^+ p=?
^{137}Tb	−41000# 600#					600# ms		$11/2^-$ #	96		p ?; β^+ ?
* ^{137}I	T : supersedes 74Ru08=24.5(0.2) from same group										**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{138}Sb	-55150#	300#		500#	ms	(>300 ns)	2 ⁻ #	03 94Be24 I	β^- ?; β^-n ?
^{138}Te	-65930#	210#		1.4	s	0.4	0 ⁺	03	β^- =100; β^-n =6.3 21
^{138}I	-72330	80		6.23	s	0.03	(2 ⁻)	03 93Ru01 D	β^- =100; β^-n =5.46 18
^{138}Xe	-80150	40		14.08	m	0.08	0 ⁺	03	β^- =100
^{138}Cs	-82887	9		33.41	m	0.18	3 ⁻	03	β^- =100
$^{138}\text{Cs}^m$	-82807	9	79.9 0.3	2.91	m	0.08	6 ⁻	03	IT=81 2; β^- =19 2
$^{138}\text{Cs}^x$	-82847	25	40 23	R=?			fsmix		
^{138}Ba	-88261.6	0.4		STABLE			0 ⁺	03	IS=71.698 42
$^{138}\text{Ba}^m$	-86171.1	0.4	2090.54 0.06	800	ns	100	6 ⁺	03	IT=100
^{138}La	-86525	4		102	Gy	1	5 ⁺	03	IS=0.090 1; ... *
$^{138}\text{La}^m$	-86452	4	72.57 0.03	116	ns	5	(3 ⁺)	03	IT=100
^{138}Ce	-87569	10		STABLE		(>150 Ty)	0 ⁺	03 01Da22 T	IS=0.251 2; 2 β^+ ?
$^{138}\text{Ce}^m$	-85440	10	2129.17 0.12	8.65	ms	0.20	7 ⁻	03	IT=100
^{138}Pr	-83132	14		1.45	m	0.05	1 ⁺	03	β^+ =100
$^{138}\text{Pr}^m$	-82783	17	348 23 BD	2.12	h	0.04	7 ⁻	03	β^+ =100
^{138}Nd	-82018	12		5.04	h	0.09	0 ⁺	03	β^+ =100
$^{138}\text{Nd}^m$	-78843	12	3174.9 0.4	410	ns	50	(10 ⁺)	03	IT=100
^{138}Pm	-74940	27		10	s	2	1 ⁺ #	03	β^+ =100
$^{138}\text{Pm}^m$	-74911	13	30 30 BD *	3.24	m	0.05	5 ⁻ #	03	β^+ =100
$^{138}\text{Pm}^n$			non existent EU	3.24	m	0.05	(3 ⁺)	03 81De38 I	β^+ =100 *
^{138}Sm	-71498	12		3.1	m	0.2	0 ⁺	03	β^+ =100
^{138}Eu	-61750	28		12.1	s	0.6	(6 ⁻)	03	β^+ =100
^{138}Gd	-55780#	200#		4.7	s	0.9	0 ⁺	03	β^+ =100
$^{138}\text{Gd}^m$	-53550#	200#	2232.7 1.1	6	μs	1	(8 ⁻)	03	
^{138}Tb	-43630#	400#		800#	ms	(>200 ns)	0 ⁺	03 00So11 I	β^+ ?; p=0 *
^{138}Dy	-34940#	600#		200#	ms		0 ⁺		β^+ ? *
* ^{138}La	D : ... ; β^+ =65.6 5; β^- =34.4 5 **								
* $^{138}\text{Pm}^n$	D : arguments for a second isomer, of intermediate spin, are not convincing **								
* ^{138}Tb	D : from 93Li40 **								
^{139}Sb	-50320#	500#		300#	ms	(>300 ns)	7/2 ⁺ #	01 94Be24 I	β^- ?
^{139}Te	-60800#	400#		500#	ms	(>300 ns)	5/2 ⁻ #	01 94Be24 I	β^- ?; β^-n ?
^{139}I	-68840	30		2.282	s	0.010	7/2 ⁺ #	01 93Ru01 T	β^- =100; β^-n =10.0 3 *
^{139}Xe	-75644	21		39.68	s	0.14	3/2 ⁻	01	β^- =100
^{139}Cs	-80701	3		9.27	m	0.05	7/2 ⁺	01	β^- =100
^{139}Ba	-84913.7	0.4		83.1	m	0.3	(7/2 ⁻)	01	β^- =100
^{139}La	-87231.4	2.4		STABLE			7/2 ⁺	01	IS=99.910 1
^{139}Ce	-86952	7		137.641	d	0.020	3/2 ⁺	01	ϵ =100
$^{139}\text{Ce}^m$	-86198	7	754.24 0.08	56.54	s	0.13	11/2 ⁻	01 94It.A T	IT=100
^{139}Pr	-84823	8		4.41	h	0.04	5/2 ⁺	01	β^+ =100
^{139}Nd	-81992	26		29.7	m	0.5	3/2 ⁺	01	β^+ =100
$^{139}\text{Nd}^m$	-81761	26	231.15 0.05	5.50	h	0.20	11/2 ⁻	01	β^+ =88.2 4; IT=11.8 4
^{139}Pm	-77496	13		4.15	m	0.05	(5/2 ⁺)	01	β^+ =100
$^{139}\text{Pm}^m$	-77307	13	188.7 0.3	180	ms	20	(11/2 ⁻)	01	IT \approx 100; β^+ =0.16#
^{139}Sm	-72380	11		2.57	m	0.10	1/2 ⁺	01	β^+ =100
$^{139}\text{Sm}^m$	-71923	11	457.40 0.22	10.7	s	0.6	11/2 ⁻	01	IT=93.7 5; β^+ =6.3 5
^{139}Eu	-65398	13		17.9	s	0.6	(11/2 ⁻)	01	β^+ =100
^{139}Gd	-57530#	200#		5.7	s	0.3	9/2 ⁻ #	01 99Xi04 T	β^+ =100; β^+p ? *
$^{139}\text{Gd}^m$	-57280#	250#	150#	4.8	s	0.9	1/2 ⁺ #	01	β^+ =100; β^+p ? *
^{139}Tb	-48170#	300#		1.6	s	0.2	11/2 ⁻ #	01	β^+ =100; β^+p ?
^{139}Dy	-37690#	500#		600	ms	200	7/2 ⁺ #	01	β^+ =100; β^+p ?
* ^{139}I	T : average 93Ru01=2.280(0.011) 80Al15=2.29(0.02) **								
* ^{139}Gd	T : average 99Xi04=5.8(0.9) 88Be.A=5.8(0.4); other 83Ni05=4.9(1.0) not used **								
* ^{139}Gd	T : since it corresponds to a mixture of ground-state and isomer **								
* $^{139}\text{Gd}^m$	D : assuming that the delayed protons reported by 83Ni05 are from both states **								

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁴⁰ Te	-56960#	300#		300# ms (>300 ns)	0 ⁺	98	94Be24 I	β^- ?; β^-_n ?
¹⁴⁰ I	-64270#	200#		860 ms	40	(3) ^(-#)	95	β^- =100; β^-_n =9.3 10
¹⁴⁰ Xe	-72990	60		13.60 s	0.10	0 ⁺	02	β^- =100
¹⁴⁰ Cs	-77051	8		63.7 s	0.3	1 ⁻	95	β^- =100
¹⁴⁰ Ba	-83271	8		12.752 d	0.003	0 ⁺	98	β^- =100
¹⁴⁰ La	-84321.0	2.4		1.6781 d	0.0003	3 ⁻	95	β^- =100
¹⁴⁰ Ce	-88083.3	2.5		STABLE		0 ⁺	95	IS=88.450 51
¹⁴⁰ Ce ^m	-85975.5	2.5	2107.85	0.03	7.3 μ s	1.5	6 ⁺	
¹⁴⁰ Pr	-84695	6		3.39 m	0.01	1 ⁺	95	β^+ =100
¹⁴⁰ Pr ^m	-83932	6	763.3	0.7	3.05 μ s	0.20	(8) ⁻	
¹⁴⁰ Nd	-84252	28		3.37 d	0.02	0 ⁺	95	ϵ =100
¹⁴⁰ Nd ^m	-82031	28	2221.4	0.1	600 μ s	50	7 ⁻	IT=100
¹⁴⁰ Pm	-78210	40		9.2 s	0.2	1 ⁺	95	β^+ =100
¹⁴⁰ Pm ^m	-77783	13	420	40	5.95 m	0.05	8 ⁻	β^+ =100
¹⁴⁰ Sm	-75456	12		14.82 m	0.12	0 ⁺	95	β^+ =100
¹⁴⁰ Eu	-66990	50		1.51 s	0.02	1 ⁺	95	β^+ =100
¹⁴⁰ Eu ^m	-66780	50	210	15	125 ms	2	5 ^{-#}	IT \approx 100; β^+ <1 *
¹⁴⁰ Gd	-61782	28		15.8 s	0.4	0 ⁺	95	β^+ =100
¹⁴⁰ Tb	-50480	800		2.4 s	0.2	5	97	β^+ =100; β^+_p =0.26 13
¹⁴⁰ Dy	-42840#	500#		700# ms		0 ⁺	02	β^+ ?
¹⁴⁰ Dy ^m	-40670#	500#	2166.1	0.5	7.0 μ s	0.5	(8) ⁻	β^+ ?
¹⁴⁰ Ho	-29310#	500#		6 ms	3	8 ^{+#}	02	p=?; β^+ =1#
* ¹⁴⁰ Eu ^m	E : less than 50 keV above 185.3 level, from ENSDF, thus 185.3 + 25(15) **							
¹⁴¹ Te	-51560#	400#		100# ms (>300 ns)	5/2 ^{-#}	01	94Be24 I	β^- ?; β^-_n ?
¹⁴¹ I	-60520#	200#		430 ms	20	7/2 ^{+#}	01	β^- =100; β^-_n =21 3
¹⁴¹ Xe	-68330	90		1.73 s	0.01	5/2 ^(-#)	01	β^- =100; β^-_n =0.044 5
¹⁴¹ Cs	-74477	11		24.84 s	0.16	7/2 ⁺	01	β^- =100; β^-_n =0.035 3
¹⁴¹ Ba	-79726	8		18.27 m	0.07	3/2 ⁻	01	β^- =100
¹⁴¹ La	-82938	5		3.92 h	0.03	(7/2 ⁺)	01	β^- =100
¹⁴¹ Ce	-85440.1	2.5		32.508 d	0.013	7/2 ⁻	01	β^- =100
¹⁴¹ Pr	-86020.9	2.5		STABLE		5/2 ⁺	01	IS=100.
¹⁴¹ Nd	-84198	4		2.49 h	0.03	3/2 ⁺	01	β^+ =100
¹⁴¹ Nd ^m	-83441	4	756.51	0.05	62.0 s	0.8	11/2 ⁻	70Ab05 D
¹⁴¹ Pm	-80523	14		20.90 m	0.05	5/2 ⁺	01	β^+ =100
¹⁴¹ Pm ^m	-79895	14	628.40	0.10	630 ns	20	11/2 ⁻	IT=100
¹⁴¹ Sm	-75939	9		10.2 m	0.2	1/2 ⁺	01	β^+ =100
¹⁴¹ Sm ^m	-75763	9	176.0	0.3	22.6 m	0.2	11/2 ⁻	β^+ \approx 100; IT=0.31 3
¹⁴¹ Eu	-69927	13		40.7 s	0.7	5/2 ⁺	01	β^+ =100
¹⁴¹ Eu ^m	-69831	13	96.45	0.07	2.7 s	0.3	11/2 ⁻	IT=86 3; β^+ =14 3
¹⁴¹ Gd	-63224	20		14 s	4	(1/2 ⁺)	01	β^+ =100; β^+_p =0.03 1
¹⁴¹ Gd ^m	-62846	20	377.8	0.2	24.5 s	0.5	(11/2 ⁻)	β^+ =89 2; IT=11 2
¹⁴¹ Tb	-54540	110		3.5 s	0.2	(5/2 ⁻)	01	β^+ =100
¹⁴¹ Tb ^m	-54540#	230#	0#	200#	7.9 s	0.6	11/2 ^{-#}	88Be.A I
¹⁴¹ Dy	-45320#	300#		900 ms	200	(9/2 ⁻)	01	β^+ =100; β^+_p =?
¹⁴¹ Ho	-34370#	500#		4.1 ms	0.3	(7/2 ⁻)	02	p=?; β^+ =1#
¹⁴¹ Ho ^m	-34300#	500#	66	2	6.4 μ s	0.8	(1/2 ⁺)	02
* ¹⁴¹ Tb ^m	I : existence discussed in 88Be.A. Provisionally accepted **							
* ¹⁴¹ Ho ^m	T : from 01Se03=6.5(+0.7-0.9) **							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁴² Te	-47430# 600#		50# ms (>300 ns)	0 ⁺	00	94Be24 I	β^- ?	
¹⁴² I	-55720# 400#		200 ms	2 ⁻ #	00		β^- =100; β^- n=25#	
¹⁴² Xe	-65480 100		1.22 s 0.02	0 ⁺	00	03Be05 TD	β^- =100; β^- n=0.36 3	
¹⁴² Cs	-70515 11		1.689 s 0.011	0 ⁻	00	93Ru01 T	β^- =100; β^- n=0.090 4 *	
¹⁴² Ba	-77823 6		10.6 m 0.2	0 ⁺	00		β^- =100 *	
¹⁴² La	-80035 6		91.1 m 0.5	2 ⁻	00		β^- =100	
¹⁴² Ce	-84538.5 3.0		STABLE (>50 Py)	0 ⁺	00		IS=11.114 51; α ?; $2\beta^-$? *	
¹⁴² Pr	-83792.7 2.5		19.12 h 0.04	2 ⁻	00		β^- \approx 100; ϵ =0.0164 8	
¹⁴² Pr ^m	-83789.0 2.5	3.694 0.003	14.6 m 0.5	5 ⁻	00		IT=100	
¹⁴² Nd	-85955.2 2.3		STABLE	0 ⁺	00		IS=27.2 5	
¹⁴² Pm	-81157 25		40.5 s 0.5	1 ⁺	00		β^+ =100	
¹⁴² Pm ^m	-80274 25	883.17 0.16	2.0 ms 0.2	(8) ⁻	00		IT=100	
¹⁴² Sm	-78993 6		72.49 m 0.05	0 ⁺	00		β^+ =100	
¹⁴² Eu	-71320 30		2.36 s 0.10	1 ⁺	00	91Fi03 T	β^+ =100 *	
¹⁴² Eu ^m	-70856 12	460 30 BD	1.223 m 0.008	8 ⁻	00		β^+ =100	
¹⁴² Gd	-66960 28		70.2 s 0.6	0 ⁺	00		β^+ =100	
¹⁴² Tb	-57060# 300#		597 ms 17	1 ⁺	00		β^+ =100; β^+ p=0.0022 11	
¹⁴² Tb ^m	-56780# 300#	280.2 1.0	303 ms 17	(5 ⁻)	00		IT \approx 100; β^+ <0.5	
¹⁴² Dy	-49960# 360#		2.3 s 0.3	0 ⁺	00		β^+ =100; β^+ p=0.06 3	
¹⁴² Ho	-37470# 500#		400 ms 100	(6to9)	02		β^+ \approx 100; β^+ p=?; p \approx 0	
* ¹⁴² Cs	T : average 93Ru01=1.684(0.014) 77Re05=1.70(0.02)							**
* ¹⁴² Ba	D : β^- n=0.091(0.003)% in ENSDF'00 contradicts $Q(\beta^-)$ =-2955(7) keV							**
* ¹⁴² Ce	T : lower limit is for α decay; for $\beta\beta$ decay 01Da22>260 Py							**
* ¹⁴² Eu	T : average 91Fi03=2.34(0.12) 75Ke08=2.4(0.2)							**
¹⁴³ I	-51640# 400#		100# ms (>300 ns)	7/2 ⁺ #	02	94Be24 I	β^- ?; β^- n=40#	
¹⁴³ Xe	-60450# 200#		511 ms 6	5/2 ⁻	02	03Be05 TD	β^- =100; β^- n=1.00 15	
¹⁴³ Cs	-67671 24		1.791 s 0.007	3/2 ⁺	02		β^- =100; β^- n=1.64 7	
¹⁴³ Ba	-73936 13		14.5 s 0.3	5/2 ⁻	02		β^- =100	
¹⁴³ La	-78187 15		14.2 m 0.1	(7/2) ⁺	02		β^- =100	
¹⁴³ Ce	-81612.0 3.0		33.039 h 0.006	3/2 ⁻	02		β^- =100	
¹⁴³ Pr	-83073.5 2.6		13.57 d 0.02	7/2 ⁺	02		β^- =100	
¹⁴³ Nd	-84007.4 2.3		STABLE	7/2 ⁻	02		IS=12.2 2	
¹⁴³ Pm	-82966 3		265 d 7	5/2 ⁺	02		ϵ =100; e^+ <5.7e-6	
¹⁴³ Pm ^m	-82006 3	959.73 0.13	24.0 ns 0.7	11/2 ⁻	02		IT=100	
¹⁴³ Sm	-79523 4		8.75 m 0.08	3/2 ⁺	02		β^+ =100	
¹⁴³ Sm ^m	-78769 4	753.99 0.16	66 s 2	11/2 ⁻	02		IT \approx 100; β^+ =0.24 6	
¹⁴³ Sm ⁿ	-76729 4	2793.8 0.13	30 ms 3	23/2 ⁽⁻⁾	02		IT=100	
¹⁴³ Eu	-74242 11		2.59 m 0.02	5/2 ⁺	02		β^+ =100	
¹⁴³ Eu ^m	-73852 11	389.51 0.04	50.0 μ s 0.5	11/2 ⁻	02		IT=100	
¹⁴³ Gd	-68230 200		39 s 2	(1/2) ⁺	02	78Fi02 D	β^+ =100; β^+ p=?; β^+ α =? *	
¹⁴³ Gd ^m	-68080 200	152.6 0.5	110.0 s 1.4	(11/2 ⁻)	02	78Fi02 D	β^+ =100; β^+ p=?; β^+ α =? *	
¹⁴³ Tb	-60430 60		12 s 1	(11/2 ⁻)	01		β^+ =100	
¹⁴³ Tb ^m	-60430# 120#	0# 100#	* < 21 s	5/2 ⁺ #	01		β^+ ?	
¹⁴³ Dy	-52320# 200#		5.6 s 1.0	(1/2 ⁺)	01	03Xu04 TJ	β^+ =100; β^+ p=? *	
¹⁴³ Dy ^m	-52010# 200#	310.7 0.6	3.0 s 0.3	(11/2 ⁻)	01	03Xu04 JTD	β^+ =100; β^+ p=? *	
¹⁴³ Ho	-42280# 400#		300# ms (>200 ns)	11/2 ⁻ #	01	00So11 I	β^+ ?	
¹⁴³ Er	-31350# 600#		200# ms	9/2 ⁻ #			β^+ ?	
* ¹⁴³ Gd	D : 78Fi02: β^+ p and/or β^+ α for ¹⁴³ Gd+ ¹⁴³ Gd ^m =0.001%, 39 particles detected							**
* ¹⁴³ Dy	T : others: 84Ni03=3.2(0.6) 83Ni05=4.1(0.3) in two different experiments							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁴⁴ I	-46580# 500#		50# ms (>300 ns)	1 ⁻ #	01	94Be24 I	β^- ?; β^- n=40#	
¹⁴⁴ Xe	-57280# 300#		388 ms	0 ⁺	01	03Be05 TD	β^- =100; β^- n=3.0 3	
¹⁴⁴ Cs	-63270 26		994 ms	4	1 ^(-#)	01	β^- =100; β^- n=3.20 21	
¹⁴⁴ Cs ^m	-62970# 200#	300# 200#	< 1 s	(> 3)	01		β^- =?; IT ?	
¹⁴⁴ Ba	-71769 13		11.5 s	0.2	0 ⁺	01	β^- =100 *	
¹⁴⁴ La	-74890 50		40.8 s	0.4	(3 ⁻)	01	β^- =100	
¹⁴⁴ Ce	-80437 3		284.91 d	0.05	0 ⁺	01	β^- =100	
¹⁴⁴ Pr	-80756 3		17.28 m	0.05	0 ⁻	01	β^- =100	
¹⁴⁴ Pr ^m	-80697 3	59.03 0.03	7.2 m	0.3	3 ⁻	01	IT \approx 100; β^- =0.07	
¹⁴⁴ Nd	-83753.2 2.3		2.29 Py	0.16	0 ⁺	01	IS=23.8 3; α =100	
¹⁴⁴ Pm	-81421 3		363 d	14	5 ⁻	01	ϵ =100; e^+ <8e-5	
¹⁴⁴ Pm ^m	-80580 3	840.90 0.05	780 ns	200	(9 ⁺)	01	IT=100	
¹⁴⁴ Pm ⁿ	-72825 4	8595.8 2.2	2.7 μ s		(27 ⁺)	01	IT=100	
¹⁴⁴ Sm	-81972.0 2.8		STABLE		0 ⁺	01	IS=3.07 7; $2\beta^+$?	
¹⁴⁴ Sm ^m	-79648.4 2.8	2323.60 0.08	880 ns	25	6 ⁺	01	IT=100	
¹⁴⁴ Eu	-75622 11		10.2 s	0.1	1 ⁺	01	β^+ =100	
¹⁴⁴ Eu ^m	-74494 11	1127.6 0.6	1.0 μ s	0.1	(8 ⁻)	01	IT=100	
¹⁴⁴ Gd	-71760 28		4.47 m	0.06	0 ⁺	01	β^+ =100	
¹⁴⁴ Tb	-62368 28		1 s		1 ⁺	01	β^+ =100; β^+ p ?	
¹⁴⁴ Tb ^m	-61971 28	396.9 0.5	4.25 s	0.15	(6 ⁻)	01	IT=66; β^+ =34; β^+ p ?	
¹⁴⁴ Tb ⁿ	-61892 28	476.2 0.5	2.8 μ s	0.3	(8 ⁻)	01	IT=100	
¹⁴⁴ Tb ^p	-61851 28	517.1 0.5	670 ns	60	(9 ⁺)	01	IT=100	
¹⁴⁴ Dy	-56580 30		9.1 s	0.4	0 ⁺	01	β^+ =100; β^+ p=?	
¹⁴⁴ Ho	-45200# 300#		700 ms	100		01	β^+ =100; β^+ p=?	
¹⁴⁴ Er	-36910# 400#		400# ms (>200 ns)	0 ⁺	01	00So11 I	β^+ ?	
* ¹⁴⁴ Ba	D : β^- n=3.6 7 in ENSDF'01 belongs in fact to ¹⁴⁴ Cs							**
¹⁴⁵ Xe	-52100# 300#		188 ms	4	3/2 ⁻ #	97	β^- =100; β^- n=5.0 6	
¹⁴⁵ Cs	-60057 11		582 ms	6	3/2 ⁺	93	β^- =100; β^- n=14.3 8 *	
¹⁴⁵ Ba	-67410 70		4.31 s	0.16	5/2 ⁻	98	β^- =100	
¹⁴⁵ La	-72990 90		24.8 s	2.0	(5/2 ⁺)	98	β^- =100	
¹⁴⁵ Ce	-77100 40		3.01 m	0.06	(3/2 ⁻)	93	β^- =100	
¹⁴⁵ Pr	-79632 7		5.984 h	0.010	7/2 ⁺	93	β^- =100	
¹⁴⁵ Nd	-81437.1 2.3		STABLE		7/2 ⁻	93	IS=8.3 1	
¹⁴⁵ Pm	-81274 3		17.7 y	0.4	5/2 ⁺	93	ϵ =100; α =2.8e-7	
¹⁴⁵ Sm	-80657.7 2.8		340 d	3	7/2 ⁻	02	ϵ =100	
¹⁴⁵ Sm ^m	-71871.5 2.9	8786.2 0.7	990 ns	170	(49/2 ⁺)	02	IT=100	
¹⁴⁵ Eu	-77998 4		5.93 d	0.04	5/2 ⁺	93	β^+ =100	
¹⁴⁵ Eu ^m	-77282 4	716.0 0.3	490 ns		11/2 ⁻	93	IT=100	
¹⁴⁵ Gd	-72927 19		23.0 m	0.4	1/2 ⁺	01	β^+ =100	
¹⁴⁵ Gd ^m	-72178 19	749.1 0.2	85 s	3	11/2 ⁻	01	IT=94.3 5; β^+ =5.7 5	
¹⁴⁵ Tb	-65880 60		20# m		(3/2 ⁺)	96	β^+ ?	
¹⁴⁵ Tb ^m	-65880# 120#	0# 100#	* 30.9 s	0.7	(11/2 ⁻)	96	β^+ =100 *	
¹⁴⁵ Dy	-58290 50		9.5 s	1.0	(1/2 ⁺)	93	β^+ =100; β^+ p=? *	
¹⁴⁵ Dy ^m	-58170 50	118.2 0.2	14.1 s	0.7	(11/2 ⁻)	93	β^+ =100 *	
¹⁴⁵ Ho	-49180# 300#		* 2.4 s	0.1	(11/2 ⁻)	93	β^+ =100	
¹⁴⁵ Ho ^m	-49080# 320#	100# 100#	* 100# ms		5/2 ⁺ #		β^+ ?; IT ?	
¹⁴⁵ Er	-39690# 400#		900 ms	300	1/2 ⁺ #	98	β^+ =100; β^+ p=?	
¹⁴⁵ Tm	-27880# 400#		3.1 μ s	0.3	(11/2 ⁻)	02	p=100 *	
* ¹⁴⁵ Cs	T : average 93Ru01=579(6) 82Ra13=594(13)							**
* ¹⁴⁵ Tb ^m	T : average 93A103=31.6(0.6) 82No08=29.5(1.0) and 82A107=29.5(1.5)							**
* ¹⁴⁵ Dy	T : average 93A103=10.5(1.5) 93To04=6(2) and 84Sc.C=10(1)							**
* ¹⁴⁵ Dy ^m	T : average 93To04=14.5(1.0) 82No08=13.6(1.0)							**
* ¹⁴⁵ Tm	T : average 03Ka04=3.1(0.3) 98Ba13=3.5(1.0) J : not adopted by ENSDF'02							**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁴⁶ Xe	-48670#	400#			146 ms	6	0 ⁺	97 03Be05	TD β^- =100; β^- n=6.9 15
¹⁴⁶ Cs	-55620	70			323 ms	6	1 ⁻	97 93Ru01	T β^- =100; β^- n=14.2 5 *
¹⁴⁶ Ba	-65000	70			2.22 s	0.07	0 ⁺	97 93Ru01	D β^- =100 *
¹⁴⁶ La	-69120	70			6.27 s	0.10	2 ⁻	97 93Ru01	D β^- =100 *
¹⁴⁶ La ^m	-68990	150	130		10.0 s	0.1	(6 ⁻)	97 79Ke02	E β^- =100 *
¹⁴⁶ Ce	-75680	70			13.52 m	0.13	0 ⁺	97	β^- =100
¹⁴⁶ Pr	-76710	60			24.15 m	0.18	(2) ⁻	97	β^- =100
¹⁴⁶ Nd	-80931.1	2.3			STABLE		0 ⁺	97	IS=17.2 3; $2\beta^-$?; α ?
¹⁴⁶ Pm	-79460	5			5.53 y	0.05	3 ⁻	99	ϵ =66.0 13; β^- =34.0 13
¹⁴⁶ Sm	-81002	4			103 My	5	0 ⁺	97	α =100
¹⁴⁶ Eu	-77122	6			4.61 d	0.03	4 ⁻	97	β^+ =100
¹⁴⁶ Eu ^m	-76456	6	666.37	0.16	235 μ s	3	9 ⁺	97	IT=100
¹⁴⁶ Gd	-76093	5			48.27 d	0.10	0 ⁺	01	ϵ =100
¹⁴⁶ Tb	-67770	50			8 s	4	1 ⁺	97	β^+ =100
¹⁴⁶ Tb ^m	-67620#	110#	150#	100#	24.1 s	0.5	5 ⁻	97 93Al03	T β^+ =100
¹⁴⁶ Tb ⁿ	-66840#	110#	930#	100#	1.18 ms	0.02	(10 ⁺)	97	IT=100 *
¹⁴⁶ Dy	-62554	27			33.2 s	0.7	0 ⁺	97 93Al03	T β^+ =100
¹⁴⁶ Dy ^m	-59618	27	2935.7	0.6	150 ms	20	10 ⁺ #	97	IT=100
¹⁴⁶ Ho	-51570#	200#			3.6 s	0.3	(10 ⁺)	97	β^+ =100; β^+ p=?
¹⁴⁶ Er	-44710#	300#			1.7 s	0.6	0 ⁺	97 93To05	D β^+ =100; β^+ p=?
¹⁴⁶ Tm	-31280#	400#			240 ms	30	(6 ⁻)	02	p≈100; β^+ ?
¹⁴⁶ Tm ^m	-31200#	400#	71	6	72 ms	23	(10 ⁺)	02	p=?; β^+ =16#
* ¹⁴⁶ Cs	T : average 93Ru01=321(2) 76Lu02=343(7) **								
* ¹⁴⁶ Ba	D : 93Ru01 β^- n<0.02% is not relevant since $Q(\beta^-$ n) is negative: =-190(100) **								
* ¹⁴⁶ La	D : 93Ru01 β^- n<0.007% is not relevant since $Q(\beta^-$ n) is negative: =-180(80) **								
* ¹⁴⁶ La ^m	E : derived from $Q(^{146}\text{La}^m)$ =6660(120) in 79Ke02 **								
* ¹⁴⁶ Tb ⁿ	E : 779.6 keV above ¹⁴⁶ Tb ^m , from ENSDF **								
¹⁴⁷ Xe	-43260#	400#			130 ms	80	3/2 ⁻ #	98 03Be05	TD β^- =100; β^- n=4.0 23 *
¹⁴⁷ Cs	-52020	50			225 ms	5	(3/2 ⁺)	92 93Ru01	D β^- =100; β^- n=28.5 17
¹⁴⁷ Ba	-60600#	210#			893 ms	1	(3/2 ⁺)	98 93Ru01	D β^- =100 *
¹⁴⁷ La	-66850	50			4.015 s	0.008	(5/2 ⁺)	98 93Ru01	D β^- =100; β^- n=0.040 3 *
¹⁴⁷ Ce	-72030	30			56.4 s	1.0	(5/2 ⁻)	92	β^- =100
¹⁴⁷ Pr	-75455	23			13.4 m	0.4	(3/2 ⁺)	92	β^- =100
¹⁴⁷ Nd	-78151.9	2.3			10.98 d	0.01	5/2 ⁻	92	β^- =100
¹⁴⁷ Pm	-79047.9	2.4			2.6234 y	0.0002	7/2 ⁺	96	β^- =100
¹⁴⁷ Sm	-79272.1	2.4			106.0 Gy	1.1	7/2 ⁻	92 70Gu14	T IS=14.99 18; α =100 *
¹⁴⁷ Eu	-77550	3			24.1 d	0.6	5/2 ⁺	99	β^+ ≈100; α =0.0022 6
¹⁴⁷ Gd	-75363	3			38.06 h	0.12	7/2 ⁻	99	β^+ =100
¹⁴⁷ Gd ^m	-66775	3	8587.8	0.4	510 ns	20	(49/2 ⁺)	99	IT=100
¹⁴⁷ Tb	-70752	12			1.64 h	0.03	1/2 ⁺ #	99 97Wa04	T β^+ =100
¹⁴⁷ Tb ^m	-70701	12	50.6	0.9	1.87 m	0.05	(11/2) ⁻	99 93Al03	T β^+ =100 *
¹⁴⁷ Dy	-64188	20			40 s	10	1/2 ⁺	92 84To07	D β^+ =100; β^+ p≈0.05
¹⁴⁷ Dy ^m	-63438	20	750.5	0.4	55 s	1	11/2 ⁻	92	β^+ =65 4; IT=35 4
¹⁴⁷ Ho	-55837	28			5.8 s	0.4	(11/2 ⁻)	92	β^+ =100; β^+ p ?
¹⁴⁷ Er	-47050#	300#			2.5 s		(1/2 ⁺)	92	β^+ =100; β^+ p=?
¹⁴⁷ Er ^m	-46950#	300#	100#	50#	2.5 s	0.2	(11/2 ⁻)	92	β^+ =100 *
¹⁴⁷ Tm	-36370#	300#			580 ms	30	11/2 ⁻	02	β^+ =85 5; p=15 5
¹⁴⁷ Tm ^m	-36300#	300#	60	5	360 μ s	40	3/2 ⁺	02	p=100
* ¹⁴⁷ Xe	D : from β^- n<8% **								
* ¹⁴⁷ Ba	D : 93Ru01 β^- n=0.06(3)% contradicts $Q(\beta^-$ n)=-340(120) **								
* ¹⁴⁷ La	J : from 96Ur02 **								
* ¹⁴⁷ Sm	T : average 70Gu14=106(2) 65Va16=108(2) 64Do01=104(3) 61Wr02=105(2) **								
* ¹⁴⁷ Tb ^m	T : average 93Al03=1.92(0.07) 73Bo13=1.83(0.06) E : from 87Li09 **								
* ¹⁴⁷ Er ^m	E : estimated from 11/2 ⁻ level in isotones ¹⁴¹ Sm=175 ¹⁴³ Gd=152 ¹⁴⁵ Dy=118 **								

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{148}Cs	-47300	580			146 ms	6		00		β^- =100; β^- -n=25.1 25
^{148}Ba	-58010	80			612 ms	17	0^+	00		β^- =100; β^- -n=0.4 3
^{148}La	-63130	60			1.26 s	0.08	(2^-)	00		β^- =100; β^- -n=0.15 3
^{148}Ce	-70391	29			56 s	1	0^+	00		β^- =100
^{148}Pr	-72531	26			2.29 m	0.02	1^-	00		β^- =100
$^{148}\text{Pr}^m$	-72480#	40#	50#	30#	* 2.01 m	0.07	(4)	00	ABBW E	β^- =100 *
^{148}Nd	-77413.4	2.8			STABLE	(>3.0 Ey)	0^+	00	82Be20 T	IS=5.7 1; $2\beta^-$?; α ?
^{148}Pm	-76872	6			5.368 d	0.002	1^-	00		β^- =100
$^{148}\text{Pm}^m$	-76734	6	137.9	0.3	41.29 d	0.11	$5^-, 6^-$	00		β^- =95.8 6; IT=4.2 6
^{148}Sm	-79342.2	2.4			7 Py	3	0^+	00		IS=11.24 10; α =100
^{148}Eu	-76302	10			54.5 d	0.5	5^-	00		β^+ =100; α =9.4e-7 28
^{148}Gd	-76275.8	2.8			74.6 y	3.0	0^+	00		α =100; $2\beta^+$?
^{148}Tb	-70540	14			60 m	1	2^-	00		β^+ =100
$^{148}\text{Tb}^m$	-70450	14	90.1	0.3	2.20 m	0.05	$(9)^+$	00		β^+ =100
$^{148}\text{Tb}^n$	-61921	14	8618.6	1.0	1.310 μs	0.007	(27^+)	00		IT=100
^{148}Dy	-67859	11			3.3 m	0.2	0^+	00		β^+ =100
^{148}Ho	-58020	130			2.2 s	1.1	(1^+)	00		β^+ =100
$^{148}\text{Ho}^m$	-57620#	160#	400#	100#	9.49 s	0.12	$(6)^-$	00	93Al03 T	β^+ =100; β^+ p=0.08 1 *
$^{148}\text{Ho}^n$	-57330#	160#	690#	100#	2.35 ms	0.04	(10^+)	00		IT=100 *
^{148}Er	-51650#	200#			4.6 s	0.2	0^+	00		β^+ =100; β^+ p \approx 0.15
^{148}Tm	-39270#	400#			700 ms	200	(10^+)	00		β^+ =100
^{148}Yb	-30350#	600#			250# ms		0^+			β^+ ?
* $^{148}\text{Pr}^m$	E : derived from ENSDF estimate $E < 90$ keV **									
* $^{148}\text{Ho}^m$	T : average $93\text{Al}03=9.30(0.20)$ $89\text{Ta}11=9.59(0.15)$ **									
* $^{148}\text{Ho}^n$	E : 694.4 keV above $^{148}\text{Ho}^m$, from ENSDF **									
^{149}Cs	-43850#	200#			150# ms	(>50 ms)	$3/2^+\#$	95	87Ra12 I	β^- ?; β^- -n ?
^{149}Ba	-53490#	200#			344 ms	7	$3/2^- \#$	95		β^- =100; β^- -n=0.43 12
^{149}La	-60800#	320#			1.05 s	0.03	$5/2^+\#$	95	93Ru01 D	β^- =100; β^- -n=1.4 3
^{149}Ce	-66700	100			5.3 s	0.2	$3/2^- \#$	98		β^- =100
^{149}Pr	-71060	80			2.26 m	0.07	$(5/2^+)$	95		β^- =100
^{149}Nd	-74380.9	2.8			1.728 h	0.001	$5/2^-$	95		β^- =100
^{149}Pm	-76071	4			53.08 h	0.05	$7/2^+$	95		β^- =100
$^{149}\text{Pm}^m$	-75831	4	240.214	0.007	35 μs	3	$11/2^-$			
^{149}Sm	-77141.9	2.4			STABLE	(>2 Py)	$7/2^-$	95		IS=13.82 7; α ?
^{149}Eu	-76447	4			93.1 d	0.4	$5/2^+$	95		ϵ =100
^{149}Gd	-75133	4			9.28 d	0.10	$7/2^-$	01		β^+ =100; α =4.3e-4 10
^{149}Tb	-71496	4			4.118 h	0.025	$1/2^+$	99		β^+ =83.3 17; α =16.7 17
$^{149}\text{Tb}^m$	-71460	4	35.78	0.13	4.16 m	0.04	$11/2^-$	99		β^+ \approx 100; α =0.022 3
^{149}Dy	-67715	9			4.20 m	0.14	$7/2^{(-)}$	95	88Ah02 J	β^+ =100
$^{149}\text{Dy}^m$	-65054	9	2661.1	0.4	490 ms	15	$(27/2^-)$	95		IT=99.3 3; β^+ =0.7 3
$^{149}\text{Dy}^n$	-60230	30	7490	30	28 ns	2	$(47/2^+)$	95		IT=100 *
^{149}Ho	-61688	18			21.1 s	0.2	$(11/2^-)$	95		β^+ =100
$^{149}\text{Ho}^m$	-61639	18	48.80	0.20	56 s	3	$(1/2^+)$	95		β^+ =100
^{149}Er	-53742	28			4 s	2	$(1/2^+)$	95		β^+ =100; β^+ p=7 2
$^{149}\text{Er}^m$	-53000	28	741.8	0.2	8.9 s	0.2	$(11/2^-)$	95		β^+ =96.5 7; IT=3.5 7;... *
^{149}Tm	-44040#	300#			900 ms	200	$(11/2^-)$	95		β^+ =100; β^+ p=0.26 15
^{149}Yb	-33500#	500#			700 ms	200	$(1/2^+, 3/2^+)$	95	01Xu06 TD	β^+ =100; β^+ p=?
* $^{149}\text{Dy}^n$	E : 7409.9 above level at ≈ 80 keV **									
* $^{149}\text{Er}^m$	D : ... ; β^+ p=0.18 7 **									
^{150}Cs	-38960#	300#			100# ms	(>50 ms)		97	87Ra12 I	β^- ?; β^- -n ?
^{150}Ba	-50600#	400#			300 ms		0^+	95		β^- =100; β^- -n ?
^{150}La	-57040#	400#			510 ms	30	(3^+)	97	95Ok02 TJ	β^- =100; β^- -n=2.7 3
^{150}Ce	-64820	50			4.0 s	0.6	0^+	95		β^- =100
^{150}Pr	-68304	26			6.19 s	0.16	$(1)^-$	96		β^- =100
^{150}Nd	-73690	3			6.7 Ey	0.7	0^+	96	97De40 TD	IS=5.6 2; $2\beta^-$ =100 *
^{150}Pm	-73603	20			2.68 h	0.02	(1^-)	95		β^- =100

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁵² Ba	-42600#	500#		100# ms	0 ⁺	97		β^- ?
¹⁵² La	-50070#	400#		200# ms (>300 ns)		97	94Be24 I	β^- ?
¹⁵² Ce	-59110#	200#		1.1 s	0.3	0 ⁺	97 90Ta07 T	β^- =100 *
¹⁵² Pr	-63810	120		3.63 s	0.12	4 ⁺	97 99To04 J	β^- =100
¹⁵² Nd	-70158	25		11.4 m	0.2	0 ⁺	97	β^- =100
¹⁵² Pm	-71262	26		* 4.12 m	0.08	1 ⁺	97	β^- =100
¹⁵² Pm ^m	-71120	80	140	BD * 7.52 m	0.08	4 ⁻	97	β^- =100
¹⁵² Pm ⁿ	-71010#	150#	250#	* 13.8 m	0.2	(8)	97	β^- ≈100; IT=? *
¹⁵² Sm	-74768.8	2.5		STABLE		0 ⁺	97	IS=26.75 16
¹⁵² Eu	-72894.5	2.5		13.537 y	0.006	3 ⁻	97	β^+ =72.1 3; β^- =27.9 3
¹⁵² Eu ^m	-72848.9	2.5	45.5998	9.3116 h	0.0013	0 ⁻	97	β^- =72 4; β^+ =28 4
¹⁵² Eu ⁿ	-72746.6	2.5	147.86	96 m	1	8 ⁻	97	IT=100
¹⁵² Gd	-74714.2	2.5		108 Ty	8	0 ⁺	97	IS=0.20 1; α =100; 2 β^+ ?
¹⁵² Tb	-70720	40		17.5 h	0.1	2 ⁻	98	β^+ =100; α <7e-7
¹⁵² Tb ^m	-70220	40	501.74	4.2 m	0.1	8 ⁺	98	IT=78.8 8; β^+ =21.2 8
¹⁵² Dy	-70124	5		2.38 h	0.02	0 ⁺	99	ϵ ≈100; α =0.100 7
¹⁵² Ho	-63608	14		161.8 s	0.3	2 ⁻	97	β^+ =88 3; α =12 3
¹⁵² Ho ^m	-63448	14	160	50.0 s	0.4	9 ⁺	97	β^+ =89.2 17; α =10.8 17
¹⁵² Ho ⁿ	-60588	14	3019.59	8.4 μs	0.3	19 ⁻	97	IT=100
¹⁵² Er	-60500	11		10.3 s	0.1	0 ⁺	97	α =90 4; β^+ =10 4
¹⁵² Tm	-51770	70		* 8.0 s	1.0	(2#) ⁻	97	β^+ =100
¹⁵² Tm ^m	-51670#	110#	100#	* 5.2 s	0.6	(9) ⁺	97	β^+ =100
¹⁵² Yb	-46310	210		3.04 s	0.06	0 ⁺	97	β^+ =100; β^+ p ?
¹⁵² Lu	-33420#	200#		650 ms	70	(5 ⁻ , 6 ⁻)	97 88Ni02 T	β^+ =100; β^+ p=15 7 *
* ¹⁵² Ce	T : average 90Ta07=1.4(0.2) 91Ay.A=0.8(0.3) **							
* ¹⁵² Pm ⁿ	E : ENSDF: "Probably feeds 7.52 m level" at 140 keV **							
* ¹⁵² Lu	T : average 88Ni02=600(100) 87To02=700(100) **							
¹⁵³ Ba	-37620#	800#		80# ms		5/2 ⁻ #		β^- ?
¹⁵³ La	-46930#	600#		150# ms (>300 ns)		5/2 ⁺ #	98 94Be24 I	β^- ?
¹⁵³ Ce	-55350#	400#		500# ms (>300 ns)		3/2 ⁻ #	98 94Be24 I	β^- ?
¹⁵³ Pr	-61630	100		4.28 s	0.11	5/2 ⁻ #	98	β^- =100
¹⁵³ Nd	-67349	27		31.6 s	1.0	(3/2) ⁻	98	β^- =100
¹⁵³ Pm	-70685	11		5.25 m	0.02	5/2 ⁻	98	β^- =100
¹⁵³ Sm	-72565.8	2.5		46.284 h	0.004	3/2 ⁺	98	β^- =100 *
¹⁵³ Sm ^m	-72467.4	2.5	98.37	10.6 ms	0.3	11/2 ⁻	98	IT=100
¹⁵³ Eu	-73373.5	2.5		STABLE		5/2 ⁺	98	IS=52.19 3
¹⁵³ Gd	-72889.8	2.5		240.4 d	1.0	3/2 ⁻	98	ϵ =100
¹⁵³ Gd ^m	-72794.6	2.5	95.1737	3.5 μs	0.4	(9/2 ⁺)	98	IT=100
¹⁵³ Gd ⁿ	-72718.6	2.5	171.189	76.0 μs	1.4	(11/2 ⁻)	98	IT=100
¹⁵³ Tb	-71320	4		2.34 d	0.01	5/2 ⁺	98	β^+ =100
¹⁵³ Tb ^m	-71157	4	163.175	186 μs	4	11/2 ⁻	98	IT=100
¹⁵³ Dy	-69150	5		6.4 h	0.1	7/2 ⁽⁻⁾	99	β^+ ≈100; α =0.0094 14
¹⁵³ Ho	-65019	6		2.01 m	0.03	11/2 ⁻	98	β^+ ≈100; α =0.051 25
¹⁵³ Ho ^m	-64950	6	68.7	9.3 m	0.5	1/2 ⁺	98	β^+ ≈100; α =0.18 8
¹⁵³ Er	-60488	9		37.1 s	0.2	7/2 ⁽⁻⁾	98 85Ah.1 J	α =53 3; β^+ =47 3 *
¹⁵³ Tm	-54015	18		1.48 s	0.01	(11/2 ⁻)	98	α =91 3; β^+ =9 3
¹⁵³ Tm ^m	-53972	18	43.2	2.5 s	0.2	(1/2 ⁺)	98	α =92 3; β^+ =?
¹⁵³ Yb	-47060#	200#		4.2 s	0.2	7/2 ⁻ #	98 88Wi05 D	β^+ =?; α =50#; ... *
¹⁵³ Yb ^m	-44360#	220#	2700	15 μs	1	(27/2 ⁻)	98	*
¹⁵³ Lu	-38410	210		900 ms	200	11/2 ⁻	98 97Ir01 D	α =70#; β^+ =?; p=0 *
¹⁵³ Lu ^m	-38330	210	80	1# s		1/2 ⁺	98 97Ir01 ED	β^+ ?; α ?; p=0 *
¹⁵³ Lu ⁿ	-35780	210	2632.9	15 μs	3	27/2 ⁻	98	
¹⁵³ Hf	-27300#	500#		400# ms (>200 ns)		1/2 ⁺ #	00So11 I	β^+ ?
¹⁵³ Hf ^m	-26550#	510#	750#	500# ms		11/2 ⁻ #		β^+ ?; IT ?
* ¹⁵³ Sm	T : see also 99Sc12=46.274(7) **							
* ¹⁵³ Er	J : and 89Ot.A **							
* ¹⁵³ Yb	D : ... ; β^+ p=0.008 2 **							
* ¹⁵³ Yb ^m	E : in ENSDF 2578.2 + x **							
* ¹⁵³ Lu	D : p decay is from 97Ir01 **							

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)		
¹⁵⁴ La	-42380#	600#		100#	ms				β^- ?		
¹⁵⁴ Ce	-52700#	500#		300#	ms (>300 ns)	0 ⁺	98	94Be24 I	β^- ?		
¹⁵⁴ Pr	-58200	150		2.3	s	0.1	(3 ⁺ , 2 ⁺)	98	β^- =100		
¹⁵⁴ Nd	-65690	110		25.9	s	0.2	0 ⁺	98	β^- =100		
¹⁵⁴ Nd ^m	-65210#	190#	480#	150#	1.3	μ s	0.5		98		
¹⁵⁴ Nd ⁿ	-64340	110	1349	10	>1	μ s	(5 ⁻)	98			
¹⁵⁴ Pm	-68500	40			* & 1.73	m	0.10	(0, 1)	98	β^- =100	
¹⁵⁴ Pm ^m	-68380	110	120	120	BD * & 2.68	m	0.07	(3, 4)	98	β^- =100	
¹⁵⁴ Sm	-72461.6	2.5			STABLE	(>2.3 Ey)	0 ⁺		98	IS=22.75 29; 2 β^- ?	
¹⁵⁴ Eu	-71744.4	2.5			8.593	y	0.004	3 ⁻	98	β^- ≈100; ϵ =0.02 1	
¹⁵⁴ Eu ^m	-71599.1	2.5	145.3	0.3	46.3	m	0.4	(8 ⁻)	98	IT=100	
¹⁵⁴ Gd	-73713.2	2.5			STABLE		0 ⁺		98	IS=2.18 3	
¹⁵⁴ Tb	-70160	50			*	21.5	h	0.4	0 ⁽⁺⁾ #	98	β^- ≈100; β^- <0.1
¹⁵⁴ Tb ^m	-70150	50	12	7	*	9.4	h	0.4	3 ⁻	98	ABBW E β^+ =78.2 7; IT=21.8 7;...
¹⁵⁴ Tb ⁿ	-69960#	160#	200#	150#	*	22.7	h	0.5	7 ⁻	98	β^+ =98.2 6; IT=1.8 6
¹⁵⁴ Dy	-70398	8				3.0	My	1.5	0 ⁺	99	α =100; 2 β^+ ?
¹⁵⁴ Ho	-64644	8				11.76	m	0.19	2 ⁻	98	β^+ ≈100; α =0.019 5
¹⁵⁴ Ho ^m	-64406	28	238	30	AD	3.10	m	0.14	8 ⁺	98	β^+ =100; α <0.001; IT≈0
¹⁵⁴ Er	-62612	5				3.73	m	0.09	0 ⁺	01	β^+ ≈100; α =0.47 13
¹⁵⁴ Tm	-54429	14			*	8.1	s	0.3	(2 ⁻)	98	α =54 5; β^+ =46 5
¹⁵⁴ Tm ^m	-54360	50	70	50	BD *	3.30	s	0.07	(9 ⁺)	98	α =58 5; β^+ =42 5
¹⁵⁴ Yb	-49934	17				409	ms	2	0 ⁺	98	α =92.6 12; β^+ =7.4 12
¹⁵⁴ Lu	-39570#	200#				1#	s		(2 ⁻)	98	β^+ ?
¹⁵⁴ Lu ^m	-39510#	200#	58	13	AD	1.12	s	0.08	(9 ⁺)	98	88Vi02 D β^+ ≈100; β^+ p=?: ...
¹⁵⁴ Lu ⁿ	-37300#	600#	> 2562			35	μ s	3	(17 ⁺)	98	IT=100
¹⁵⁴ Hf	-32730#	500#				2	s	1	0 ⁺	98	β^+ ≈100; α ≈0
* ¹⁵⁴ Tb ^m	D : ... ; β^- <0.1									**	
* ¹⁵⁴ Tb ⁿ	E : less than 25 keV, from ENSDF									**	
* ¹⁵⁴ Tm ^m	D : IT decay has not been observed									**	
* ¹⁵⁴ Lu ^m	D : ... ; β^+ α =?: α =0.002#									**	
* ¹⁵⁴ Lu ⁿ	D : β^+ p and β^+ α modes observed by 88Vi02; β^+ p confirmed by 90Sh.A									**	
¹⁵⁵ La	-38800#	800#			60#	ms		5/2 ⁺ #		β^- ?	
¹⁵⁵ Ce	-48400#	600#			200#	ms (>300 ns)		5/2 ⁻ #	97	94Be24 I β^- ?	
¹⁵⁵ Pr	-55780#	300#			1#	s (>300 ns)		5/2 ⁻ #	97	95Cz.A I β^- ?	
¹⁵⁵ Nd	-62470#	150#			8.9	s	0.2	3/2 ⁻ #	94	β^- =100	
¹⁵⁵ Pm	-66970	30			41.5	s	0.2	(5/2 ⁻)	94	β^- =100	
¹⁵⁵ Sm	-70197.2	2.6			22.3	m	0.2	3/2 ⁻	94	β^- =100	
¹⁵⁵ Eu	-71824.5	2.5			4.7611	y	0.0013	5/2 ⁺	94	β^- =100	
¹⁵⁵ Gd	-72077.1	2.5			STABLE			3/2 ⁻	97	IS=14.80 12	
¹⁵⁵ Gd ^m	-71956.1	2.5	121.05	0.19	32.0	ms	0.3	11/2 ⁻	97	IT=100	
¹⁵⁵ Tb	-71254	12			5.32	d	0.06	3/2 ⁺	94	ϵ =100	
¹⁵⁵ Dy	-69160	12			9.9	h	0.2	3/2 ⁻	99	β^+ =100	
¹⁵⁵ Dy ^m	-68926	12	234.33	0.03	6	μ s		11/2 ⁻	99	IT=100	
¹⁵⁵ Ho	-66040	18			48	m	1	5/2 ⁺	94	β^+ =100	
¹⁵⁵ Ho ^m	-65898	18	141.97	0.11	880	μ s	80	11/2 ⁻	94	IT=100	
¹⁵⁵ Er	-62215	7			5.3	m	0.3	7/2 ⁻	94	β^+ ≈100; α =0.022 7	
¹⁵⁵ Tm	-56635	13			21.6	s	0.2	(11/2 ⁻)	95	β^+ =98.1 3; α =1.9 3	
¹⁵⁵ Tm ^m	-56594	14	41	6	45	s	3	(1/2 ⁺)	95	β^+ >92; α <8	
¹⁵⁵ Yb	-50503	17			1.793	s	0.019	(7/2 ⁻)	94	96Pa01 T α =89 4; β^+ =11 4	
¹⁵⁵ Lu	-42554	20			& 68.6	ms	1.6	(11/2 ⁻)	94	97Da07 TD α =88 4; β^+ ?	
¹⁵⁵ Lu ^m	-42534	21	20	6	& 138	ms	8	(1/2 ⁺)	94	97Da07 TJD α =76 16; β^+ ?	
¹⁵⁵ Lu ⁿ	-40773	20	1781.0	2.0	AD	2.70	ms	0.03	(25/2 ⁻)	94	96Pa01 T α ≈100; IT ?
¹⁵⁵ Hf	-34100#	400#			890	ms	120	7/2 ⁻ #	94	β^+ ≈100; α ?	
¹⁵⁵ Ta	-23670#	500#			13	μ s	4	(11/2 ⁻)	02	p=100	
* ¹⁵⁵ Yb	T : average 96Pa01=1.80(0.02) 91To08=1.75(0.05)									**	
* ¹⁵⁵ Lu	T : average 96Pa01=70(1) 97Da07=63(2) 91To09=66(7) 79Ho10=70(6)									**	
* ¹⁵⁵ Lu	D : α : average 97Da07=90(2)% 79Ho10=79(4)% with Birge ratio B=4.4									**	
* ¹⁵⁵ Lu ^m	T : average 97Da07=150(24) 96Pa01=136(9) 91To09=140(20)									**	
* ¹⁵⁵ Lu ⁿ	T : average 96Pa01=2.71(0.03) 81Ho.A=2.62(0.07)									**	

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)		
¹⁵⁶ Ce	-45400#	600#		150#	ms	0 ⁺			β^- ?		
¹⁵⁶ Pr	-51910#	400#		500#	ms (>300 ns)			95Cz.A I	β^- ?		
¹⁵⁶ Nd	-60530	200		5.49	s	0.07	0 ⁺	03	β^- =100		
¹⁵⁶ Nd ^m	-59100	200	1432	5	135	ns	5 ⁻	03	IT=100		
¹⁵⁶ Pm	-64220	30		26.70	s	0.10	4 ⁻	03	β^- =100		
¹⁵⁶ Sm	-69370	10		9.4	h	0.2	0 ⁺	03	β^- =100		
¹⁵⁶ Sm ^m	-67972	10	1397.55	0.09	185	ns	7	5 ⁻	03	IT=100	
¹⁵⁶ Eu	-70093	6		15.19	d	0.08	0 ⁺	03	β^- =100		
¹⁵⁶ Gd	-72542.2	2.5		STABLE			0 ⁺	03	IS=20.47 9		
¹⁵⁶ Gd ^m	-70404.6	2.5	2137.60	0.05	1.3	μ s	0.1	7 ⁻	03	IT=100	
¹⁵⁶ Tb	-70098	4		5.35	d	0.10	3 ⁻	03	β^+ \approx 100; β^- ?		
¹⁵⁶ Tb ^m	-70044	5	54	3	24.4	h	1.0	(7 ⁻)	03	IT=100 *	
¹⁵⁶ Tb ⁿ	-70010	4	88.4	0.2	5.3	h	0.2	(0 ⁺)	03	IT=?; β^+ =?	
¹⁵⁶ Dy	-70530	7		STABLE		(>1 Ey)	0 ⁺	03	58Ri23 T	IS=0.06 1; α ?; $2\beta^+$? *	
¹⁵⁶ Ho	-65350	40		56	m	1	4 ⁻	03		β^+ =100	
¹⁵⁶ Ho ^m	-65300	40	52.4	0.5	9.5	s	1.5	1 ⁻	03	IT=?; β^+ ?	
¹⁵⁶ Ho ⁿ	-65250#	60#	100#	50#	7.8	m	0.3	(9 ⁺)	03	β^+ =75; IT ?	
¹⁵⁶ Er	-64213	24		19.5	m	1.0	0 ⁺	03		β^+ =100; $\alpha=17e-6$ 4	
¹⁵⁶ Tm	-56840	16		83.8	s	1.8	2 ⁻	03		β^+ \approx 100; $\alpha=0.064$ 10	
¹⁵⁶ Tm ^m	-56636	16	203.6	0.5	400	ns		(11 ⁻)	03	IT=100	
¹⁵⁶ Tm ⁿ			non existent	RN	19	s	3	9 ⁺	03	91To08 I *	
¹⁵⁶ Yb	-53264	11		26.1	s	0.7	0 ⁺	03		β^+ =90 2; $\alpha=10$ 2	
¹⁵⁶ Lu	-43750	70		* 494	ms	12	(2 ⁻)	03		α =?; β^+ =5#	
¹⁵⁶ Lu ^m	-43530#	110#	220#	80#	* 198	ms	2	(9 ⁺)	03	96Pa01 D	$\alpha=94$ 6; β^+ ? *
¹⁵⁶ Hf	-37850	210		23	ms	1	0 ⁺	03	96Pa01 D	$\alpha=97$ 3; β^+ ? *	
¹⁵⁶ Hf ^m	-35890	210	1959.0	1.0	AD 480	μ s	40	8 ⁺	03	96Pa01 T	$\alpha=100$ *
¹⁵⁶ Ta	-25800#	400#		144	ms	24	(2 ⁻)	03		p \approx 100; β^+ ?	
¹⁵⁶ Ta ^m	-25700#	400#	100	8	AD 360	ms	40	(9 ⁺)	03	β^+ =95.8 9; p=4.2 9 *	
* ¹⁵⁶ Tb ^m										E: derived from E3 24h to 4 ⁺ 49.630 level and $E(IT)< B(L)=9$ keV **	
* ¹⁵⁶ Dy										T: lower limit is for α decay **	
* ¹⁵⁶ Tm ⁿ										I: see also the discussion in ENSDF'03 **	
* ¹⁵⁶ Lu ^m										D: derived from original $\alpha=98(9)\%$ **	
* ¹⁵⁶ Hf										D: derived from original $\alpha=100(6)\%$ **	
* ¹⁵⁶ Hf ^m										T: average 96Pa01=520(10) 81Ho.A=444(17) **	
* ¹⁵⁶ Ta ^m										T: 96Pa01=375(54) 93Li34=320(80) **	
¹⁵⁷ Ce	-40670#	700#		50#	ms		7/2 ⁺ #			β^- ?	
¹⁵⁷ Pr	-48970#	400#		300#	ms		5/2 ⁻ #			β^- ?	
¹⁵⁷ Nd	-56790#	200#		2#	s	(>300 ns)	5/2 ⁻ #	97	95Cz.A I	β^- ?	
¹⁵⁷ Pm	-62370	110		10.56	s	0.10	(5/2 ⁻)	96		β^- =100	
¹⁵⁷ Sm	-66730	50		8.03	m	0.07	(3/2 ⁻)	96		β^- =100	
¹⁵⁷ Eu	-69467	5		15.18	h	0.03	5/2 ⁺	96		β^- =100	
¹⁵⁷ Gd	-70830.7	2.5		STABLE			3/2 ⁻	96		IS=15.65 2	
¹⁵⁷ Tb	-70770.6	2.5		71	y	7	3/2 ⁺	96		$\varepsilon=100$	
¹⁵⁷ Dy	-69428	7		8.14	h	0.04	3/2 ⁻	97		β^+ =100	
¹⁵⁷ Dy ^m	-69229	7	199.38	0.07	21.6	ms	1.6	11/2 ⁻	97	IT=100 *	
¹⁵⁷ Ho	-66829	24		12.6	m	0.2	7/2 ⁻	96		β^+ =100	
¹⁵⁷ Er	-63420	28		18.65	m	0.10	3/2 ⁻	96		β^+ =100	
¹⁵⁷ Er ^m	-63265	28	155.4	0.3	76	ms	6	(9/2 ⁺)	96	IT=100	
¹⁵⁷ Tm	-58709	28		3.63	m	0.09	1/2 ⁺	97		β^+ =100	
¹⁵⁷ Yb	-53442	10		38.6	s	1.0	7/2 ⁻	96		β^+ =99.5; $\alpha=0.5$	
¹⁵⁷ Lu	-46483	19		6.8	s	1.8	(1/2 ⁺ , 3/2 ⁺)	96		β^+ ?; α =? *	
¹⁵⁷ Lu ^m	-46462	19	21.0	2.0	AD 4.79	s	0.12	(11/2 ⁻)	96	β^+ =?; $\alpha=6$ 2	
¹⁵⁷ Hf	-38750#	200#		115	ms	1	7/2 ⁻	96	96Pa01 T	$\alpha=86$ 9; β^+ =14 9	
¹⁵⁷ Ta	-29630	210		10.1	ms	0.4	1/2 ⁺	02		α =?; p=3.4 12; ... *	
¹⁵⁷ Ta ^m	-29610	210	22	5	4.3	ms	0.1	11/2 ⁻	02	α =?; β^+ =1#; p=0	
¹⁵⁷ Ta ⁿ	-28040	210	1593	9	AD 1.7	ms	0.1	(25/2 ⁻)	02	$\alpha=100$	
* ¹⁵⁷ Dy ^m										T: as adopted by ENSDF evaluator from 3 inconsistent results **	
* ¹⁵⁷ Lu										T: ENSDF'96 average of very discrepant 91To09=5.7(0.5) 91Le15,92Po14=9.6(8) **	
* ¹⁵⁷ Ta										D: ...; β^+ =1# **	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁵⁸ Pr	-44730# 600#		200# ms				β^- ?
¹⁵⁸ Nd	-54400# 400#		700# ms (>300 ns)	0 ⁺	97	95Cz.A I	β^- ?
¹⁵⁸ Pm	-59090 130		4.8 s	0.5	96		β^- =100
¹⁵⁸ Sm	-65210 80		5.30 m	0.03	96		β^- =100
¹⁵⁸ Eu	-67210 80		45.9 m	0.2	(1 ⁻)	96	β^- =100
¹⁵⁸ Gd	-70696.8 2.5		STABLE		96		IS=24.84 7
¹⁵⁸ Tb	-69477.2 2.6		180 y	11	3 ⁻	96	β^+ =83.4 7; β^- =16.6 7
¹⁵⁸ Tb ^m	-69366.9 2.9	110.3 1.2	10.70 s	0.17	0 ⁻	96	IT≈100; β^- <0.6; ... *
¹⁵⁸ Tb ⁿ	-69088.8 2.6	388.37 0.15	395 μs		7 ⁻		
¹⁵⁸ Dy	-70412 3		STABLE		0 ⁺	96	IS=0.10 1; α ?; 2 β^+ ?
¹⁵⁸ Ho	-66191 27		11.3 m	0.4	5 ⁺	97	β^+ ≈100; α ?
¹⁵⁸ Ho ^m	-66124 27	67.200 0.010	28 m	2	2 ⁻	97	IT>81; β^+ <19
¹⁵⁸ Ho ⁿ	-66010# 80#	180# 70#	21.3 m	2.3	(9 ⁺)	97	β^+ >93; IT<7#
¹⁵⁸ Er	-65304 25		2.29 h	0.06	0 ⁺	96	ε =100
¹⁵⁸ Tm	-58703 25		3.98 m	0.06	2 ⁻	96	β^+ =100
¹⁵⁸ Tm ^m	-58650# 100#	50# 100#	20 ns		(5 ⁺)	96 81Dr07 T	IT ? *
¹⁵⁸ Yb	-56015 8		1.49 m	0.13	0 ⁺	96	β^+ ≈100; α ≈0.0021 12
¹⁵⁸ Lu	-47214 15		10.6 s	0.3	2 ⁻	96 95Ga.A J	β^+ =99.09 20; ... *
¹⁵⁸ Hf	-42104 18		2.84 s	0.07	0 ⁺	96 96Pa01 TD	β^+ =55 3; α =45 3 *
¹⁵⁸ Ta	-31020# 200#		49 ms	8	(2 ⁻)	96 97Da07 TJD	α =96 4; β^+ ? *
¹⁵⁸ Ta ^m	-30880# 200#	140 12 AD	36.0 ms	0.8	(9 ⁺)	96 97Da07 TJE	α =93 6; β^+ ?; IT ? *
¹⁵⁸ W	-23700# 500#		1.37 ms	0.17	0 ⁺	96 00Ma95 T	α =100 *
¹⁵⁸ W ^m	-21810# 500#	1889 8 AD	143 μs	19	8 ⁺	00Ma95 T	α =100 *
* ¹⁵⁸ Tb ^m	D : ... ; β^+ <0.01 **						
* ¹⁵⁸ Tm ^m	I : T≈20 s in 81Dr07 was a typo. Value in Fig. 2 was correct. See 96Dr.A **						
* ¹⁵⁸ Lu	D : ... ; α =0.91 20 **						
* ¹⁵⁸ Hf	T : average 96Pa01=2.85(0.07) 73To02=2.8(0.2) **						
* ¹⁵⁸ Ta	T : average 97Da07=72(12) 96Pa01=46(4) with Birge ratio B=2 **						
* ¹⁵⁸ Ta	D : derived from original α ≈100(8)% **						
* ¹⁵⁸ Ta ^m	T : average 97Da07=37.7(1.5) 96Pa01=35(1) 79Ho10=36.8(1.6) **						
* ¹⁵⁸ W	T : average 00Ma95=1.5(0.2) 96Pa01=0.9(+0.4-0.3) **						
* ¹⁵⁸ W ^m	T : average 00Ma95=140(20) 96Pa01=160(50) **						
¹⁵⁹ Pr	-41450# 700#		100# ms				β^- ?
¹⁵⁹ Nd	-50220# 500#		500# ms				β^- ?
¹⁵⁹ Pm	-56850# 200#		1.47 s	0.15	5/2 ⁻ #	03	β^- =100
¹⁵⁹ Sm	-62210 100		11.37 s	0.15	5/2 ⁻	03	β^- =100
¹⁵⁹ Eu	-66053 7		18.1 m	0.1	5/2 ⁺	03	β^- =100
¹⁵⁹ Gd	-68568.5 2.5		18.479 h	0.004	3/2 ⁻	03	β^- =100
¹⁵⁹ Tb	-69539.0 2.6		STABLE		3/2 ⁺	03	IS=100.
¹⁵⁹ Dy	-69173.5 2.7		144.4 d	0.2	3/2 ⁻	03	ε =100
¹⁵⁹ Dy ^m	-68820.7 2.7	352.77 0.14	122 μs	3	11/2 ⁻	03	IT=100
¹⁵⁹ Ho	-67336 4		33.05 m	0.11	7/2 ⁻	03	β^+ =100
¹⁵⁹ Ho ^m	-67130 4	205.91 0.05	8.30 s	0.08	1/2 ⁺	03	IT=100
¹⁵⁹ Er	-64567 4		36 m	1	3/2 ⁻	03	β^+ =100
¹⁵⁹ Er ^m	-64384 4	182.602 0.024	337 ns	14	9/2 ⁺	03	IT=100
¹⁵⁹ Er ⁿ	-64138 4	429.05 0.03	590 ns	60	11/2 ⁻	03	IT=100
¹⁵⁹ Tm	-60570 28		9.13 m	0.16	5/2 ⁺	03	β^+ =100
¹⁵⁹ Yb	-55843 18		1.72 m	0.10	5/2 ⁽⁻⁾	03 93Al03 T	β^+ =100 *
¹⁵⁹ Lu	-49710 40		12.1 s	1.0	1/2 ⁺ #	03	β^+ ≈100; α =0.1#
¹⁵⁹ Lu ^m	-49610# 90#	100# 80#	10# s		11/2 ⁻ #		β^+ ?; IT ?; α ?
¹⁵⁹ Hf	-42854 17		5.20 s	0.10	7/2 ⁻ #	03 96Pa01 T	β^+ =65 7; α =35 7 *
¹⁵⁹ Ta	-34448 21		1.04 s	0.09	(1/2 ⁺)	97Da07 TJ	β^+ ?; α =34 5 *
¹⁵⁹ Ta ^m	-34385 20	64 5 AD	514 ms	9	(11/2 ⁻)	03 96Pa01 T	α =55 1; β^+ ? *
¹⁵⁹ W	-25230# 400#		8.2 ms	0.7	7/2 ⁻ #	03 96Pa01 TD	α =82 16; β^+ ? *
* ¹⁵⁹ Yb	T : supersedes 80Al14=1.40(0.20) from same group **						
* ¹⁵⁹ Hf	J : 7/2 ⁻ is not measured in 00D118, p.7: "a 7/2 ⁻ assignment is assumed" **						
* ¹⁵⁹ Ta	T : average 97Da07=0.83(0.18) 96Pa01=1.10(0.10) **						
* ¹⁵⁹ Ta ^m	T : average 97Da07=500(11) 96Pa01=544(16); other 02Ro17=620(50) **						
* ¹⁵⁹ W	D : derived from original α =92(23)% **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)				
^{160}Nd	-47420#	600#	300# ms	0^+		85Si25 I	$\beta^- ?$	*			
^{160}Pm	-53100#	300#	2# s				$\beta^- ?$				
^{160}Sm	-60420#	200#	9.6 s	0.3	0^+	97	$\beta^- = 100$				
^{160}Eu	-63370#	200#	38 s	4	$1^{(-)}$	97	$\beta^- = 100$				
^{160}Gd	-67948.6	2.6	STABLE	(>31 Ey)	0^+	97	IS=21.86 19; $2\beta^- ?$				
^{160}Tb	-67842.9	2.6	72.3 d	0.2	3^-	97	$\beta^- = 100$				
^{160}Dy	-69678.1	2.5	STABLE		0^+	97	IS=2.34 8				
^{160}Ho	-66388	15	25.6 m	0.3	5^+	97	$\beta^+ = 100$				
$^{160}\text{Ho}^m$	-66328	15	59.98	0.03	5.02	h	0.05	IT=65 3; $\beta^+ = 35$ 3			
$^{160}\text{Ho}^n$	-66191	22	197	16	3	s	(9 ⁺)	97	IT=100	*	
^{160}Er	-66058	24	28.58 h	0.09	0^+	97	$\epsilon = 100$				
^{160}Tm	-60300	30	9.4 m	0.3	1^-	97	$\beta^+ = 100$				
$^{160}\text{Tm}^m$	-60230	40	70	20	74.5	s	1.5	5 ^(#)	97	IT=85 5; $\beta^+ = 15$ 5	
^{160}Yb	-58170	17	4.8 m	0.2	0^+	97	$\beta^+ = 100$				
^{160}Lu	-50270	60	*	*	36.1	s	0.3	$2^- \#$	97	$\beta^+ = 100$; $\alpha < 1e-4$	
$^{160}\text{Lu}^m$	-50270#	120#	0#	100#	*	40	s	1	97	$\beta^+ \approx 100$; $\alpha ?$	
^{160}Hf	-45937	12	13.6 s	0.2	0^+	97	$\beta^+ = 99.3$ 2; $\alpha = 0.7$ 2				
^{160}Ta	-35880	90	1.70 s	0.20	(2#) ⁻	96Pa01	TJD	$\beta^+ ?$; $\alpha = ?$	*		
$^{160}\text{Ta}^m$	-35560#	110#	310#	90#	1.55	s	0.04	(9) ⁺	97	$\beta^+ = 66\%$; $\alpha = ?$	*
^{160}W	-29360	210	90 ms	5	0^+	97	96Pa01	TD	$\alpha = 87$ 8; $\beta^+ ?$	*	
^{160}Re	-16660#	400#	860 μs	120	(2 ⁻)	02	92Pa05	J	p=91 5; $\alpha = 9$ 5	*	
^{160}Nd	I : seen in the thermal fission of ^{252}Cf								**		
$^{160}\text{Ho}^n$	E : less than 55 keV above 169.55 level, from ENSDF								**		
^{160}Ta	J : from α correlation with ^{156}Lu line								**		
$^{160}\text{Ta}^m$	J : from α correlation with $^{156}\text{Lu}^m$ line								**		
^{160}W	T : average 96Pa01=91(5) 81Ho10=81(15)								**		
^{160}Re	J : protons from $d_{3/2}$ orbital								**		
^{161}Nd	-42960#	700#	200# ms		$1/2^- \#$			$\beta^- ?$			
^{161}Pm	-50430#	500#	700# ms		$5/2^- \#$			$\beta^- ?$			
^{161}Sm	-56980#	300#	4.8 s	0.8	$7/2^+ \#$	00		$\beta^- = 100$			
^{161}Eu	-61780#	300#	26 s	3	$5/2^+ \#$	00		$\beta^- = 100$			
^{161}Gd	-65512.7	2.7	3.646 m	0.003	$5/2^-$	00	94It.A	T	$\beta^- = 100$		
^{161}Tb	-67468.2	2.6	6.906 d	0.019	$3/2^+$	00		$\beta^- = 100$			
^{161}Dy	-68061.1	2.5	STABLE		$5/2^+$	00		IS=18.91 24			
^{161}Ho	-67203	3	2.48 h	0.05	$7/2^-$	00		$\epsilon = 100$			
$^{161}\text{Ho}^m$	-66992	3	211.16	0.03	6.76	s	0.07	$1/2^+$	00	IT=100	
^{161}Er	-65209	9	3.21 h	0.03	$3/2^-$	00		$\beta^+ = 100$			
$^{161}\text{Er}^m$	-64813	9	396.44	0.04	7.5	μs	0.7	$11/2^-$	00	IT=100	
^{161}Tm	-61899	28	30.2 m	0.8	$7/2^+$	00		$\beta^+ = 100$			
$^{161}\text{Tm}^m$	-61892	28	7.4	0.2	5#	m		$1/2^+$	00	$\beta^+ ?$; IT ?	
^{161}Yb	-57844	16	4.2 m	0.2	$3/2^-$	00		$\beta^+ = 100$			
^{161}Lu	-52562	28	77 s	2	$1/2^+$	00		$\beta^+ = 100$			
$^{161}\text{Lu}^m$	-52400	30	166	18	7.3	ms	0.4	(9/2 ⁻)	00	ABBW E	
^{161}Hf	-46319	23	18.2 s	0.5	$3/2^- \#$	00		$\beta^+ \approx 100$; $\alpha < 0.13$			
^{161}Ta	-38730#	60#	*	&	3#	s		$1/2^+ \#$			
$^{161}\text{Ta}^m$	-38684	23	50#	50#	2.89	s	0.12	$11/2^- \#$	00	$\beta^+ = 95\%$; $\alpha = ?$	
^{161}W	-30410#	200#	409 ms	16	$7/2^- \#$	00	96Pa01	T	$\alpha = 73$ 3; $\beta^+ = 27$ 3	*	
^{161}Re	-20880	210	370 μs	40	$1/2^+$	02	97Ir01	D	p=97 2; $\alpha ?$	*	
$^{161}\text{Re}^m$	-20750	210	123.8	1.3	15.6	ms	0.9	$11/2^-$	02	$\alpha = ?$; p=4.8 6	
$^{161}\text{Lu}^m$	E : less than K binding energy (61 keV) above 135.6 level, from ENSDF								**		
^{161}W	T : average 96Pa01=409(18) 79Ho10=410(40)								**		
^{161}Re	D : derived from original p=100(7)%								**		

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)		
¹⁶² Pm	-46310#	700#			500# ms					β^- ?	
¹⁶² Sm	-54750#	500#			2.4 s	0.5	0 ⁺	00As.A	TD	β^- =100	
¹⁶² Eu	-58650#	300#			10.6 s	1.0		99		β^- =100	
¹⁶² Gd	-64287	5			8.4 m	0.2	0 ⁺	99		β^- =100	
¹⁶² Tb	-65680	40			7.60 m	0.15	1 ⁻	99		β^- =100	
¹⁶² Dy	-68186.8	2.5			STABLE		0 ⁺	99		IS=25.51 26	
¹⁶² Ho	-66047	4			15.0 m	1.0	1 ⁺	99		β^+ =100	
¹⁶² Ho ^m	-65941	8	106	7	67.0 m	0.7	6 ⁻	99		IT=62; β^+ =38	
¹⁶² Er	-66343	3			STABLE	(>140 Ty)	0 ⁺	99	56Po16	T	IS=0.14 1; α ?; 2 β^+ ?
¹⁶² Tm	-61484	26			21.70 m	0.19	1 ⁻	99		β^+ =100	
¹⁶² Tm ^m	-61350	50	130	40	24.3 s	1.7	5 ⁺	99	ABBW	E	IT ?; β^+ =18 4
¹⁶² Yb	-59832	16			18.87 m	0.19	0 ⁺	99		β^+ =100	
¹⁶² Lu	-52840	80			* 1.37 m	0.02	1 ⁽⁻⁾	99	98Ge13	J	β^+ =100
¹⁶² Lu ^m	-52720#	220#	120#	200#	* 1.5 m		4 ⁻ #	99		β^+ ≈100; IT ?	
¹⁶² Lu ^m	-52540#	220#	300#	200#	* 1.9 m			99		β^+ ≈100; IT ?	
¹⁶² Hf	-49173	10			39.4 s	0.9	0 ⁺	99		β^+ ≈100; α =0.008 1	
¹⁶² Ta	-39780	50			3.57 s	0.12	3 ⁺ #	99		β^+ ≈100; α =0.074 10	
¹⁶² W	-34002	18			1.36 s	0.07	0 ⁺	99		β^+ ?; α =45.2 16	
¹⁶² Re	-22350#	200#			107 ms	13	(2 ⁻)	99		α =94 6; β^+ ?	
¹⁶² Re ^m	-22180#	200#	173	10	AD	77 ms	9	(9 ⁺)	99	α =91 5; β^+ ?	
¹⁶² Os	-14500#	500#			1.87 ms	0.18	0 ⁺	99	00Ma95	T	α =100
* ¹⁶² Ho ^m	E : about 10 keV above level at 96.1(0.1), from ENSDF; error from NUBASE										
* ¹⁶² Er	T : lower limit is for α decay										
* ¹⁶² Tm ^m	E : above 66.90 level and less than 192 keV, from ENSDF										
* ¹⁶² Os	T : average 00Ma95=1.9(0.2) 96Bi07=1.5(+0.7-0.5) 89Ho12=1.9(0.7)										
¹⁶³ Pm	-43150#	800#			200# ms		5/2 ⁻ #			β^- ?	
¹⁶³ Sm	-50900#	700#			1# s		1/2 ⁻ #			β^- ?	
¹⁶³ Eu	-56630#	500#			6# s		5/2 ⁺ #			β^- ?	
¹⁶³ Gd	-61490#	300#			68 s	3	7/2 ⁺ #	00		β^- =100	
¹⁶³ Tb	-64601	5			19.5 m	0.3	3/2 ⁺	00		β^- =100	
¹⁶³ Dy	-66386.5	2.5			STABLE		5/2 ⁻	00		IS=24.90 16	
¹⁶³ Ho	-66383.9	2.5			4.570 ky	0.025	7/2 ⁻	00		ϵ =100	
¹⁶³ Ho ^m	-66086.0	2.5	297.88	0.07	1.09 s	0.03	1/2 ⁺	00		IT=100	
¹⁶³ Er	-65174	5			75.0 m	0.4	5/2 ⁻	00		β^+ =100	
¹⁶³ Er ^m	-64729	5	445.5	0.6	580 ns	100	(11/2 ⁻)	00		IT=100	
¹⁶³ Tm	-62735	6			1.810 h	0.005	1/2 ⁺	00		β^+ =100	
¹⁶³ Yb	-59304	16			11.05 m	0.25	3/2 ⁻	00		β^+ =100	
¹⁶³ Lu	-54791	28			3.97 m	0.13	1/2 ⁽⁺⁾	01		β^+ =100	
¹⁶³ Hf	-49286	28			40.0 s	0.6	3/2 ⁻ #	00		β^+ =100; α <0.0001	
¹⁶³ Ta	-42540	40			10.6 s	1.8	1/2 ⁺ #	00		β^+ ≈100; α ≈0.2	
¹⁶³ W	-34910	50			2.8 s	0.2	3/2 ⁻ #	00		β^+ ?; α =13 2	
¹⁶³ Re	-26007	20			390 ms	70	(1/2 ⁺)	00		β^+ ?; α =32 3	
¹⁶³ Re ^m	-25892	20	115	4	AD	214 ms	5	(11/2 ⁻)	00	α =66 4; β^+ ?	
¹⁶³ Os	-16120#	400#			5.5 ms	0.6	7/2 ⁻ #	00		α ≈100; β^+ ?; β^+ p ?	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁶⁴ Sm	-48180#	800#	500# ms	0 ⁺			β^- ?	
¹⁶⁴ Eu	-53100#	600#	2# s				β^- ?	
¹⁶⁴ Gd	-59750#	400#	45 s	3	0 ⁺	01	β^- =100	
¹⁶⁴ Tb	-62080	100	3.0 m	0.1	(5 ⁺)	01	β^- =100	
¹⁶⁴ Dy	-65973.3	2.5	STABLE		0 ⁺	01	IS=28.18 37	
¹⁶⁴ Ho	-64987.1	2.8	29 m	1	1 ⁺	01	ϵ =60 5; β^- =40 5	
¹⁶⁴ Ho ^m	-64847.3	2.8	139.77 0.08 38.0 m	1.0	6 ⁻	01	IT=100	
¹⁶⁴ Er	-65950	3	STABLE		0 ⁺	01	IS=1.61 3; α ?; 2 β^+ ?	
¹⁶⁴ Tm	-61888	28	* 2.0 m	0.1	1 ⁺	01	ϵ =61 1; e^+ =39 1	
¹⁶⁴ Tm ^m	-61878	29	* 5.1 m	0.1	6 ⁻	01	IT \approx 80; β^+ \approx 20	
¹⁶⁴ Yb	-61023	16	75.8 m	1.7	0 ⁺	01	ϵ =100	
¹⁶⁴ Lu	-54642	28	3.14 m	0.03	1 ⁽⁻⁾	01	β^+ =100	
¹⁶⁴ Hf	-51822	20	111 s	8	0 ⁺	01	β^+ =100	
¹⁶⁴ Ta	-43283	28	14.2 s	0.3	(3 ⁺)	01	β^+ =100	
¹⁶⁴ W	-38234	12	6.3 s	0.2	0 ⁺	01	β^+ =96.2 12; α =3.8 12	
¹⁶⁴ Re	-27640#	160#	* &		high	95Pa.A	J α ?	
¹⁶⁴ Re ^m	-27520	100	* & 530 ms	230	(2#) ⁻	01	96Pa01 JD α =?; β^+ =42#	
¹⁶⁴ Os	-20460	210	21 ms	1	0 ⁺	01	α =?; β^+ =2#	
¹⁶⁴ Ir	-7270#	410#	& 1# ms		2 ⁻ #		p ?; α ?; β^+ ?	
¹⁶⁴ Ir ^m	-7000#	400#	& 94 μ s	27	9 ⁺ #	02	02Ma61 T p=?; α ?; β^+ ?	
* ¹⁶⁴ Tm ^m	E : less than 20 keV, from ENSDF							**
* ¹⁶⁴ Lu	J : negative parity proposed by 98Ge13; odd-odd ¹⁶⁰ Tm ¹⁶² Tm ¹⁶² Lu have 1 ⁻ ground-state							**
* ¹⁶⁴ Ta	D : was erroneously considered as alpha emitter, instead of ¹⁶³ Ta by 83Sc18							**
* ¹⁶⁴ Re ^m	J : from α correlation with ¹⁶⁰ Ta line							**
* ¹⁶⁴ Ir ^m	T : average 02Ma61=58(+46-18) 01Ke05=110(+60-30)							**
¹⁶⁵ Sm	-43800#	900#	200# ms		5/2 ⁻ #		β^- ?	
¹⁶⁵ Eu	-50560#	700#	1# s		5/2 ⁺ #		β^- ?	
¹⁶⁵ Gd	-56470#	500#	10.3 s	1.6	1/2 ⁻ #	99	β^- =100	
¹⁶⁵ Tb	-60660#	200#	2.11 m	0.10	3/2 ⁺ #	92	β^- =100	
¹⁶⁵ Dy	-63617.9	2.5	2.334 h	0.001	7/2 ⁺	92	β^- =100	
¹⁶⁵ Dy ^m	-63509.7	2.5	108.160 0.003 1.257 m	0.006	1/2 ⁻	92	IT=97.76 11; β^- =2.24 11	
¹⁶⁵ Ho	-64904.6	2.5	STABLE		7/2 ⁻	92	IS=100.	
¹⁶⁵ Er	-64528	3	10.36 h	0.04	5/2 ⁻	92	ϵ =100	
¹⁶⁵ Tm	-62936	3	30.06 h	0.03	1/2 ⁺	92	β^+ =100	
¹⁶⁵ Yb	-60287	28	9.9 m	0.3	5/2 ⁻	92	β^+ =100	
¹⁶⁵ Lu	-56442	27	* 10.74 m	0.10	1/2 ⁺	99	β^+ =100	
¹⁶⁵ Hf	-51636	28	76 s	4	(5/2 ⁻)	92	β^+ =100	
¹⁶⁵ Ta	-45855	17	31.0 s	1.5	5/2 ⁻ #	92	β^+ =100	
¹⁶⁵ Ta ^p	-45800	30	60 30 AD		9/2 ⁻ #			
¹⁶⁵ W	-38862	25	5.1 s	0.5	3/2 ⁻ #	99	β^+ \approx 100; α <0.2	
¹⁶⁵ Re	-30657	28	* & 1# s		1/2 ⁺ #	99	β^+ ?; α ?	
¹⁶⁵ Re ^m	-30610	23	AD * & 2.1 s	0.3	11/2 ⁻ #	99	β^+ =87 3; α =13 3	
¹⁶⁵ Os	-21650#	200#	71 ms	3	(7/2 ⁻)	99	α >60; β^+ <40	
¹⁶⁵ Ir	-11630#	220#	< 1# μ s		1/2 ⁺ #	02	p ?; α ?	
¹⁶⁵ Ir ^m	-11440	210	180# 50# 300 μ s	60	11/2 ⁻	02	p=87 4; α =13 4	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁶⁶ Eu	-46600# 800#		400# ms				β^- ?	
¹⁶⁶ Gd	-54400# 600#		4.8 s	1.0	0+	00As.A TD	β^- =100	
¹⁶⁶ Tb	-57760 100		25.6 s	2.2		97 00As.A T	β^- =100 *	
¹⁶⁶ Dy	-62590.1 2.6		81.6 h	0.1	0+	92	β^- =100	
¹⁶⁶ Ho	-63076.9 2.5		26.83 h	0.02	0-	92	β^- =100	
¹⁶⁶ Ho ^m	-63070.9 2.5	5.985 0.018	1.20 ky	0.18	(7)-	92	β^- =100	
¹⁶⁶ Er	-64931.6 2.5		STABLE		0+	92	IS=33.61 35	
¹⁶⁶ Tm	-61894 12		7.70 h	0.03	2+	92	β^+ =100	
¹⁶⁶ Tm ^m	-61772 14	122 8	340 ms	25	6-	96Dr07 TJE	IT=100 *	
¹⁶⁶ Yb	-61588 8		56.7 h	0.1	0+	92	ϵ =100	
¹⁶⁶ Lu	-56021 30		2.65 m	0.10	6(-)	92 98Ge13 J	β^+ =100	
¹⁶⁶ Lu ^m	-55990 30	34.37 0.05	1.41 m	0.10	3(-)	92 98Ge13 J	β^+ =58 5; IT=42 5	
¹⁶⁶ Lu ⁿ	-55980 30	42.9 0.5	2.12 m	0.10	0(-)	92 98Ge13 J	β^+ >80; IT<20	
¹⁶⁶ Hf	-53859 28		6.77 m	0.30	0+	92	β^+ =100	
¹⁶⁶ Ta	-46098 28		34.4 s	0.5	(2)+	92	β^+ =100	
¹⁶⁶ W	-41892 10		19.2 s	0.6	0+	00	β^+ ≈100; α =0.035 12	
¹⁶⁶ Re	-31850# 90#		& 2# s		2-#		β^+ ?; α ?	
¹⁶⁶ Re ^m	-31700 70	150# 50#	& 2.5 s	0.2	9+#	92 92Me10 T	β^+ ?; α =5 2 *	
¹⁶⁶ Re ^p	-31700# 100#	150# 50#			low			
¹⁶⁶ Os	-25438 18		216 ms	9	0+	92 96Pa01 T	α =72 13; β^+ =28 13 *	
¹⁶⁶ Ir	-13210# 200#		10.5 ms	2.2	(2)-	02	α =93 3; p=7 3	
¹⁶⁶ Ir ^m	-13030# 200#	172 6 p	15.1 ms	0.9	(9+)	02	α =98.2 6; p=1.8 6	
¹⁶⁶ Pt	-4790# 500#		300 μs	100	0+	97 96Bi07 TD	α =100	
* ¹⁶⁶ Tb	T : supersedes 94Ts.A=21(6) same group							**
* ¹⁶⁶ Tm ^m	E : less than 25 keV above 109.34 level							**
* ¹⁶⁶ Re ^m	T : average 92Me10=2.3(0.2) 84Sc06=2.8(0.3)							**
* ¹⁶⁶ Re ^m	D : α intensity is derived from 2% < α < 8% as discussed in ENSDF							**
* ¹⁶⁶ Os	T : average 96Pa01=220(7) 91Se01=194(17)							**
¹⁶⁷ Eu	-43590# 800#		200# ms		5/2+#		β^- ?	
¹⁶⁷ Gd	-50700# 600#		3# s		5/2-#		β^- ?	
¹⁶⁷ Tb	-55840# 400#		19 s	3	3/2+#	00 99As03 T	β^- =100	
¹⁶⁷ Dy	-59940 60		6.20 m	0.08	(1/2)-	00	β^- =100	
¹⁶⁷ Ho	-62287 6		3.1 h	0.1	7/2-	00	β^- =100	
¹⁶⁷ Ho ^m	-62028 6	259.34 0.11	6.0 μs	1.0	3/2+	00	IT=100	
¹⁶⁷ Er	-63296.7 2.5		STABLE		7/2+	00	IS=22.93 17	
¹⁶⁷ Er ^m	-63088.9 2.5	207.801 0.005	2.269 s	0.006	1/2-	00	IT=100	
¹⁶⁷ Tm	-62548.3 2.7		9.25 d	0.02	1/2+	00	ϵ =100	
¹⁶⁷ Tm ^m	-62368.8 2.7	179.480 0.019	1.16 μs	0.06	(7/2)+	00	IT=100	
¹⁶⁷ Tm ⁿ	-62255.5 2.7	292.820 0.020	0.9 μs	0.1	7/2-	00	IT=100	
¹⁶⁷ Yb	-60594 5		17.5 m	0.2	5/2-	00	β^+ =100	
¹⁶⁷ Lu	-57500 30		51.5 m	1.0	7/2+	00	β^+ =100	
¹⁶⁷ Lu ^m	-57500# 40#	0# 30#	> 1 m		1/2(-#)	00	IT ?; β^+ ?	
¹⁶⁷ Hf	-53468 28		2.05 m	0.05	(5/2)-	00	β^+ =100	
¹⁶⁷ Ta	-48351 28		1.33 m	0.07	(3/2+)	00	β^+ =100	
¹⁶⁷ W	-42089 19		19.9 s	0.5	3/2-#	00	β^+ =99.96 1; α =0.04 1 *	
¹⁶⁷ Re	-34840# 50#		& 3.4 s	0.4	9/2-#	00	α ≈100; β^+ ?	
¹⁶⁷ Re ^m	-34710 40	130# 40#	& 5.9 s	0.3	1/2+#	00	β^+ ≈99; α ≈1	
¹⁶⁷ Os	-26500 70		810 ms	60	3/2-#	00	α =57 8; β^+ =43 8	
¹⁶⁷ Ir	-17079 19		35.2 ms	2.0	1/2+	02	α =48 6; p=32 4; β^+ ?	
¹⁶⁷ Ir ^m	-16903 19	175.3 2.2 p	30.0 ms	0.6	11/2-	02	α =80 10; β^+ ?; ... *	
¹⁶⁷ Pt	-6540# 410#		700 μs	200	7/2-#	00	α =100	
* ¹⁶⁷ W	J : lowest observed state by 92Th06 is 13/2+							**
* ¹⁶⁷ Ir ^m	D : ... ; p=0.4 1							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁶⁸ Gd	-48100# 700#		300# ms	0 ⁺		85Si25 I	β^- ? *
¹⁶⁸ Tb	-52500# 500#		8.2 s 1.3	4 ⁻ #	99		β^- =100
¹⁶⁸ Dy	-58560 140		8.7 m 0.3	0 ⁺	99		β^- =100
¹⁶⁸ Ho	-60070 30		2.99 m 0.07	3 ⁺	94		β^- =100
¹⁶⁸ Ho ^m	-60010 30	59 1	132 s 4	(6 ⁺)	94	90Ch37 E	IT \approx 100; β^- <0.5
¹⁶⁸ Er	-62996.7 2.5		STABLE	0 ⁺	94		IS=26.78 26
¹⁶⁸ Tm	-61317.7 2.9		93.1 d 0.2	3 ⁺	94		β^+ \approx 100; β^- =0.010 7
¹⁶⁸ Yb	-61575 4		STABLE (>130 Ty)	0 ⁺	94	56Po16 T	IS=0.13 1; α ?; $2\beta^+$? *
¹⁶⁸ Lu	-57060 50		5.5 m 0.1	6 ⁽⁻⁾	94	98Ge13 J	β^+ =100
¹⁶⁸ Lu ^m	-56880 100	180 110	6.7 m 0.4	3 ⁺	94		β^+ >95; IT<5
¹⁶⁸ Hf	-55361 28		25.95 m 0.20	0 ⁺	94		$\epsilon\approx$ 98; $e^+\approx$ 2
¹⁶⁸ Ta	-48394 28		2.0 m 0.1	(2 ⁻ , 3 ⁺)	94		β^+ =100
¹⁶⁸ W	-44890 16		51 s 2	0 ⁺	94		β^+ \approx 100; α =0.0032 10
¹⁶⁸ Re	-35790 30		4.4 s 0.1	(5 ⁺ , 6 ⁺ , 7 ⁺)	94		β^+ \approx 100; $\alpha\approx$ 0.005
¹⁶⁸ Re ^m		non existent	6.6 s 1.5			92Me10 I	
¹⁶⁸ Os	-29991 12		2.06 s 0.06	0 ⁺	94	96Pa01 T	β^+ =51 3; α =49 3 *
¹⁶⁸ Ir	-18740# 150#		161 ms 21	high	94	96Pa01 TJD	α =82 14
¹⁶⁸ Ir ^m	-18690 110	50# 100#	125 ms 40	low	94	96Pa01 TJ	α =?; β^+ ?
¹⁶⁸ Pt	-11040 210		2.00 ms 0.18	0 ⁺	94	98Ki20 T	$\alpha\approx$ 100; β^+ =0.7# *
* ¹⁶⁸ Gd	I : seen in the thermal fission of ²⁵² Cf **						
* ¹⁶⁸ Yb	T : lower limit is for α decay **						
* ¹⁶⁸ Os	T : average 96Pa01=2.1(0.1) 84Sc06=2.0(0.2) 82En03=2.2(0.1) 78Ca11=1.9(0.1) **						
* ¹⁶⁸ Os	T : 84Sc06 supersedes 78Sc26=2.4(0.2) from same group **						
* ¹⁶⁸ Pt	T : average 98Ki20=2.0(0.2) 96Bi07=2.0(0.4) **						
¹⁶⁹ Gd	-43900# 800#		1# s	7/2 ⁻ #			β^- ?
¹⁶⁹ Tb	-50100# 600#		2# s	3/2 ⁺ #			β^- ?
¹⁶⁹ Dy	-55600 300		39 s 8	(5/2 ⁻)	91		β^- =100
¹⁶⁹ Ho	-58803 20		4.7 m 0.1	7/2 ⁻	91		β^- =100
¹⁶⁹ Er	-60928.7 2.5		9.40 d 0.02	1/2 ⁻	91		β^- =100
¹⁶⁹ Tm	-61280.0 2.5		STABLE	1/2 ⁺	91		IS=100.
¹⁶⁹ Yb	-60370 4		32.026 d 0.005	7/2 ⁺	91		ϵ =100
¹⁶⁹ Yb ^m	-60346 4	24.199 0.003	46 s 2	1/2 ⁻	91		IT=100
¹⁶⁹ Lu	-58077 5		34.06 h 0.05	7/2 ⁺	91		β^+ =100
¹⁶⁹ Lu ^m	-58048 5	29.0 0.5	160 s 10	1/2 ⁻	91		IT=100
¹⁶⁹ Hf	-54717 28		3.24 m 0.04	(5/2 ⁻)	91		β^+ =100
¹⁶⁹ Ta	-50290 28		4.9 m 0.4	(5/2 ⁺)	91	98Zh03 J	β^+ =100
¹⁶⁹ W	-44918 15		76 s 6	(5/2 ⁻)	91		β^+ =100
¹⁶⁹ Re	-38386 28		8.1 s 0.5	9/2 ⁻ #	91	92Me10 TD	β^+ =?; α =0.005 3 *
¹⁶⁹ Re ^m	-38241 17	145 29	15.1 s 1.6	1/2 ⁺ #	91	92Me10 TD	β^+ ?; $\alpha\approx$ 0.2 *
¹⁶⁹ Os	-30721 25		3.46 s 0.11	3/2 ⁻ #	91	96Pa01 T	β^+ =89 1; α =11 1 *
¹⁶⁹ Ir	-22081 26		& 780 ms 360	1/2 ⁺ #		99Po09 TD	α =50 18; β^+ ? *
¹⁶⁹ Ir ^m	-21927 22	154 24	& 308 ms 22	11/2 ⁻ #	91	96Pa01 TD	α =81 7; β^+ =19 7 *
¹⁶⁹ Pt	-12380# 200#		3.7 ms 1.5	3/2 ⁻ #	91	96Pa01 T	α =?; β^+ =1# *
¹⁶⁹ Au	-1790# 300#		150# μ s	1/2 ⁺ #			α ?; β^+ ? *
* ¹⁶⁹ Re	D : α =0.005(3)% derived from original α =0.001% - 0.01% **						
* ¹⁶⁹ Re ^m	T : average 92Me10=16.3(0.8) 84Sc06=12.9(1.1) **						
* ¹⁶⁹ Os	T : average 96Pa01=3.6(0.2) 95Hi02=3.2(0.3) 84Sc06=3.5(0.2) 82En03=3.4(0.2) **						
* ¹⁶⁹ Ir ^m	T : also 99Po09=323(+90-66) D : average 99Po09=84(8)% 96Pa01=72(13)% **						
* ¹⁶⁹ Pt	T : average 96Pa01=5(3) 81Ho10=2.5(+2.5-1.0) **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁷⁰ Tb	-46340# 700#		3# s				$\beta^- ?$
¹⁷⁰ Dy	-53660# 200#		30# s	0 ⁺	02		$\beta^- ?$
¹⁷⁰ Ho	-56240 50		2.76 m	0.05	6 ⁺ #	02	$\beta^- = 100$
¹⁷⁰ Ho ^m	-56140 60	100 80	43 s	2	(1 ⁺)	02	$\beta^- = 100$
¹⁷⁰ Er	-60114.6 2.8		STABLE	(>320 Py)	0 ⁺	02	96De60 T IS=14.93 27; ... *
¹⁷⁰ Tm	-59800.6 2.5		128.6 d	0.3	1 ⁻	02	$\beta^- \approx 100$; $\epsilon = 0.131$ 10 *
¹⁷⁰ Tm ^m	-59617.4 2.5	183.197 0.004	4.12 μ s	0.13	(3) ⁺	02	IT=100
¹⁷⁰ Yb	-60769.0 2.4		STABLE		0 ⁺	02	IS=3.04 15
¹⁷⁰ Yb ^m	-59510.5 2.4	1258.46 0.14	370 ns	15	4 ⁻	02	IT=100
¹⁷⁰ Lu	-57310 17		2.012 d	0.020	0 ⁺	02	$\beta^+ = 100$
¹⁷⁰ Lu ^m	-57217 17	92.91 0.09	670 ms	100	(4) ⁻	02	IT=100
¹⁷⁰ Hf	-56254 28		16.01 h	0.13	0 ⁺	02	$\epsilon = 100$
¹⁷⁰ Ta	-50138 28		6.76 m	0.06	(3) ⁽⁺⁾	02	$\beta^+ = 100$
¹⁷⁰ W	-47293 15		2.42 m	0.04	0 ⁺	02	$\beta^+ \approx 100$; $\alpha < 1\%$
¹⁷⁰ Re	-38918 26		9.2 s	0.2	(5) ⁺	02	$\beta^+ \approx 100$; $\alpha < 0.01\%$
¹⁷⁰ Os	-33928 11		7.46 s	0.23	0 ⁺	02	$\beta^+ = ?$; $\alpha = 8.6$ 18
¹⁷⁰ Ir	-23320# 100#		910 ms	150	low#	02	$\beta^+ ?$; $\alpha = 5.2$ 17
¹⁷⁰ Ir ^m	-23050 70	270# 70#	440 ms	60	high#	02	$\alpha = 36$ 10; $\beta^+ ?$; IT ?
¹⁷⁰ Pt	-16306 19		13.8 ms	0.5	0 ⁺	02	$\alpha = ?$; $\beta^+ = 2\%$
¹⁷⁰ Au	-3610# 200#		310 μ s	50	(2) ⁻	02	p=85 10; $\alpha = 15$ 10
¹⁷⁰ Au ^m	-3340# 200#	274 16	630 μ s	60	(9) ⁺	02	02Ma61 TD p=75 15; $\alpha = ?$; $\beta^+ ?$ *
* ¹⁷⁰ Er	D : ... ; $2\beta^- ?$; $\alpha ?$ **						
* ¹⁷⁰ Au ^m	T : from 02Ke.C=620(+60-50); other 02Ma61=570(+310-150) **						
¹⁷¹ Tb	-43500# 800#		500# ms		3/2 ⁺ #		$\beta^- ?$
¹⁷¹ Dy	-50110# 300#		6# s		7/2 ⁻ #		$\beta^- ?$
¹⁷¹ Ho	-54520 600		53 s	2	7/2 ⁻ #	02	$\beta^- = 100$
¹⁷¹ Er	-57724.9 2.8		7.516 h	0.002	5/2 ⁻	02	$\beta^- = 100$
¹⁷¹ Er ^m	-57526.3 2.8	198.6 0.1	210 ns	10	1/2 ⁻	02	IT=100
¹⁷¹ Tm	-59215.6 2.6		1.92 y	0.01	1/2 ⁺	02	$\beta^- = 100$
¹⁷¹ Tm ^m	-58790.6 2.6	424.9560 0.0015	2.60 μ s	0.02	7/2 ⁻	02	IT=100
¹⁷¹ Yb	-59312.1 2.4		STABLE		1/2 ⁻	02	IS=14.28 57
¹⁷¹ Yb ^m	-59216.8 2.4	95.282 0.002	5.25 ms	0.24	7/2 ⁺	02	IT=100
¹⁷¹ Yb ⁿ	-59189.7 2.4	122.416 0.002	265 ns	20	5/2 ⁻	02	IT=100
¹⁷¹ Lu	-57833.5 2.8		8.24 d	0.03	7/2 ⁺	02	$\beta^+ = 100$
¹⁷¹ Lu ^m	-57762.4 2.8	71.13 0.08	79 s	2	1/2 ⁻	02	IT=100
¹⁷¹ Hf	-55431 29		12.1 h	0.4	7/2 ⁽⁺⁾	02	$\beta^+ = 100$
¹⁷¹ Hf ^m	-55409 29	21.93 0.09	29.5 s	0.9	1/2 ⁽⁻⁾	02	IT \approx 100; $\beta^+ ?$
¹⁷¹ Ta	-51720 28		23.3 m	0.3	(5/2 ⁻)	02	$\beta^+ = 100$
¹⁷¹ W	-47086 28		2.38 m	0.04	(5/2 ⁻)	02	$\beta^+ = 100$
¹⁷¹ Re	-41250 28		15.2 s	0.4	(9/2 ⁻)	02	$\beta^+ = 100$
¹⁷¹ Os	-34293 19		8.3 s	0.2	(5/2 ⁻)	02	$\beta^+ ?$; $\alpha = 1.80$ 21
¹⁷¹ Ir	-26430 40		3.6 s	1.0	1/2 ⁺ #	02	$\alpha \approx 100$; $\beta^+ ?$
¹⁷¹ Ir ^m	-26250# 50#	180# 30#	1.40 s	0.10	(11/2 ⁻)	02	99Ba84 J $\alpha = 58$ 11; $\beta^+ ?$; p ?
¹⁷¹ Pt	-17470 90		44 ms	7	3/2 ⁻ #	02	$\alpha = ?$; $\beta^+ = 2\%$
¹⁷¹ Au	-7565 26		30 μ s	5	(1/2 ⁺)	02	03Ba20 T p \approx 100; $\alpha ?$ *
¹⁷¹ Au ^m	-7315 20	250 16	1.014 ms	0.019	11/2 ⁻	02	03Ba20 TJ $\alpha = 54$ 4; p=46 4
¹⁷¹ Hg	3500# 300#		80 μ s	30	3/2 ⁻ #	02	$\alpha \approx 100$; $\beta^+ = 0.01\%$
* ¹⁷¹ Au	T : average 03Ba20=37(+7-5) 99Po09=17(+9-5); Birge ratio B=2.0 **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁷² Dy	-47730# 400#		3# s	0 ⁺			β^- ?	
¹⁷² Ho	-51400# 400#		25 s	3	95		β^- =100	
¹⁷² Er	-56489 5		49.3 h	0.3	95		β^- =100	
¹⁷² Tm	-57380 6		63.6 h	0.2	95		β^- =100	
¹⁷² Yb	-59260.3 2.4		STABLE		95		IS=21.83 67	
¹⁷² Lu	-56741.3 3.0		6.70 d	0.03	95		β^+ =100	
¹⁷² Lu ^m	-56699 3	41.86 0.04	3.7 m	0.5	95		IT=100	
¹⁷² Lu ⁿ	-56632 3	109.41 0.10	440 μ s	12			(1) ⁺	
¹⁷² Hf	-56404 24		1.87 y	0.03	95		ϵ =100	
¹⁷² Hf ^m	-54398 24	2005.58 0.11	163 ns	3			(8 ⁻)	
¹⁷² Ta	-51330 28		36.8 m	0.3	95		(3 ⁺)	
¹⁷² W	-49097 28		6.6 m	0.9	95		0 ⁺	
¹⁷² Re	-41520 50		15 s	3	95		(5)	
¹⁷² Re ^m	-41520# 110#	0# 100#	55 s	5	95		(2)	
¹⁷² Os	-37238 15		19.2 s	0.9	95	95Hi02 D	β^+ =?; α =1.1 2	
¹⁷² Ir	-27520# 110#		4.4 s	0.3	95		(3 ⁺)	
¹⁷² Ir ^m	-27240 30	280# 100#	2.0 s	0.1	95		(7 ⁺)	
¹⁷² Pt	-21101 13		98.4 ms	2.4	95	02Ro17 T	α =77 21; β^+ ? *	
¹⁷² Au	-9280# 160#		4.7 ms	1.1	95	96Pa01 TJ	α =?; p<2 *	
¹⁷² Hg	-1090 210		420 μ s	240		99Se14 TD	α =100	
* ¹⁷² Pt	T : average 02Ro17=104(7) 96Pa01=96(3) 82En03=90(10) 81De22=120(10) and							**
* ¹⁷² Pt	T : 75Ga25=100(10) D : derived from original α =94(32)%							**
* ¹⁷² Au	T : average 96Pa01=6.3(1.5) 93Se09=4(1)							**
* ¹⁷² Au	J : from α correlation with ¹⁶⁸ Ir line							**
¹⁷³ Dy	-43780# 500#		2# s	9/2 ⁺ #			β^- ?	
¹⁷³ Ho	-49100# 400#		10# s	7/2 ⁻ #			β^- ?	
¹⁷³ Er	-53650# 200#		1.434 m	0.017	95	94It.A T	β^- =100	
¹⁷³ Tm	-56259 5		8.24 h	0.08	95		β^- =100	
¹⁷³ Tm ^m	-55941 5	317.73 0.20	10 μ s				(7/2 ⁻)	
¹⁷³ Yb	-57556.3 2.4		STABLE		95		IS=16.13 27	
¹⁷³ Yb ^m	-57157.4 2.5	398.9 0.5	2.9 μ s	0.1			1/2 ⁻	
¹⁷³ Lu	-56885.8 2.4		1.37 y	0.01	95		ϵ =100	
¹⁷³ Lu ^m	-56762.1 2.4	123.672 0.013	74.2 μ s				5/2 ⁻	
¹⁷³ Hf	-55412 28		23.6 h	0.1	95		β^+ =100	
¹⁷³ Ta	-52397 28		3.14 h	0.13	95		β^+ =100	
¹⁷³ W	-48727 28		7.6 m	0.2	95		β^+ =100	
¹⁷³ Re	-43554 28		2.0 m	0.3	95		β^+ =100	
¹⁷³ Os	-37438 15		22.4 s	0.9	95	95Hi02 TD	β^+ \approx 100; α =0.4 2	
¹⁷³ Ir	-30272 14		9.0 s	0.8	95		(3/2 ⁺ , 5/2 ⁺)	
¹⁷³ Ir ^m	-30019 28	253 27	2.20 s	0.05	95		(11/2 ⁻)	
¹⁷³ Pt	-21940 60		365 ms	7	95	02Ro17 T	α =84 6; β^+ =16 6 *	
¹⁷³ Au	-12820 26		25 ms	1	03		(1/2 ⁺)	
¹⁷³ Au ^m	-12606 22	214 23	14.0 ms	0.9	03		α =86 13; β^+ =6# *	
¹⁷³ Hg	-2570# 210#		1.1 ms	0.4	03		α =89 11; β^+ =4#	
* ¹⁷³ Pt	T : average 02Ro17=370(13) 96Pa01=376(11) 82En03=360(20) and 81De22=325(20)							**
* ¹⁷³ Au	D : from 94(+6-19)%; and for isomer ¹⁷³ Au ^m 92(+8-13)%							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁷⁴ Ho	-45500# 500#		8# s				β^- ?	
¹⁷⁴ Er	-51950# 300#		3.2 m	0.2	0+	99	β^- =100	
¹⁷⁴ Tm	-53870 40		5.4 m	0.1	(4) ⁻	99	β^- =100	
¹⁷⁴ Yb	-56949.6 2.4		STABLE		0+	99	IS=31.83 92	
¹⁷⁴ Lu	-55575.3 2.4		3.31 y	0.05	1 ⁻	99	β^+ =100	
¹⁷⁴ Lu ^m	-55404.5 2.4	170.83 0.05	142 d	2	6 ⁻	99	IT=99.38 2; ϵ =0.62 2	
¹⁷⁴ Hf	-55846.6 2.8		2.0 Py	0.4	0+	99	IS=0.16 1; α =100; $2\beta^+$?	
¹⁷⁴ Hf ^m	-54049 3	1797.5 2.0	2.39 μ s	0.04	(8) ⁻	99	IT=100	
¹⁷⁴ Ta	-51741 28		1.14 h	0.08	3+	99	β^+ =100	
¹⁷⁴ W	-50227 28		33.2 m	2.1	0+	99	β^+ =100	
¹⁷⁴ Re	-43673 28		2.40 m	0.04		99	β^+ =100	
¹⁷⁴ Os	-39996 11		44 s	4	0+	99	β^+ \approx 100; α =0.024 7	
¹⁷⁴ Ir	-30869 28		7.9 s	0.6	(3) ⁺	99	β^+ =99.5 3; α =0.5 3	
¹⁷⁴ Ir ^m	-30676 26	193 11 AD	4.9 s	0.3	(7) ⁺	99	β^+ =97.5 3; α =2.5 3	
¹⁷⁴ Pt	-25319 12		889 ms	17	0+	99	α =76 8; β^+ ?	
¹⁷⁴ Au	-14200# 100#		139 ms	3	low	99	α =90 6; β^+ ?	
¹⁷⁴ Au ^m	-13840 70	360# 70#	171 ms	29	high	99	α =?; β^+ ?	
¹⁷⁴ Hg	-6647 20		2.0 ms	0.4	0+	99	α \approx 100; β^+ =0.4#	
* ¹⁷⁴ Au	T : others 96Pa01=171(29) 83Sc24=120(20)							**
¹⁷⁵ Ho	-42800# 600#		5# s		7/2 ⁻ #		β^- ?	
¹⁷⁵ Er	-48650# 400#		1.2 m	0.3	(9/2 ⁺)	98	β^- =100	
¹⁷⁵ Tm	-52320 50		15.2 m	0.5	1/2 ⁺	98	β^- =100	
¹⁷⁵ Yb	-54700.6 2.4		4.185 d	0.001	7/2 ⁻	93	β^- =100	
¹⁷⁵ Yb ^m	-54185.7 2.4	514.869 0.007	68.2 ms	0.3	1/2 ⁻	93	IT=100	
¹⁷⁵ Lu	-55170.7 2.2		STABLE		7/2 ⁺	93	IS=97.41 2	
¹⁷⁵ Lu ^m	-53780 4	1391 3	930 μ s	80	19/2 ⁺	98	Wh02 ETJ IT=100	
¹⁷⁵ Hf	-54483.8 2.8		70 d	2	5/2 ⁻	93	ϵ =100	
¹⁷⁵ Ta	-52409 28		10.5 h	0.2	7/2 ⁺	93	β^+ =100	
¹⁷⁵ W	-49633 28		35.2 m	0.6	(1/2 ⁻)	93	β^+ =100	
¹⁷⁵ Re	-45288 28		5.89 m	0.05	(5/2 ⁻)	93	β^+ =100	
¹⁷⁵ Os	-40105 14		1.4 m	0.1	(5/2 ⁻)	93	β^+ =100	
¹⁷⁵ Ir	-33429 20		9 s	2	(5/2 ⁻)	93	β^+ =99.15 28; α =0.85 28	
¹⁷⁵ Ir ^p	-33357 17	72 17 AD			am			
¹⁷⁵ Pt	-25690 19		2.52 s	0.08	5/2 ⁻ #	93	α =64 5; β^+ ?	
¹⁷⁵ Au	-17440 40		& 100# ms		1/2 ⁺ #	02Ro17 D	α =?; β^+ ?	
¹⁷⁵ Au ^m	-17240# 50#	200# 30#	& 156 ms	3	11/2 ⁻ #	93	α =82 17; β^+ ?	
¹⁷⁵ Hg	-7990 100		10.8 ms	0.4	5/2 ⁻ #	93	α =?; β^+ =1#	
* ¹⁷⁵ Au	D : from analysis of data in 02Ro17, we assign the 6412 line to ¹⁷⁵ Au							**
* ¹⁷⁵ Au ^m	T : average 02Ro17=158(3) 01Ko44=143(8); others 96Pa01=185(30) 83Sc24=200(22)							**
* ¹⁷⁵ Hg	T : others 97Uu01=13(+6-4) 96Pa01=8(8) outweighed, not used							**
¹⁷⁶ Er	-46500# 400#		20# s		0+		β^- ?	
¹⁷⁶ Tm	-49370 100		1.85 m	0.03	(4) ⁺	98	β^- =100	
¹⁷⁶ Yb	-53494.1 2.6		STABLE	(>160 Py)	0+	98	IS=12.76 41; ...	
¹⁷⁶ Yb ^m	-52444.1 2.6	1050.0 0.3	11.4 s	0.3	(8) ⁻	98	IT=?; β^- <10#	
¹⁷⁶ Lu	-53387.4 2.2		38.5 Gy	0.7	7 ⁻	98	IS=2.59 2; β^- =100	
¹⁷⁶ Lu ^m	-53264.5 2.2	122.855 0.006	3.664 h	0.019	1 ⁻	98	β^- \approx 100; ϵ =0.095 16	
¹⁷⁶ Hf	-54577.5 2.2		STABLE		0+	98	IS=5.26 7	
¹⁷⁶ Ta	-51370 30		8.09 h	0.05	(1) ⁻	98	β^+ =100	
¹⁷⁶ Ta ^m	-51270 30	103.0 1.0	1.1 ms	0.1	(+)	98	IT=100	
¹⁷⁶ Ta ⁿ	-48550 60	2820 50	0.97 ms	0.07	(20) ⁻	98	IT=100	
... A-group is continued on next page ...								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
¹⁷⁶ W	-50642	28	2.5 h	0.1	0 ⁺	98	$\epsilon=100$	
¹⁷⁶ Re	-45063	28	5.3 m	0.3	3 ⁺	98	$\beta^+=100$	
¹⁷⁶ Os	-42098	28	3.6 m	0.5	0 ⁺	98	$\beta^+=100$	
¹⁷⁶ Ir	-33861	20	8.3 s	0.6		98	$\beta^+=96.9$ 6; $\alpha=3.1$ 6	
¹⁷⁶ Pt	-28928	14	6.33 s	0.15	0 ⁺	98	$\beta^+ ?$; $\alpha=38$ 3	
¹⁷⁶ Au	-18540#	110#	1.08 s	0.17	(5 ⁻)	98	ABW J $\alpha=?$; $\beta^+=40$ #	
¹⁷⁶ Au ^m	-18380	30	860 ms	160	(7 ⁺)	02Ro17	T $\alpha=?$; $\beta^+=40$ #	
¹⁷⁶ Hg	-11779	14	20.4 ms	1.5	0 ⁺	98	02Ro17 T $\alpha=90$ 9; $\beta^+ ?$	
¹⁷⁶ Tl	550#	200#	10# ms				$\alpha ?$	
* ¹⁷⁶ Yb	D : ... ; 2 $\beta^- ?$; $\alpha ?$							**
* ¹⁷⁶ Lu	T : arithmetic average 03Gr02=40.8(0.3) 98Ni07=36.9(0.2) 92Da03=37.3(0.5)							**
* ¹⁷⁶ Lu	T : 90Ge05=40.5(0.9) 83Sa44=37.8(0.2) 82Sg01=35.9(0.5) 80No01=40.8(2.4)							**
* ¹⁷⁶ Lu	T : 72Ko50=37.9(0.3) (a weighed average would yield Birge ratio B=4.6)							**
* ¹⁷⁶ Ta ⁿ	E : 2774.8(1.5) + x, and x estimated 50(50) by NUBASE							**
* ¹⁷⁶ Au	J : from α decay to ¹⁷² Ir 168.4 level							**
* ¹⁷⁶ Au ^m	J : from α decay to ¹⁷² Ir ^m							**
* ¹⁷⁶ Hg	T : average 02Ro17=20(2) 99He25=21(3) 99Po09=21(4); others not used							**
* ¹⁷⁶ Hg	T : 96Pa01=18(10) and 83Sc24=34(+18-9)							**
¹⁷⁷ Er	-42800#	500#	3#	s	1/2 ⁻ #		$\beta^- ?$	
¹⁷⁷ Tm	-47470#	300#	90	s	6	(7/2 ⁻)	03 $\beta^-=100$	
¹⁷⁷ Yb	-50989.2	2.6	1.911 h	0.003		(9/2 ⁺)	03 $\beta^-=100$	
¹⁷⁷ Yb ^m	-50657.7	2.6	331.5	0.3	6.41 s	0.02	(1/2 ⁻)	03 IT=100
¹⁷⁷ Lu	-52389.0	2.2			6.647 d	0.004	7/2 ⁺	03 $\beta^-=100$
¹⁷⁷ Lu ^m	-51418.8	2.2	970.1750	0.0024	160.44 d	0.06	23/2 ⁻	03 $\beta^-=78.6$ 8; IT=21.4 8
¹⁷⁷ Lu ⁿ	-48489	10	3900	10	7 m	2	39/2 ⁻	03 03Al.1 ET $\beta^-=?$; IT ?
¹⁷⁷ Lu ^p	-52238.6	2.2	150.3967	0.0010	130 ns	3	9/2 ⁻	03 IT=100
¹⁷⁷ Lu ^q	-51819.3	2.2	569.7068	0.0016	155 μ s	7	1/2 ⁺	03 IT=100
¹⁷⁷ Hf	-52889.6	2.1			STABLE		7/2 ⁻	03 IS=18.60 9
¹⁷⁷ Hf ^m	-51574.1	2.1	1315.4504	0.0008	1.09 s	0.05	23/2 ⁺	03 IT=100
¹⁷⁷ Hf ⁿ	-50149.6	2.1	2740.02	0.15	51.4 m	0.5	37/2 ⁻	03 IT=100
¹⁷⁷ Hf ^p	-51547.2	2.1	1342.38	0.20	55.9 μ s	1.2	(19/2 ⁻)	03 IT=100
¹⁷⁷ Ta	-51724	4			56.56 h	0.06	7/2 ⁺	03 $\beta^+=100$
¹⁷⁷ Ta ^m	-51538	4	186.15	0.06	3.62 μ s	0.10	5/2 ⁻	03 IT=100
¹⁷⁷ Ta ⁿ	-50369	4	1355.01	0.19	5.31 μ s	0.25	21/2 ⁻	03 IT=100
¹⁷⁷ Ta ^p	-51651	4	73.36	0.15	410 ns	7	9/2 ⁻	03 IT=100
¹⁷⁷ Ta ^q	-47068	4	4656.3	0.5	133 μ s	4	49/2 ⁻	03 IT=100
¹⁷⁷ W	-49702	28			132 m	2	1/2 ⁻	03 $\beta^+=100$
¹⁷⁷ Re	-46269	28			14 m	1	5/2 ⁻	03 $\beta^+=100$
¹⁷⁷ Re ^m	-46184	28	84.71	0.10	50 μ s	10	5/2 ⁺	03 IT=100
¹⁷⁷ Os	-41950	16			3.0 m	0.2	1/2 ⁻	03 $\beta^+=100$
¹⁷⁷ Ir	-36047	20			30 s	2	5/2 ⁻	03 $\beta^+\approx 100$; $\alpha=0.06$ 1
¹⁷⁷ Pt	-29370	15			10.6 s	0.4	5/2 ⁻	03 $\beta^+=94.3$ 5; $\alpha=5.7$ 5
¹⁷⁷ Pt ^m	-29223	15	147.4	0.4	2.2 μ s	0.3	1/2 ⁻	03 IT=100
¹⁷⁷ Au	-21550	13			1.46 s	0.03	(1/2 ⁺ , 3/2 ⁺)	03 01Ko44 TJD $\alpha\approx 100$; $\beta^+ ?$
¹⁷⁷ Au ^m	-21334	28	216	26	1.180 s	0.012	11/2 ⁻	03 01Ko44 ETJ $\alpha\approx 100$; $\beta^+ ?$
¹⁷⁷ Au ⁿ	-21093	28	457	26	7 ns	4	(9/2 ⁻)	03 02Ro17 ETJ IT=100
¹⁷⁷ Hg	-12780	80			127.3 ms	1.8	5/2 ⁻ #	03 $\alpha=85$; $\beta^+=15$
¹⁷⁷ Tl	-3328	25			18 ms	5	(1/2 ⁺)	03 $\alpha=73$ 13; p=27 13
¹⁷⁷ Tl ^m	-2521	17	807	18	230 μ s	40	(11/2 ⁻)	03 p=51 8; $\alpha=49$ 8
* ¹⁷⁷ Au ^m	E : 157.9 keV above 5/2 ⁺ level at estimated 44(28) keV by NUBASE							**
* ¹⁷⁷ Au ⁿ	E : 240.8 keV above 11/2 ⁻ level T : < 15 ns							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{178}Tm	-44120#	400#	30#	s			β^- ?
^{178}Yb	-49698	10	74	m 3	0 ⁺	94	β^- =100
^{178}Lu	-50343.0	2.9	28.4	m 0.2	1 ⁽⁺⁾	94	β^- =100
$^{178}\text{Lu}^m$	-50219	4	123.8	2.6	RQ	94	β^- =100
^{178}Hf	-52444.3	2.1	STABLE			94	IS=27.28 7
$^{178}\text{Hf}^m$	-51296.9	2.1	1147.423	0.005		94	IT=100
$^{178}\text{Hf}^n$	-49998.6	2.1	2445.69	0.11		94	IT=100
$^{178}\text{Hf}^p$	-49870.8	2.2	2573.5	0.5		94	IT=100
^{178}Ta	-50507	15	9.31	m 0.03	1 ⁺	94	β^+ =100
$^{178}\text{Ta}^m$	-50410#	50#	2.36	h 0.08	(7) ⁻	94	β^+ =100
$^{178}\text{Ta}^n$	-48940#	50#	59	ms 3	(15) ⁻	94	IT=100
$^{178}\text{Ta}^p$	-47510#	50#	290	ms 12	(21) ⁻	96Ko13 TJE	*
^{178}W	-50416	15	21.6	d 0.3	0 ⁺	94	ϵ =100
^{178}Re	-45653	28	13.2	m 0.2	(3) ⁺	94	β^+ =100
^{178}Os	-43546	16	5.0	m 0.4	0 ⁺	94	β^+ =100
^{178}Ir	-36252	20	12	s 2		95	β^+ =100
^{178}Pt	-31998	11	21.1	s 0.6	0 ⁺	94	β^+ =92.3 3; α =7.7 3
^{178}Au	-22330	60	2.6	s 0.5		94	β^+ ≤60; α >40
^{178}Hg	-16317	13	269	ms 3	0 ⁺	94	α =?; β^+ =30#
^{178}Tl	-4750#	110#	255	ms 10		02Ro17 TD	α =?; β^+ =47#
^{178}Pb	3568	24	230	μs 150	0 ⁺	01Ro.B T	α ≈100; β^+ ?
* $^{178}\text{Ta}^n$	E : 1470.6 keV above $^{178}\text{Ta}^m$, from ENSDF						
* $^{178}\text{Ta}^n$	T : average 96Ko13=58(4) 79Du02=60(5)						
* $^{178}\text{Ta}^p$	E : 2902 keV above the (7) ⁻ $^{178}\text{Ta}^m$ isomer						
* ^{178}Hg	T : others 96Pa01=287(23) 91Se01=250(25) and 79Ha10=260(30)						
* ^{178}Pb	T : two events at 202 and 147 μs						
^{179}Tm	-41600#	500#	20#	s	1/2 ⁺ #		β^- ?
^{179}Yb	-46420#	300#	8.0	m 0.4	(1/2) ⁻	94	β^- =100
^{179}Lu	-49064	5	4.59	h 0.06	7/2 ⁽⁺⁾	94	β^- =100
$^{179}\text{Lu}^m$	-48472	5	592.4	0.4		94	IT=100
^{179}Hf	-50471.9	2.1	STABLE			94	IS=13.62 2
$^{179}\text{Hf}^m$	-50096.9	2.1	375.0367	0.0025		94	IT=100
$^{179}\text{Hf}^n$	-49366.1	2.1	1105.84	0.19		94	IT=100
^{179}Ta	-50366.3	2.2	1.82	y 0.03	7/2 ⁺	00	ϵ =100
$^{179}\text{Ta}^m$	-49049.0	2.2	1317.3	0.4		00	IT=100
$^{179}\text{Ta}^n$	-47727.0	2.3	2639.3	0.5		00	IT=100
^{179}W	-49304	16	37.05	m 0.16	(7/2) ⁻	94	β^+ =100
$^{179}\text{W}^m$	-49082	16	6.40	m 0.07	(1/2) ⁻	94	IT≈100; β^+ =0.28 3
^{179}Re	-46586	24	19.5	m 0.1	(5/2) ⁺	95	β^+ =100
$^{179}\text{Re}^m$	-46521	24	95	μs 25	(5/2) ⁻		
^{179}Os	-43020	18	6.5	m 0.3	(1/2) ⁻	94	β^+ =100
^{179}Ir	-38077	11	79	s 1	(5/2) ⁻	98	β^+ =100
^{179}Pt	-32264	9	21.2	s 0.4	1/2 ⁻	94	β^+ ≈100; α =0.24 3
^{179}Au	-24952	17	7.1	s 0.3	5/2 ⁻ #	94	β^+ =78.0 9; α =22.0 9
$^{179}\text{Au}^p$	-24853	18			(11/2) ⁻		
^{179}Hg	-16922	27	1.09	s 0.04	5/2 ⁻ #	94	α ≈53; β^+ =?; β^+ p≈0.15
^{179}Tl	-8300	40	270	ms 30	(1/2) ⁺	01	α =?; β^+ =30#
$^{179}\text{Tl}^m$	-7440#	50#	1.60	ms 0.16	(9/2) ⁻	01	α ≈100; IT ?; β^+ ?
^{179}Pb	2000#	200#	3#	ms	5/2 ⁻ #		α ?
* ^{179}Hg	T : average 02Ro17=1.08(0.09) 71Ha03=1.09(0.04)						
* ^{179}Tl	T : average 02Ro17=415(55) 98To14=230(40) 83Sc24=160(+90-40)						
* ^{179}Tl	J : from α decay to $^{175}\text{Au}^m$						
* $^{179}\text{Tl}^m$	T : average 02Ro17=1.7(0.2) 98To14=1.8(0.4) 96Pa01=0.7(+6-4) 83Sc24=1.4(0.5)						

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁸⁰ Yb	-44400#	400#		2.4	m	0.5	0 ⁺	94	β^- =100	
¹⁸⁰ Lu	-46690	70		5.7	m	0.1	5 ⁺	94	95Me03 J β^- =100	
¹⁸⁰ Lu ^m	-46680	70	13.9	0.3	1	s	3 ⁻	95Me03	EJT β^- ?; IT ?	
¹⁸⁰ Hf	-49788.4	2.1			STABLE		0 ⁺	94	IS=35.08 16	
¹⁸⁰ Hf ^m	-48646.9	2.1	1141.48	0.04	5.5	h	0.1	8 ⁻	94 IT \approx 100; β^- =0.3 1	
¹⁸⁰ Ta	-48936.2	2.2			8.152	h	0.006	1 ⁺	94 ϵ =86 3; β^- =14 3	
¹⁸⁰ Ta ^m	-48860.9	1.8	75.3	1.3	RQ	STABLE	(>1.2 Py)	9 ⁻	94 IS=0.012 2; β^- ?	
¹⁸⁰ Ta ⁿ	-47485.2	2.4	1451.0	1.0	45	μ s	2	15 ⁻	96Dr02 TE	
¹⁸⁰ W	-49644	4			STABLE	(>700 Py)	0 ⁺	94	03Da05 T IS=0.12 1; α ?; $2\beta^+$? *	
¹⁸⁰ W ^m	-48115	4	1529.04	0.03	5.47	ms	0.09	8 ⁻	94 IT=100	
¹⁸⁰ Re	-45840	21			2.44	m	0.06	(1) ⁻	94 β^+ =100	
¹⁸⁰ Os	-44359	20			21.5	m	0.4	0 ⁺	94 β^+ =100	
¹⁸⁰ Ir	-37978	22			1.5	m	0.1	(4,5) ⁽⁺⁾	94 β^+ =100	
¹⁸⁰ Pt	-34436	11			52	s	3	0 ⁺	94 $\beta^+\approx$ 100; $\alpha\approx$ 0.3	
¹⁸⁰ Au	-25596	21			8.1	s	0.3	94	$\beta^+\leq$ 98.2; $\alpha\geq$ 1.8	
¹⁸⁰ Hg	-20245	14			2.56	s	0.02	0 ⁺	94 93Wa03 T $\beta^+=$ 52 4; $\alpha=$ 48 4	
¹⁸⁰ Tl	-9400#	120#			1.5	s	0.2	94	98To14 TD β^+ ?; $\alpha=$ 7 3; ... *	
¹⁸⁰ Pb	-1939	21			5	ms	3	0 ⁺	00 96To08 TD $\alpha=$ 100 **	
* ¹⁸⁰ W	T : lower limit is for α decay, also 03Ce01>270 Py 97Ge15>74 Py **									
* ¹⁸⁰ W	T : indication in 03Da05 for 1.1(+0.8-0.4) Ey, but important background **									
* ¹⁸⁰ W	T : 03Da09>80 Py for $2\beta^-$ decay **									
* ¹⁸⁰ Tl	D : ... ; β^+ SF \approx 1.0e-4 **									
* ¹⁸⁰ Tl	D : $\alpha=(2-12)\%$ from 02An.A **									
¹⁸¹ Yb	-40850#	400#			1#	m		3/2 ⁻ #	β^- ?	
¹⁸¹ Lu	-44740#	300#			3.5	m	0.3	(7/2 ⁺)	91 β^- =100	
¹⁸¹ Hf	-47411.9	2.1			42.39	d	0.06	1/2 ⁻	91 β^- =100	
¹⁸¹ Hf ^m	-46817	4	595	3	80	μ s	5	(9/2 ⁺)	01Sh36 ETJ IT=100	
¹⁸¹ Hf ⁿ	-46372	10	1040	10	100	μ s		(17/2 ⁺)	01Sh36 ETJ IT=100	
¹⁸¹ Hf ^p	-45674	10	1738	10	1.5	ms	0.5	(27/2 ⁻)	01Sh36 ETJ IT=100	
¹⁸¹ Ta	-48441.6	1.8			STABLE			7/2 ⁺	92 IS=99.988 2	
¹⁸¹ Ta ^m	-48435.4	1.8	6.238	0.020	6.05	μ s	0.12	9/2 ⁻	92 IT=100	
¹⁸¹ Ta ⁿ	-46957	3	1485	3	25	μ s	2	21/2 ⁻	98Wh02 ETJ IT=100	
¹⁸¹ Ta ^p	-46212	3	2230	3	210	μ s	20	29/2 ⁻	98Wh02 ETJ IT=100	
¹⁸¹ W	-48254	5			121.2	d	0.2	9/2 ⁺	91 $\epsilon=$ 100	
¹⁸¹ Re	-46511	13			19.9	h	0.7	5/2 ⁺	91 β^+ =100	
¹⁸¹ Os	-43550	30			105	m	3	1/2 ⁻	92 β^+ =100	
¹⁸¹ Os ^m	-43500	30	48.9	0.2	2.7	m	0.1	(7/2) ⁻	92 95Ro09 E β^+ =100	
¹⁸¹ Ir	-39472	26			4.90	m	0.15	(5/2) ⁻	93 β^+ =100	
¹⁸¹ Pt	-34375	15			52.0	s	2.2	1/2 ⁻	99 95Bi01 D $\beta^+\approx$ 100; $\alpha=$ 0.074 10	
¹⁸¹ Au	-27871	20			13.7	s	1.4	(3/2 ⁻)	99 $\beta^+=?$; $\alpha=$ 2.7 5	
¹⁸¹ Hg	-20661	15			3.6	s	0.1	1/2 ⁽⁻⁾	99 $\beta^+=$ 69 5; $\alpha=$ 31 5; ... *	
¹⁸¹ Hg ^p	-20460#	40#	210#	40#				13/2 ⁺		
¹⁸¹ Tl	-12801	9			3.2	s	0.3	1/2 ⁺ #	91 98To14 TD $\alpha=?$; β^+ ? *	
¹⁸¹ Tl ^m	-11944	29	857	29	AD	1.7	ms	0.4	9/2 ⁻ #	98To14 TD β^+ ?; $\alpha=?$; IT ? *
¹⁸¹ Pb	-3140	90			&	45	ms	20	5/2 ⁻ #	96To01 T $\alpha=?$; $\beta^+=$ 2# *
¹⁸¹ Pb ^m	non existent		RN	&					13/2 ⁺ #	91 96To01 I *
* ¹⁸¹ Hg	D : ... ; β^+ p=0.016 4; β^+ $\alpha=$ 11e-6 4 **									
* ¹⁸¹ Tl	T : average 98To14=3.2(0.3) 92Bo.D=3.4(0.6) **									
* ¹⁸¹ Tl ^m	T : average 98To14=1.4(0.5) 84Sc.A=2.7(1.0) **									
* ¹⁸¹ Pb	T : supersedes 89To01=50(+40-30) from same group **									
* ¹⁸¹ Pb ^m	I : proved by 96To01 not to exist **									

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)		
¹⁸² Lu	-41880#	200#		2.0	m	0.2	(0, 1, 2)	95	$\beta^- = 100$		
¹⁸² Hf	-46059	6		9	My	2	0 ⁺	95	$\beta^- = 100$		
¹⁸² Hf ^m	-44886	6	1172.88	0.18	61.5	m	1.5	8 ⁻	95	$\beta^- = 58.3$; IT=42.3	
¹⁸² Ta	-46433.3	1.8		114.43	d	0.03	3 ⁻	95	$\beta^- = 100$		
¹⁸² Ta ^m	-46417.0	1.8	16.263	0.003	283	ms	3	5 ⁺	95	IT=100	
¹⁸² Ta ⁿ	-45913.7	1.8	519.572	0.018	15.84	m	0.10	10 ⁻	95	IT=100	
¹⁸² W	-48247.5	0.8			STABLE	(>170 Ey)	0 ⁺	95	03Da05 T	IS=26.50 16; α ?	
¹⁸² Re	-45450	100			64.0	h	0.5	7 ⁺	95	$\beta^+ = 100$	
¹⁸² Re ^m	-45388	20	60	100	BD *	12.7	h	0.2	2 ⁺	95	$\beta^+ = 100$
¹⁸² Os	-44609	22			22.10	h	0.25	0 ⁺	95	$\epsilon = 100$	
¹⁸² Ir	-39052	21			15	m	1	(3 ⁺)	95	95Sa42 J	$\beta^+ = 100$
¹⁸² Pt	-36169	16			2.2	m	0.1	0 ⁺	95	$\beta^+ \approx 100$; $\alpha = 0.038$ 2	
¹⁸² Au	-28301	20			15.5	s	0.4	(2 ⁺)	95	01Ib02 J	$\beta^+ \approx 100$; $\alpha = 0.13$ 5
¹⁸² Hg	-23576	10			10.83	s	0.06	0 ⁺	95	97Ba21 D	$\beta^+ \approx 86.2$ 9; $\alpha = 13.8$ 9; ...
¹⁸² Tl	-13350	80			2.0	s	0.3	2 ⁻ #	95	92Bo.D T	$\beta^+ > 96$; $\alpha < 4$
¹⁸² Tl ^m	-13250#	130#	100#	100#	*	2.9	s	0.5	(7 ⁺)	91Bo22 TJ	$\alpha \approx 100$; $\beta^+ ?$
¹⁸² Tl ^p	-12750#	160#	600#	140#	*						
¹⁸² Pb	-6826	14			60	ms	40	0 ⁺	95	$\alpha = ?$; $\beta^+ = 2\#$	
* ¹⁸² W	T : also 03Ce01 > 25 Ey 97Ge15 > 8.3 Ey									**	
* ¹⁸² Au	T : average 95Bi01=14.5(1.3)(for β^+), 15.3(1.0)(for α) and 92Ro21=15.6(0.4)									**	
* ¹⁸² Hg	D : ... ; $\beta^+ p < 1e-5$									**	
* ¹⁸² Hg	D : α average 97Ba21=13.3(0.5) 80Sc09=15.2(0.8); $\beta^+ p$ is from 71Ho07									**	
* ¹⁸² Tl ^m	T : average 91Bo22=3.1(1.0) 92Bo.D=2.8(0.6)									**	
¹⁸³ Lu	-39520#	300#			58	s	4	(7/2 ⁺)	91	$\beta^- = 100$	
¹⁸³ Hf	-43290	30			1.067	h	0.017	(3/2 ⁻)	91	$\beta^- = 100$	
¹⁸³ Ta	-45296.1	1.8			5.1	d	0.1	7/2 ⁺	91	$\beta^- = 100$	
¹⁸³ Ta ^m	-45222.9	1.8	73.174	0.012	107	ns	11	9/2 ⁻	91	IT=100	
¹⁸³ W	-46367.0	0.8			STABLE	(>80 Ey)	1/2 ⁻	01	03Da05 T	IS=14.31 4; α ?	
¹⁸³ W ^m	-46057.5	0.8	309.493	0.003	5.2	s	0.3	11/2 ⁺	01	IT=100	
¹⁸³ Re	-45811	8			70.0	d	1.4	5/2 ⁺	99	$\epsilon = 100$	
¹⁸³ Re ^m	-43903	8	1907.6	0.3	1.04	ms	0.04	(25/2 ⁺)	99	IT=100	
¹⁸³ Os	-43660	50			13.0	h	0.5	9/2 ⁺	91	$\beta^+ = 100$	
¹⁸³ Os ^m	-43490	50	170.71	0.05	9.9	h	0.3	1/2 ⁻	91	$\beta^+ \approx 85$ 2; IT=15 2	
¹⁸³ Ir	-40197	25			58	m	5	5/2 ⁻	91	61Di04 T	$\beta^+ \approx 100$; $\alpha = 0.05\#$
¹⁸³ Pt	-35772	16			6.5	m	1.0	1/2 ⁻	93	95Bi01 D	$\beta^+ \approx 100$; $\alpha = 0.0096$ 5
¹⁸³ Pt ^m	-35738	16	34.50	0.08	43	s	5	(7/2 ⁻)	93	$\beta^+ \approx 100$; $\alpha < 4e-4$; IT ?	
¹⁸³ Au	-30187	10			42.8	s	1.0	5/2 ⁻	99	94Pa37 J	$\beta^+ \approx 100$; $\alpha = 0.55$ 25
¹⁸³ Au ^m	-30114	10	73.3	0.4	> 1	μ s		(1/2 ⁺)	99	IT=100	
¹⁸³ Au ^p	-29956	10	230.6	0.6	< 1	μ s		(11/2 ⁻)	99	IT=100	
¹⁸³ Hg	-23800	8			9.4	s	0.7	1/2 ⁻	01	$\beta^+ \approx 88.3$ 20; $\alpha = 11.7$ 20; ...	
¹⁸³ Hg ^m	-23560#	40#	240#	40#	EU	5#	s	13/2 ⁺ #	01Sc41 I	$\beta^+ ?$	
¹⁸³ Hg ^p	-23602	13	198	14	AD			13/2 ⁺ #			
¹⁸³ Tl	-16587	10			6.9	s	0.7	1/2 ⁺ #	02	$\beta^+ = ?$; $\alpha = 2\#$	
¹⁸³ Tl ^m	-15944	16	643	14	AD	60	ms	15	9/2 ⁻ #	02	$\alpha \approx 1.5$; $\beta^+ ?$; IT ?
¹⁸³ Tl ⁿ	-15611	20	976.8	17	1.48	μ s	0.10	(13/2 ⁺)	02	01Mu26 EJ	IT=100
¹⁸³ Pb	-7569	28			535	ms	30	(3/2 ⁻)	03	$\alpha = ?$; $\beta^+ = 10\#$	
¹⁸³ Pb ^m	-7475	28	94	8	AD	415	ms	20	(13/2 ⁺)	03	$\alpha \approx 100$; $\beta^+ ?$
* ¹⁸³ W	T : also 03Ce01 > 13 Ey 97Ge15 > 1.9 Ey									**	
* ¹⁸³ Ir	T : average 61Di04=55(7) 61La05=60(6)									**	
* ¹⁸³ Hg	D : ... ; $\beta^+ p = 2.6e-4$ 8									**	
* ¹⁸³ Hg ^m	I : 2001Sc41= no isomer seen with same characteristics as ¹⁸⁵ Hg or ¹⁸⁷ Hg									**	
* ¹⁸³ Hg ^m	I : no isomer in same odd-N ¹⁸¹ Pt and ¹⁷⁹ Os									**	
* ¹⁸³ Tl ⁿ	E : 346.8(0.3) keV above ¹⁸³ Tl ^m									**	

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
^{184}Lu	-36410#	400#		20	s	3	(3 ⁺)	90 95Kr04 TJ	β^- =100
$^{184}\text{Lu}^m$			non existent	20	s		high	95Kr04 I	
^{184}Hf	-41500	40		4.12	h	0.05	0 ⁺	90	β^- =100
$^{184}\text{Hf}^m$	-40230	40	1272.4 0.4	48	s	10	8 ⁻	95Kr04 TE	β^- =100
^{184}Ta	-42841	26		8.7	h	0.1	(5 ⁻)	90	β^- =100
^{184}W	-45707.3	0.9		STABLE	(>180 Ey)		0 ⁺	90 03Da05 T	IS=30.64 2; α ? *
^{184}Re	-44227	4		38.0	d	0.5	3 ⁽⁻⁾	90	β^+ =100
$^{184}\text{Re}^m$	-44039	4	188.01 0.04	169	d	8	8 ⁽⁺⁾	90	IT=75.4 11; ϵ =24.6 11
^{184}Os	-44256.1	1.3		STABLE	(>56 Ty)		0 ⁺	90	IS=0.02 1; α ?; $2\beta^+$? *
^{184}Ir	-39611	28		3.09	h	0.03	5 ⁻	90	β^+ =100
$^{184}\text{Ir}^m$	-39385	28	225.65 0.11	470	μs		3 ⁺		
^{184}Pt	-37332	18		17.3	m	0.2	0 ⁺	90 95Bi01 D	β^+ \approx 100; α =0.0017 7
$^{184}\text{Pt}^m$	-35493	18	1839.4 1.6	1.01	ms	0.05	8 ⁻	90	IT=100
^{184}Au	-30319	22		20.6	s	0.9	5 ⁺	03	β^+ \approx 100; α <0.016
$^{184}\text{Au}^m$	-30251	22	68.46 0.01	47.6	s	1.4	2 ⁺	03 94Ib01 EJ	β^+ =?; IT=30 10; α <0.016
$^{184}\text{Au}^n$	-30091	22	228.40 0.06	69	ns	6	3 ⁻	03	IT=100
^{184}Hg	-26349	10		30.6	s	0.3	0 ⁺	90	β^+ =98.89 6; α =1.11 6
^{184}Tl	-16890	50		9.7	s	0.6	2 ⁻ #	90 92Bo.D T	β^+ =97.9 7; α =2.1 7
$^{184}\text{Tl}^m$	-16790#	110#	100# 100#	10#	s		7 ⁺ #		β^+ ?; IT ?
$^{184}\text{Tl}^n$	-16390#	150#	500# 140#	> 20	ns		(10 ⁻)	84Sc.A T	IT ? *
^{184}Pb	-11045	14		490	ms	25	0 ⁺	03 02An.A D	α =80 15; β^+ ?
^{184}Bi	1050#	130#		6.6	ms	1.5	3 ⁺ #	02An.A T	α =?
$^{184}\text{Bi}^m$	1200#	160#	150# 100#	13	ms	2	10 ⁻ #	02An.A T	α =?
* ^{184}W	T : also 03Ce01>29 Ey 97Ge15>4.0 Ey **								
* ^{184}Os	T : lower limit is for α decay **								
* $^{184}\text{Tl}^n$	T : alpha decay from $^{188}\text{Bi}^m$ not coincident with X(K) and γ **								
* $^{184}\text{Tl}^n$	I : identified by 02Sc.A **								
^{185}Hf	-38360#	200#		3.5	m	0.6	3/2 ⁻ #	95	β^- =100
^{185}Ta	-41396	14		49.4	m	1.5	7/2 ⁺ #	95	β^- =100
$^{185}\text{Ta}^m$	-40090	30	1308 29	> 1	ms		(21/2 ⁻)	99Wh03 TJD	IT=100 *
^{185}W	-43389.7	0.9		75.1	d	0.3	3/2 ⁻	95	β^- =100
$^{185}\text{W}^m$	-43192.3	0.9	197.43 0.05	1.597	m	0.004	11/2 ⁺	95 94It.A T	IT=100
^{185}Re	-43822.2	1.2		STABLE			5/2 ⁺	95	IS=37.40 2
$^{185}\text{Re}^m$	-41698.2	2.3	2124 2	123	ns	23	(21/2)	97Sh37 T	IT=100
^{185}Os	-42809.4	1.3		93.6	d	0.5	1/2 ⁻	95	ϵ =100
$^{185}\text{Os}^m$	-42707.1	1.5	102.3 0.7	3.0	μs	0.4	7/2 ⁻ #	95	IT ?
^{185}Ir	-40336	28		14.4	h	0.1	5/2 ⁻	95	β^+ =100
^{185}Pt	-36680	40		70.9	m	2.4	(9/2 ⁺)	95	β^+ \approx 100; α =0.0050 20 *
$^{185}\text{Pt}^m$	-36580	40	103.4 0.2	33.0	m	0.8	(1/2 ⁻)	95	β^+ =?; IT<2
^{185}Au	-31867	26		4.25	m	0.06	5/2 ⁻	95	β^+ \approx 100; α =0.26 6
$^{185}\text{Au}^m$	-31770#	100#	100# 100#	6.8	m	0.3	1/2 ⁺ #	95	β^+ <100; IT ?
^{185}Hg	-26176	16		49.1	s	1.0	1/2 ⁻	95	β^+ =94 1; α =6 1
$^{185}\text{Hg}^m$	-26072	16	103.8 1.0	21.6	s	1.5	13/2 ⁺	95 87Ki.A E	IT=54 10; β^+ =46 10; α \approx 0.03 *
^{185}Tl	-19760	50		19.5	s	0.5	1/2 ⁺ #	95	β^+ =?; α ?
$^{185}\text{Tl}^m$	-19300	50	452.8 2.0	1.83	s	0.12	9/2 ⁻ #	95 77Sc03 E	IT \approx 100; α =0.10 3; β^+ ?
$^{185}\text{Tl}^n$	-18760	50	1003.0 2.0	8.3	ns	1.4	(13/2 ⁺)	95La08 T	
^{185}Pb	-11541	16		6.3	s	0.4	3/2 ⁻	95 02An15 TJD	α =50 25; β^+ ? *
$^{185}\text{Pb}^m$	-11480#	40#	60# 40#	4.07	s	0.15	13/2 ⁺	02An15 TJD	α =50 25; β^+ ? *
^{185}Bi	-2210#	50#		&	2#	ms	9/2 ⁻ #	96Da06 J	p ?; α ? *
$^{185}\text{Bi}^m$	-2143	18	70# 50#	&	49	μs	7	01Po05 T	p=85 6; α =15 6 *
* $^{185}\text{Ta}^m$	E : from 99Wh03 : less than 100 keV above 1258 level J : assuming ground-state=7/2 ⁺ **								
* ^{185}Pt	D : if the 4444(10) keV α line is from ground-state; otherwise α =0.0010(4)% from isomer **								
* $^{185}\text{Hg}^m$	E : ENSDF gives 99.3(0.5) plus "8-keV uncertainty", but missed 87Ki.A work **								
* ^{185}Pb	T : average 02An15=6.3(0.4) 80Sc09=6.1(1.1) **								
* $^{185}\text{Pb}^m$	T : average 02An15=4.3(0.2) 80Sc09=3.73(0.24) (excluding the 6.1 s activity) **								
* ^{185}Bi	T : estimated from 9/2 ⁻ isomers in odd Bi and Tl isotopes **								
* $^{185}\text{Bi}^m$	T : average 01Po05=50(8) 96Da06=44(16) **								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{186}Hf	-36430#	300#	2.6 m	1.2	0 ⁺	03	$\beta^- = 100$
^{186}Ta	-38610	60	10.5 m	0.3	(2 ⁻ , 3 ⁻)	03	$\beta^- = 100$
^{186}W	-42509.5	1.7	STABLE	(>4.1 Ey)	0 ⁺	03	03Da09 T IS=28.43 19; $2\beta^- ?$; $\alpha ?$ *
$^{186}\text{W}^m$	-40992.3	1.8	1517.2	0.6	18	μs	1 (7 ⁻) 03 IT=100
$^{186}\text{W}^n$	-38966.7	2.7	3542.8	2.1	> 3	ms	(16 ⁺) 03 IT=100 *
^{186}Re	-41930.2	1.2	3.7183 d	0.0011	1 ⁻	03	$\beta^- = 92.53$ 10; $\epsilon = 7.47$ 10
$^{186}\text{Re}^m$	-41781	7	149	7	200	ky	50 (8 ⁺) 03 IT=?; $\beta^- < 10$ *
^{186}Os	-42999.5	1.4	2.0	Py	1.1	0 ⁺	03 IS=1.59 3; $\alpha = 100$
^{186}Ir	-39173	17	16.64	h	0.03	5 ⁺	03 $\beta^+ = 100$
$^{186}\text{Ir}^m$	-39172	17	0.8	0.4	1.92	h	0.05 2 ⁻ 03 91Be25 ET $\beta^+ \approx 75$; IT ≈ 25 *
^{186}Pt	-37864	22	2.08	h	0.05	0 ⁺	03 $\beta^+ = 100$; $\alpha \approx 1.4e-4$
^{186}Au	-31715	21	10.7	m	0.5	3 ⁻	03 $\beta^+ = 100$; $\alpha = 0.0008$ 2
$^{186}\text{Au}^m$	-31487	21	227.77	0.07	110	ns	10 2 ⁺ 03 IT=100
$^{186}\text{Au}^p$			non existent	RN	< 2	m	
^{186}Hg	-28539	11	1.38	m	0.06	0 ⁺	03 $\beta^+ \approx 100$; $\alpha = 0.016$ 5
$^{186}\text{Hg}^m$	-26322	11	2217.3	0.4	82	μs	5 (8 ⁻) 03 IT=100
^{186}Tl	-20190	180			40#	s	(2 ⁻) 03 91Va04 I $\beta^+ ?$ *
$^{186}\text{Tl}^m$	-19874	9	320	180	AD	*	& 27.5 s 1.0 (7 ⁺) 03 $\beta^+ \approx 100$; $\alpha \approx 0.006$
$^{186}\text{Tl}^n$	-19501	9	690	180	AD		2.9 s 0.2 (10 ⁻) 03 IT=100 *
^{186}Pb	-14681	11	4.82	s	0.03	0 ⁺	03 $\beta^+ ?$; $\alpha = 40$ 8
^{186}Bi	-3170	80			14.8	ms	0.7 (3 ⁺) 03 02An.A T $\alpha \approx 100$; $\beta^+ ?$ *
$^{186}\text{Bi}^m$	-2900#	160#	270#	140#	*	9.8	ms 0.4 (10 ⁻) 03 02An.A T $\alpha \approx 100$; $\beta^+ ?$
* ^{186}W	T : limit is $2\beta^-$ decay; 03Da05>170 Ey 03Ce01>27 Ey 97Ge15>6.5 Ey for α decay **						
* $^{186}\text{W}^n$	T : lower limit is 3 ms; upper limit 30 s **						
* $^{186}\text{Re}^m$	T : uncertainty estimated by ENSDF'89 evaluator **						
* $^{186}\text{Ir}^m$	T : average 91Be25=1.90(0.05) 70Fi.A=2.0(0.1) **						
* $^{186}\text{Ir}^n$	E : E is positive and below 1.5 keV **						
* ^{186}Tl	I : identified as decay level from ^{190}Bi in 91Va04 **						
* $^{186}\text{Tl}^n$	E : 374.0(0.2) keV above $^{186}\text{Tl}^m$ **						
* ^{186}Bi	T : average 02An.A=14.8(0.8) 97Ba21=15.0(1.7) **						
^{187}Hf	-32980#	400#	30#	s	(>300 ns)	3/2 ⁻ #	99Be63 I $\beta^- ?$
^{187}Ta	-36770#	200#	2#	m	(>300 ns)	7/2 ⁺ #	99Be63 I $\beta^- ?$
^{187}W	-39904.8	1.7	23.72	h	0.06	3/2 ⁻	92 $\beta^- = 100$
^{187}Re	-41215.7	1.4	41.2	Gy	0.2	5/2 ⁺	91 01Ga01 T IS=62.60 2; $\beta^- = 100$; ... *
^{187}Os	-41218.2	1.4	STABLE			1/2 ⁻	92 IS=1.96 2
^{187}Ir	-39716	6	10.5	h	0.3	3/2 ⁺	91 $\beta^+ = 100$
$^{187}\text{Ir}^m$	-39530	6	186.15	0.04	30.3	ms	0.6 9/2 ⁻ 91 IT=100
^{187}Pt	-36713	28	2.35	h	0.03	3/2 ⁻	91 $\beta^+ = 100$
^{187}Au	-33005	25	8.4	m	0.3	1/2 ⁺	91 $\beta^+ \approx 100$; $\alpha = 0.003\#$
$^{187}\text{Au}^m$	-32884	25	120.51	0.16	2.3	s	0.1 9/2 ⁻ 91 IT=100
^{187}Hg	-28118	14			1.9	m	0.3 3/2 ⁻ 91 $\beta^+ = 100$; $\alpha > 1.2e-4$
$^{187}\text{Hg}^m$	-28059	20	59	16	MD	&	2.4 m 0.3 13/2 ⁺ 91 $\beta^+ = 100$; $\alpha > 2.5e-4$
^{187}Tl	-22444	8			51	s	(1/2 ⁺) 99 $\beta^+ < 100$; $\alpha ?$
$^{187}\text{Tl}^m$	-22109	8	335	3	AD		15.60 s 0.12 (9/2 ⁻) 99 IT=?; $\beta^+ ?$; $\alpha = 0.15$ 5
^{187}Pb	-14980	8			15.2	s	0.3 (3/2 ⁻) 00 $\beta^+ = 93$ 2; $\alpha = 7$ 2
$^{187}\text{Pb}^m$	-14969	11	11	11	AD	*	18.3 s 0.3 (13/2 ⁺) 00 $\beta^+ = 88$ 2; $\alpha = 12$ 2
^{187}Bi	-6373	15			32	ms	3 9/2 ⁻ # 01 $\alpha > 50$; $\beta^+ ?$
$^{187}\text{Bi}^m$	-6272	18	101	20	AD		320 μs 70 1/2 ⁺ # 01 $\alpha > 50$; $\beta^+ ?$
$^{187}\text{Bi}^n$	-6121	15	252	1	7	μs	5 (13/2 ⁺) 02Hu14 ETJ IT=100
* ^{187}Re	D : ... ; $\alpha < 0.0001$ **						
* ^{187}Re	T : others: 89Li30=42.3(0.7) outweighed and, same group, 86Li11=43.5(1.3) **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
¹⁸⁸ Hf	−30880#	500#	20# s (>300 ns)	0 ⁺	02	99Be63 I	β^- ?	
¹⁸⁸ Ta	−33810#	200#	20# s (>300 ns)	0 ⁺	02	99Be63 I	β^- ?	
¹⁸⁸ W	−38667	3	69.78 d	0.05	0 ⁺	02	β^- =100	
¹⁸⁸ Re	−39016.1	1.4	17.0040 h	0.0022	1 ⁻	02	β^- =100	
¹⁸⁸ Re ^m	−38844.0	1.4	172.069	0.009	18.59 m	0.04	(6) ⁻ 02 IT=100	
¹⁸⁸ Os	−41136.4	1.4	STABLE		0 ⁺	02	IS=13.24 8	
¹⁸⁸ Ir	−38328	7	41.5 h	0.5	1 ⁻	02	β^+ =100	
¹⁸⁸ Ir ^m	−37360	30	970	30	4.2 ms	0.2	7 ⁺ # 02 ABBW E IT≈100; β^+ ? *	
¹⁸⁸ Pt	−37823	5	10.2 d	0.3	0 ⁺	02	ϵ =100; α =2.6e−5 3	
¹⁸⁸ Au	−32301	20	8.84 m	0.06	1 ⁽⁻⁾	02	β^+ =100	
¹⁸⁸ Hg	−30202	12	3.25 m	0.15	0 ⁺	02	β^+ =100; α =3.7e−5 8	
¹⁸⁸ Hg ^m	−27478	12	2724.3	0.4	134 ns	15	(12 ⁺) 02 IT=100	
¹⁸⁸ Tl	−22350	30		*	71 s	2	(2 ⁻) 02 β^+ =100	
¹⁸⁸ Tl ^m	−22307	10	40	30	MD *	71 s	1	(7 ⁺) 02 β^+ =100
¹⁸⁸ Tl ⁿ	−22038	10	310	30	MD	41 ms	4	(9 ⁻) 02 IT≈100; β^+ ? *
¹⁸⁸ Pb	−17815	11			25.5 s	0.1	0 ⁺	02 β^+ ?; α =9.3 8
¹⁸⁸ Pb ^m	−15237	11	2578.2	0.7	830 ns	210	(8 ⁻) 02 IT=100	
¹⁸⁸ Pb ⁿ	−15102	11	2713.0	0.6	94 ns		(11 ⁻) 02 IT=100	
¹⁸⁸ Pb ^p	−15020	50	2800	50	797 ns	21		02 IT=100 *
¹⁸⁸ Bi	−7200	50		*	& 44 ms	3	3 ⁺ # 02 97Wa05 T α ?; β^+ ? *	
¹⁸⁸ Bi ^m	−7000#	150#	210#	140#	*	& 220 ms	40	(10 ⁻) 02 97Wa05 T α ?; β^+ ? *
¹⁸⁸ Po	−538	19			430 μ s	180	0 ⁺	02 α ?; β^+ ?
* ¹⁸⁸ Ir ^m	E : less than 100 keV above 923.5 level, from ENSDF							**
* ¹⁸⁸ Tl ⁿ	E : 268.8(0.5) keV above ¹⁸⁸ Tl ^m , from 91Va04							**
* ¹⁸⁸ Pb ^p	E : 2700.5 above unknown level, see ENSDF'02							**
* ¹⁸⁸ Bi	T : average 97Wa05=46(7) 84Sc.A=44(3)							**
* ¹⁸⁸ Bi ^m	T : average 97Wa05=218(50) 84Sc.A=210(90)							**
¹⁸⁹ Ta	−31830#	300#	3# s (>300 ns)	7/2 ⁺ #	99Be63 I	91	β^- ?	
¹⁸⁹ W	−35480	200	11.6 m	0.3	(3/2 ⁻)	91	97Ya03 T β^- =100 *	
¹⁸⁹ Re	−37978	8	24.3 h	0.4	5/2 ⁺	91	β^- =100	
¹⁸⁹ Os	−38985.4	1.5	STABLE		3/2 ⁻	91	IS=16.15 5	
¹⁸⁹ Os ^m	−38954.6	1.5	30.814	0.018	5.8 h	0.1	9/2 ⁻ 91 IT=100	
¹⁸⁹ Ir	−38453	13			13.2 d	0.1	3/2 ⁺ 91 ϵ =100	
¹⁸⁹ Ir ^m	−38081	13	372.18	0.04	13.3 ms	0.3	11/2 ⁻ 91 IT=100	
¹⁸⁹ Ir ⁿ	−36120	13	2333.3	0.4	3.7 ms	0.2	(25/2) ⁺ 91 IT=100	
¹⁸⁹ Pt	−36483	11			10.87 h	0.12	3/2 ⁻ 92 β^+ =100	
¹⁸⁹ Pt ^m	−36291	11	191.6	0.4	143 μ s		(13/2 ⁺)	
¹⁸⁹ Au	−33582	20			28.7 m	0.3	1/2 ⁺ 92 β^+ =100; α <3e−5	
¹⁸⁹ Au ^m	−33335	20	247.23	0.17	4.59 m	0.11	11/2 ⁻ 92 β^+ ≈100; IT=?	
¹⁸⁹ Hg	−29630	30			7.6 m	0.1	3/2 ⁻ 96 β^+ =100; α <3e−5	
¹⁸⁹ Hg ^m	−29549	18	80	30	MD	8.6 m	0.1	13/2 ⁺ 96 01Sc41 E β^+ =100; α <3e−5
¹⁸⁹ Tl	−24602	11			2.3 m	0.2	(1/2 ⁺) 99 β^+ =100	
¹⁸⁹ Tl ^m	−24319	10	283	6	AD	1.4 m	0.1	9/2 ⁽⁻⁾ 99 85Bo46 J β^+ ≈100; IT<4
¹⁸⁹ Pb	−17880	30			*	51 s	3	(3/2 ⁻) 91 ABBW J β^+ >99; α ≈0.4 *
¹⁸⁹ Pb ^m	−17840#	50#	40#	30#	*	1# m	(13/2 ⁺) ABBW J β^+ ?; IT ? *	
¹⁸⁹ Bi	−10060	50			674 ms	11	(9/2 ⁻) 98 95Ba75 J α >50; β^+ <50 *	
¹⁸⁹ Bi ^m	−9880	50	181	6	AD	6.6 ms	0.6	(1/2 ⁺) 98 95Ba75 TJ α >50; β^+ <50 *
¹⁸⁹ Bi ⁿ	−9700	50	357	1	880 ns	50	(13/2 ⁺) 01An11 ETJ IT=100 *	
¹⁸⁹ Po	−1415	22			5 ms	1	3/2 ⁻ # 99An52 TD α ?; β^+ ?	
* ¹⁸⁹ W	T : average 97Ya03=11.7(0.5) 65Ka07=11.5(0.3)							**
* ¹⁸⁹ Pb	J : from α decay to ¹⁸⁵ Hg							**
* ¹⁸⁹ Pb ^m	J : from α decay from ¹⁹³ Po ^m							**
* ¹⁸⁹ Bi	T : average 02Hu14=667(13) 97Wa05=728(40) 85Co06=680(30)							**
* ¹⁸⁹ Bi ^m	T : average 97An09=4.8(0.5) 97Wa05=5.2(0.6) 95Ba75=7.0(0.2)							**
* ¹⁸⁹ Bi ⁿ	T : from 02Hu14; also 01An11>360(120)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁹⁰ Ta	-28660#	400#			300# ms				$\beta^- ?$
¹⁹⁰ W	-34300	160			30.0 m	1.5		03	$\beta^-=100$
¹⁹⁰ W ^m	-31920	160	2381	5	< 3.1 ms			(10 ⁻) 03	IT=100
¹⁹⁰ Re	-35570	150			3.1 m	0.3		(2) ⁻ 03	$\beta^-=100$
¹⁹⁰ Re ^m	-35360	160	210	60	3.2 h	0.2		(6 ⁻) 03	$\beta^-=54.4$ 20; IT ?
¹⁹⁰ Os	-38706.3	1.5			STABLE			03	IS=26.26 2
¹⁹⁰ Os ^m	-37000.9	1.5	1705.4	0.2	9.9 m	0.1		(10) ⁻ 03	IT=100
¹⁹⁰ Ir	-36751.2	1.7			11.78 d	0.10		4 ⁻ 03	$\beta^+=100$; $e^+ < 0.002$
¹⁹⁰ Ir ^m	-36725.1	1.7	26.1	0.1	1.120 h	0.003		(1 ⁻) 03	IT=100
¹⁹⁰ Ir ⁿ	-36374.8	1.7	376.4	0.1	3.087 h	0.012		(11) ⁻ 03	$\beta^+=91.4$ 2; IT=8.6 2
¹⁹⁰ Ir ^p	-36715.0	1.7	36.154	0.025	> 2 μ s			(4) ⁺ 03	IT=100
¹⁹⁰ Ir ^q	-36433.6	1.7	317.56	0.04	90 ns			(5 ⁻) 03	IT=100
¹⁹⁰ Pt	-37323	6			650 Gy	30		0 ⁺ 03	IS=0.014 1; $\alpha=100$...
¹⁹⁰ Au	-32881	16			* 42.8 m	1.0		1 ⁻ 03	$\beta^+=100$; $\alpha < 1e-6$
¹⁹⁰ Au ^m	-32680#	150#	200#	150#	* 125 ms	20		11-# 03	IT \approx 100; $\beta^+ ?$
¹⁹⁰ Hg	-31370	16			20.0 m	0.5		0 ⁺ 03	$\epsilon \approx 100$; $e^+ < 1$; ...
¹⁹⁰ Tl	-24330	50			* 2.6 m	0.3		2 ⁽⁻⁾ 03	$\beta^+=100$
¹⁹⁰ Tl ^m	-24200#	70#	130#	90#	* 3.7 m	0.3		7 ⁽⁺⁾ 03	$\beta^+=100$
¹⁹⁰ Tl ⁿ	-24040#	90#	290#	70#	750 μ s	40		(8 ⁻) 03	IT=100
¹⁹⁰ Tl ^p	-23920#	90#	410#	70#	> 1 μ s			9 ⁻ 03	IT ?
¹⁹⁰ Pb	-20417	12			71 s	1		0 ⁺ 03	$\beta^+ ?$; $\alpha=0.40$ 4
¹⁹⁰ Pb ^m	-17802	12	2614.8	0.8	150 ns			(10) ⁺ 03	IT=100
¹⁹⁰ Pb ⁿ	-17799	23	2618	20	25 μ s			(12) ⁺ 03	IT ?
¹⁹⁰ Pb ^p	-17759	12	2658.2	0.8	7.2 μ s	0.6		(11) ⁻ 03	IT=100
¹⁹⁰ Bi	-10900	180			6.3 s	0.1		(3 ⁺) 03	$\alpha=77$ 21; $\beta^+=?$
¹⁹⁰ Bi ^m	-10483	10	420	180	MD 6.2 s	0.1		(10 ⁻) 03	$\alpha=70$ 9; $\beta^+ ?$
¹⁹⁰ Bi ⁿ	-10210	10	690	180	MD > 500 ns	100		7 ⁺ # 03	IT=100
¹⁹⁰ Po	-4563	13			2.46 ms	0.05		0 ⁺ 03	$\alpha \approx 100$; $\beta^+ = 0.1\%$
* ¹⁹⁰ Re ^m	E : from lower limit 119.12 and calculated 173 and 220 (see ENSDF'90)								
* ¹⁹⁰ Re ⁿ	E : 210(290) from difference in beta-decay								
* ¹⁹⁰ Pt	D : ... ; $2\beta^+ ?$								
* ¹⁹⁰ Hg	D : ... ; $\alpha < 3.4e-7$								
* ¹⁹⁰ Tl ⁿ	E : 161.9 keV above ¹⁹⁰ Tl ^m								
* ¹⁹⁰ Tl ^p	E : 236.2 keV above ¹⁹⁰ Tl ^m								
* ¹⁹⁰ Pb ⁿ	E : above ¹⁹⁰ Pb ^m , see ENSDF'03								
* ¹⁹⁰ Bi ⁿ	E : 273(1) keV above the (10 ⁻) isomer								
¹⁹¹ W	-31110#	200#			20# s (>300 ns)			3/2 ⁻ #	99Be63 I $\beta^- ?$
¹⁹¹ Re	-34349	10			9.8 m	0.5		(3/2 ⁺ , 1/2 ⁺) 95	$\beta^-=100$
¹⁹¹ Os	-36393.7	1.5			15.4 d	0.1		9/2 ⁻ 95	$\beta^-=100$
¹⁹¹ Os ^m	-36319.3	1.5	74.382	0.003	13.10 h	0.05		3/2 ⁻ 95	IT=100
¹⁹¹ Ir	-36706.4	1.7			STABLE			3/2 ⁺ 95	IS=37.3 2
¹⁹¹ Ir ^m	-36535.2	1.7	171.24	0.05	4.94 s	0.03		11/2 ⁻ 95	IT=100
¹⁹¹ Ir ⁿ	-34590	40	2120	40	5.5 s	0.7		95	IT=100
¹⁹¹ Pt	-35698	4			2.802 d	0.025		3/2 ⁻ 96	$\epsilon=100$
¹⁹¹ Pt ^m	-35549	4	149.04	0.02	95 μ s			13/2 ⁺	
¹⁹¹ Au	-33810	40			3.18 h	0.08		3/2 ⁺ 99	$\beta^+=100$
¹⁹¹ Au ^m	-33540	40	266.2	0.5	920 ms	110		(11/2 ⁻) 99	IT=100
¹⁹¹ Hg	-30593	23			49 m	10		3/2 ⁽⁻⁾ 00	$\beta^+=100$; $\alpha < 5e-6$
¹⁹¹ Hg ^m	-30470	30	128	22	50.8 m	1.5		13/2 ⁺ 00	$\beta^+=100$; $\alpha < 5e-6$
¹⁹¹ Tl	-26281	8			20# m			(1/2 ⁺) 95	$\beta^+ ?$
¹⁹¹ Tl ^m	-25984	7	297	7	BD 5.22 m	0.16		9/2 ⁽⁻⁾ 95	$\beta^+=100$
¹⁹¹ Pb	-20250	40			* 1.33 m	0.08		(3/2 ⁻) 95	$\beta^+ \approx 100$; $\alpha=0.013$ 5
¹⁹¹ Pb ^m	-20231	28	20	50	MD * 2.18 m	0.08		13/2 ⁽⁺⁾ 95	$\beta^+ \approx 100$; $\alpha \approx 0.02$
¹⁹¹ Bi	-13240	7			12.3 s	0.3		(9/2 ⁻) 00	$\alpha=60$ 20; $\beta^+=40$ 20
¹⁹¹ Bi ^m	-13000	9	240	4	AD 124 ms	5		(1/2 ⁺) 00	$\alpha=75$ 25; $\beta^+ \approx 25$
¹⁹¹ Po	-5054	11			22 ms	1		3/2 ⁻ # 00	$\alpha \approx 100$; $\beta^+ ?$
¹⁹¹ Po ^m	-5020	10	34	12	AD 98 ms	8		(13/2 ⁺) 00	$\alpha \approx 100$; $\beta^+ ?$
* ¹⁹¹ Ir ⁿ	E : estimated less than 150 keV above 2047.1 level, from ENSDF								
* ¹⁹¹ Hg ^m	E : original error (8 keV) increased by 20 for isomer+ground-state lines in trap								
* ¹⁹¹ Bi	T : average 03Ke04=12.4(0.4) 85Co06=12(1) 74Le02=13(1) 72Ga27=12.0(0.7)								
* ¹⁹¹ Bi ^m	T : average 03Ke04=121(+8-5) 99An36=115(10) 81Le23=150(15)								

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
¹⁹² W	−29650# 600#		10# s (>300 ns)	0 ⁺		99Be63 I	β^- ?
¹⁹² Re	−31710# 200#		16 s	1		98	β^- =100
¹⁹² Os	−35880.5 2.6		STABLE (>9.8 Ty)	0 ⁺		98	IS=40.78 19; $2\beta^-$?; α ? *
¹⁹² Os ^m	−33865.1 2.6	2015.40	0.11	5.9 s	0.1	(10 ⁻)	98 IT>87; β^- <13
¹⁹² Ir	−34833.2 1.7		73.827 d	0.013		4 ⁺	98 β^- =95.13 14; ϵ =4.87 14
¹⁹² Ir ^m	−34776.5 1.7	56.720	0.005	1.45 m	0.05	1 ⁻	98 IT≈100; β^- =0.0175
¹⁹² Ir ⁿ	−34665.1 1.7	168.14	0.12	241 y	9	(11 ⁻)	98 IT=100
¹⁹² Pt	−36292.9 2.5		STABLE	0 ⁺		98	IS=0.782 7
¹⁹² Au	−32777 16		4.94 h	0.09		1 ⁻	98 β^+ =100
¹⁹² Au ^m	−32642 16	135.41	0.25	29 ms		5# ⁺	98 IT=100
¹⁹² Au ⁿ	−32345 16	431.6	0.5	160 ms	20	(11 ⁻)	98 IT=100
¹⁹² Hg	−32011 16		4.85 h	0.20		0 ⁺	00 ϵ =100; α <4e−6
¹⁹² Tl	−25870 30		9.6 m	0.4		(2 ⁻)	99 β^+ =100
¹⁹² Tl ^m	−25710 60	160	50	10.8 m	0.2	(7 ⁺)	99 91Va04 E β^+ =100
¹⁹² Tl ^p	−25694 25	180	40	AD		(3 ⁺)	99 91Va04 E
¹⁹² Pb	−22556 13		3.5 m	0.1		0 ⁺	98 β^+ ≈100; α =0.0059 7
¹⁹² Pb ^m	−19975 13	2581.1	0.1	164 ns	7	(10 ⁺)	98 IT=100
¹⁹² Pb ⁿ	−19931 13	2625.1	1.1	1.1 μs	0.5	(12 ⁺)	98 IT=100
¹⁹² Pb ^p	−19813 13	2743.5	0.4	756 ns	21	(11 ⁻)	98 IT=100
¹⁹² Bi	−13550 30		34.6 s	0.9		(3 ⁺)	98 β^+ =88 5; α =12 5
¹⁹² Bi ^m	−13399 9	150	30	MD		(10 ⁻)	98 β^+ =90 3; α =10 3
¹⁹² Po	−8071 12		32.2 ms	0.3		0 ⁺	98 99He32 T α =?; β^+ =0.5# *
¹⁹² Po ^m	−5470# 500# 2600# 500#		1 μs			12 ⁺ #	99He32 T IT=100
* ¹⁹² Os	T : lower limit is for 0v-2 β^- decay **						
* ¹⁹² Po	T : others 98A127=31(4) 96Bi17=33.2(1.4) 81Le23=34(3) outweighed, not used **						
¹⁹³ Re	−30300# 200#		30# s (>300 ns)	5/2 ⁺ #		99Be63 I	β^- ?
¹⁹³ Os	−33392.6 2.6		30.11 h	0.01		3/2 ⁻	98 β^- =100
¹⁹³ Ir	−34533.8 1.7		STABLE			3/2 ⁺	98 IS=62.7 2
¹⁹³ Ir ^m	−34453.6 1.7	80.240	0.006	10.53 d	0.04	11/2 ⁻	98 IT=100
¹⁹³ Pt	−34477.0 1.7		50 y	6		1/2 ⁻	98 ϵ =100
¹⁹³ Pt ^m	−34327.2 1.7	149.78	0.04	4.33 d	0.03	13/2 ⁺	98 IT=100
¹⁹³ Au	−33394 11		17.65 h	0.15		3/2 ⁺	98 β^+ =100; α <1e−5
¹⁹³ Au ^m	−33104 11	290.19	0.03	3.9 s	0.3	11/2 ⁻	98 IT≈100; β^+ ≈0.03
¹⁹³ Hg	−31051 15		3.80 h	0.15		3/2 ⁻	99 β^+ =100
¹⁹³ Hg ^m	−30910 15	140.76	0.05	11.8 h	0.2	13/2 ⁺	99 β^+ =92.8 5; IT=7.2 5
¹⁹³ Tl	−27320 110		21.6 m	0.8		1/2 ⁽⁺⁾ #	99 β^+ =100
¹⁹³ Tl ^m	−26950 110	369	4	2.11 m	0.15	9/2 ⁻	99 IT=75; β^+ =25 *
¹⁹³ Pb	−22190 50		* 5# m			(3/2 ⁻)	99 ABBW J β^+ ? *
¹⁹³ Pb ^m	−22060# 90# 130# 80#		* 5.8 m	0.2		13/2 ⁽⁺⁾	99 88Me.A J β^+ =100
¹⁹³ Bi	−15873 10		67 s	3		(9/2 ⁻)	98 β^+ ?; α =3.5 15
¹⁹³ Bi ^m	−15564 12	308	7	AD		(1/2 ⁺)	98 α =90 20; β^+ ?
¹⁹³ Po	−8360 30		420 ms	40		3/2 ⁻ #	98 α =?; β^+ =5#
¹⁹³ Po ^m	−8260# 50# 100# 30#		240 ms	10		(13/2 ⁺)	98 ABBW J α =?; β^+ =3#
¹⁹³ At	−150 50		40 ms			9/2 ⁻ #	98 α =100
* ¹⁹³ Tl ^m	E : less than 13 keV above 362.5 level, from ENSDF **						
* ¹⁹³ Pb	J : from α decay from ¹⁹⁷ Po **						
* ¹⁹³ Pb	T : T=4.0 m reported in Karlsruhe charts 1981 and 1995. Not traceable **						
¹⁹⁴ Re	−27550# 300#		2# s (>300 ns)			99Be63 I	β^- ?
¹⁹⁴ Os	−32432.7 2.6		6.0 y	0.2		0 ⁺	96 β^- =100
¹⁹⁴ Ir	−32529.3 1.7		19.28 h	0.13		1 ⁻	96 β^- =100
¹⁹⁴ Ir ^m	−32382.2 1.7	147.078	0.005	31.85 ms	0.24	(4 ⁺)	96 IT=100
¹⁹⁴ Ir ⁿ	−32160 70	370	70	BD		(10, 11) ^(−) #	96 β^- =100
¹⁹⁴ Pt	−34763.1 0.9		STABLE			0 ⁺	96 IS=32.967 99
¹⁹⁴ Au	−32262 10		38.02 h	0.10		1 ⁻	96 β^+ =100
¹⁹⁴ Au ^m	−32155 10	107.4	0.5	600 ms	8	(5 ⁺)	96 IT=100
¹⁹⁴ Au ⁿ	−31786 10	475.8	0.6	420 ms	10	(11 ⁻)	96 IT=100
¹⁹⁴ Hg	−32193 13		440 y	80		0 ⁺	01 ϵ =100

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J ^π	En	Reference	Decay modes and intensities (%)	
... A-group continued ...								
¹⁹⁴ Tl	-26830	140		*	33.0	m	0.5 2 ⁻ 99 β ⁺ =100; α<1e-7	
¹⁹⁴ Tl ^m	-26530#	240#	300#	200#	*	32.8	m	0.2 (7 ⁺) 99 β ⁺ =100
¹⁹⁴ Pb	-24208	17			12.0	m	0.5 0 ⁺ 99 β ⁺ =100; α=7.3e-6 29	
¹⁹⁴ Bi	-15990	50			95	s	3 (3 ⁺) 96 β ⁺ ≈100; α=0.46 25	
¹⁹⁴ Bi ^m	-15880	50	110	70	MD *	125	s	2 (6 ⁺ , 7 ⁺) 96 β ⁺ ≈100; α ?
¹⁹⁴ Bi ⁿ	-15760#	70#	230#	90#		115	s	4 (10 ⁻) 96 β ⁺ ≈100; α=0.20 7
¹⁹⁴ Po	-11005	13			392	ms	4 0 ⁺ 96 α≈100; β ⁺ ?	
¹⁹⁴ Po ^m	-8480	13	2525	2	15	μs	2 (11 ⁻) 99He32 TJD IT=100	
¹⁹⁴ At	-1190	190			40	ms	3 ⁺ # 96 α≈100; β ⁺ ?	
¹⁹⁴ At ^m	-711	17	480	190	AD	250	ms	10 ⁻ # 96 α≈100; IT ?
¹⁹⁵ Os	-29690	500			6.5	m	3/2 ⁻ # 99 β ⁻ =100 *	
¹⁹⁵ Ir	-31689.8	1.7			2.5	h	0.2 3/2 ⁺ 99 β ⁻ =100	
¹⁹⁵ Ir ^m	-31590	5	100	5	3.8	h	0.2 11/2 ⁻ 99 β ⁻ ≈95 5; IT=5 5	
¹⁹⁵ Pt	-32796.8	0.9			STABLE		1/2 ⁻ 99 IS=33.832 10	
¹⁹⁵ Pt ^m	-32537.5	0.9	259.30	0.08	4.02	d	0.01 13/2 ⁺ 99 IT=100	
¹⁹⁵ Au	-32570.0	1.3			186.10	d	0.05 3/2 ⁺ 99 ε=100	
¹⁹⁵ Au ^m	-32251.4	1.3	318.58	0.04	30.5	s	0.2 11/2 ⁻ 99 IT=100	
¹⁹⁵ Hg	-31000	23			10.53	h	0.03 1/2 ⁻ 99 01Li17 T β ⁺ =100	
¹⁹⁵ Hg ^m	-30824	23	176.07	0.04	41.6	h	0.8 13/2 ⁺ 99 IT=54.2 20; β ⁺ =45.8 20	
¹⁹⁵ Tl	-28155	14			1.16	h	0.05 1/2 ⁺ 99 β ⁺ =100	
¹⁹⁵ Tl ^m	-27672	14	482.63	0.17	3.6	s	0.4 9/2 ⁻ 99 IT=100	
¹⁹⁵ Pb	-23714	23			15	m	3/2 ⁻ # 99 β ⁺ =100	
¹⁹⁵ Pb ^m	-23511	23	202.9	0.7	15.0	m	1.2 13/2 ⁺ 99 β ⁺ =100	
¹⁹⁵ Bi	-18024	6			183	s	4 (9/2 ⁻) 99 ABBW J β ⁺ ≈100; α=0.03 2	
¹⁹⁵ Bi ^m	-17624	8	399	6	AD	87	s	1 (1/2 ⁺) 99 ABBW J β ⁺ =67 17; α=33 17 *
¹⁹⁵ Po	-11070	40			4.64	s	0.09 3/2 ⁻ # 99 α=75 15; β ⁺ =25 15	
¹⁹⁵ Po ^m	-10964	28	110	50	AD	1.92	s	0.02 13/2 ⁺ # 99 α≈90; β ⁺ ≈10; IT<0.01
¹⁹⁵ At	-3476	9			& 328	ms	20 (1/2 ⁺) 00 03Ke04 T α≈100; β ⁺ ?	
¹⁹⁵ At ^m	-3443	8	34	7	AD &	147	ms	5 9/2 ⁻ # 00 03Ke04 T α=?; β ⁺ <25#
¹⁹⁵ Rn	5070	50			*	6	ms	3/2 ⁻ # 01Ke06 TD α=?
¹⁹⁵ Rn ^m	5118	15	50	50	*	6	ms	13/2 ⁺ # 01Ke06 TD α=?
* ¹⁹⁵ Os	I : identification of this nuclide has been questioned, see ENSDF'99							**
* ¹⁹⁵ Bi ^m	J : spins of ground-state and of isomer derived from alpha decay							**
¹⁹⁶ Os	-28280	40			34.9	m	0.2 0 ⁺ 98 β ⁻ =100	
¹⁹⁶ Ir	-29440	40			52	s	1 (0 ⁻) 98 β ⁻ =100	
¹⁹⁶ Ir ^m	-29229	20	210	40	BD	1.40	h	0.02 (10, 11 ⁻) 98 β ⁻ ≈100; IT<0.3
¹⁹⁶ Pt	-32647.4	0.9			STABLE		0 ⁺ 98 IS=25.242 41	
¹⁹⁶ Au	-31140.0	3.0			6.1669	d	0.0006 2 ⁻ 98 01Li17 T β ⁺ ≈92.8 8; β ⁻ =7.2 8	
¹⁹⁶ Au ^m	-31055	3	84.660	0.020	8.1	s	0.2 5 ⁺ 98 IT=100	
¹⁹⁶ Au ⁿ	-30544	3	595.66	0.04	9.6	h	0.1 12 ⁻ 98 IT=100	
¹⁹⁶ Hg	-31826.7	2.9			STABLE	(>2.5 Ey)	0 ⁺ 98 90Bu28 T IS=0.15 1; 2β ⁺ ?	
¹⁹⁶ Tl	-27497	12			1.84	h	0.03 2 ⁻ 98 β ⁺ =100	
¹⁹⁶ Tl ^m	-27103	12	394.2	0.5	1.41	h	0.02 (7 ⁺) 98 β ⁺ ≈95.5; IT=4.5	
¹⁹⁶ Pb	-25361	14			37	m	3 0 ⁺ 01 β ⁺ =100; α≤3e-5	
¹⁹⁶ Pb ^m	-23623	14	1738.27	0.12	< 1	μs	4 ⁺ 01 IT=100	
¹⁹⁶ Bi	-18009	24			5.1	m	0.2 (3 ⁺) 99 β ⁺ ≈100; α=0.00115 34	
¹⁹⁶ Bi ^m	-17842	25	166.6	3.0	AD	0.6	s	0.5 (7 ⁺) 99 IT=?; β ⁺ ?
¹⁹⁶ Bi ⁿ	-17739	25	270	3	AD	4.00	m	0.05 (10 ⁻) 99 β ⁺ =74.2 25; IT=25.8 25;... *
¹⁹⁶ Po	-13474	13			5.56	s	0.12 0 ⁺ 98 93Wa04 TD α=94 5; β ⁺ =6 5 *	
¹⁹⁶ Po ^m	-10984	13	2490.5	1.7	850	ns	90 (11 ⁻) 98 IT=100	
¹⁹⁶ At	-3920	60			*	253	ms	9 3 ⁺ # 98 97Pu01 T α=?; β ⁺ =4#
¹⁹⁶ At ^m	-3950	50	-30	80	AD *	20#	ms	10 ⁻ # 96En01 D IT ?
¹⁹⁶ At ⁿ	-3760	60	157.9	0.1	11	μs	5 ⁺ # 00Sm06 ET IT ?	
¹⁹⁶ Rn	1970	15			4.7	ms	1.1 0 ⁺ 98 01Ke06 T α≈100; β ⁺ =0.2#	
* ¹⁹⁶ Bi ⁿ	D : ... ; α=0.00038 10							**
* ¹⁹⁶ Po	T : average 97Pu01=5.5(0.1) 93Wa04=5.8(0.2)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^{π}	Ens	Reference	Decay modes and intensities (%)
^{197}Ir	-28268	20	5.8 m	0.5	3/2 ⁺	96	$\beta^{-}=100$
$^{197}\text{Ir}^m$	-28153	21	115	5	8.9 m	0.3	11/2 ⁻ 96 $\beta^{-}\approx 100$; IT=0.25 10
^{197}Pt	-30422.4	0.8	19.8915 h	0.0019	1/2 ⁻	96	$\beta^{-}=100$
$^{197}\text{Pt}^m$	-30022.8	0.8	399.59	0.20	95.41 m	0.18	13/2 ⁺ 96 IT=96.7 4; $\beta^{-}=3.3$ 4
^{197}Au	-31141.1	0.6	STABLE		3/2 ⁺	96	IS=100.
$^{197}\text{Au}^m$	-30732.0	0.6	409.15	0.08	7.73 s	0.06	11/2 ⁻ 96 IT=100
^{197}Hg	-30541	3	64.94 h	0.07	1/2 ⁻	96	01Li17 T $\varepsilon=100$ *
$^{197}\text{Hg}^m$	-30242	3	298.93	0.08	23.8 h	0.1	13/2 ⁺ 96 IT=91.4 7; $\varepsilon=8.6$ 7
^{197}Tl	-28341	16	2.84 h	0.04	1/2 ⁺	96	$\beta^{+}=100$
$^{197}\text{Tl}^m$	-27733	16	608.22	0.08	540 ms	10	9/2 ⁻ 96 IT=100
^{197}Pb	-24749	6	8 m	2	3/2 ⁻	01	$\beta^{+}=100$
$^{197}\text{Pb}^m$	-24429	6	319.31	0.11	43 m	1	13/2 ⁺ 01 $\beta^{+}=81$ 2; IT=19 2; ... *
$^{197}\text{Pb}^n$	-22835	6	1914.10	0.25	1.15 μs	0.20	21/2 ⁻ 01 IT=100
^{197}Bi	-19688	8	9.3 m	0.5	(9/2 ⁻)	99	$\beta^{+}=100$; $\alpha=1\text{e-}4\#$
$^{197}\text{Bi}^m$	-19000	110	690	110	AD	5.04 m	0.16 (1/2 ⁺) 99 $\alpha=55$ 40; $\beta^{+}=45$ 40; ... *
^{197}Po	-13360	50	90#	230#	80#	53.6 s	1.0 (3/2 ⁻) 96 β^{+} ?; $\alpha=44$ 7
$^{197}\text{Po}^m$	-13120#	90#	230#	80#	25.8 s	0.1	(13/2 ⁺) 96 $\alpha=84$ 9; β^{+} ?; IT=0.01#
^{197}At	-6340	50	*		350 ms	40	(9/2 ⁻) 96 $\alpha=96$ 4; $\beta^{+}=4$ 4
$^{197}\text{Au}^m$	-6293	13	50	50	AD *	3.7 s	2.5 (1/2 ⁺) 96 $\alpha\approx 100$; β^{+} ?; IT<0.004
^{197}Rn	1480	60	1670#	200#	60#	66 ms	16 (3/2 ⁻) 98 $\alpha\approx 100$; β^{+} ? *
$^{197}\text{Rn}^m$	1670#	50#	200#	60#	21 ms	5	(13/2 ⁺) 98 $\alpha\approx 100$; β^{+} ? *
* ^{197}Hg	T : other 66El09=64.14(0.05) at strong variance: Birge ratio would be $B=9.3$ **						
* $^{197}\text{Pb}^m$	D : ... ; $\alpha<3\text{e-}4$ **						
* $^{197}\text{Bi}^m$	D : ... ; IT<0.3 **						
* ^{197}Rn	T : average 96En02=65(+25-14) 95Mo14=51(+35-15) **						
* $^{197}\text{Rn}^m$	T : average 96En02=19(+8-4) 95Mo14=18(+9-5) J : from α decay to $^{193}\text{Po}^m$ **						
^{198}Ir	-25820#	200#			8 s	1	02 $\beta^{-}=100$
^{198}Pt	-29908	3			STABLE	(>320 Ty)	0 ⁺ 02 52Fr23 T IS=7.163 55; $2\beta^{-}$?; α ? *
^{198}Au	-29582.1	0.6			2.69517 d	0.00021	2 ⁻ 02 $\beta^{-}=100$
$^{198}\text{Au}^m$	-29269.9	0.6	312.2200	0.0020	124 ns	4	5 ⁺ 02 IT=100
$^{198}\text{Au}^n$	-28770.4	1.6	811.7	1.5	2.27 d	0.02	(12 ⁻) 02 IT=100
^{198}Hg	-30954.4	0.3			STABLE		0 ⁺ 02 IS=9.97 20
^{198}Tl	-27490	80			5.3 h	0.5	2 ⁻ 02 $\beta^{+}=100$
$^{198}\text{Tl}^m$	-26950	80	543.5	0.4	1.87 h	0.03	7 ⁺ 02 $\beta^{+}=54$ 2; IT=46 2
$^{198}\text{Tl}^n$	-26750	80	742.3	0.4	32.1 ms	1.0	10 ⁻ # 02 IT=100
^{198}Pb	-26050	15			2.4 h	0.1	0 ⁺ 02 $\beta^{+}=100$
$^{198}\text{Pb}^m$	-23909	15	2141.4	0.4	4.19 μs	0.10	(7 ⁻) 02 IT=100
^{198}Bi	-19369	28			10.3 m	0.3	(2 ⁺ , 3 ⁺) 02 $\beta^{+}=100$
$^{198}\text{Bi}^m$	-19085	28	280	40	MD	11.6 m	0.3 (7 ⁺) 02 $\beta^{+}=100$
$^{198}\text{Bi}^n$	-18837	28	530	40	MD	7.7 s	0.5 10 ⁻ 02 IT=100 *
^{198}Po	-15473	17			1.77 m	0.03	0 ⁺ 02 $\alpha=57$ 2; $\beta^{+}=43$ 2
$^{198}\text{Po}^m$	-13619	17	1853.63	0.18	29 ns	2	8 ⁺ 02 IT=100
$^{198}\text{Po}^n$	-12907	17	2565.92	0.20	200 ns	20	11 ⁻ 02 IT=100
$^{198}\text{Po}^p$	-12781	17	2691.86	0.20	750 ns	50	12 ⁺ 02 IT ?
^{198}At	-6670	50			4.2 s	0.3	(3 ⁺) 02 95Bi.A D $\alpha>94$; β^{+} ?
$^{198}\text{At}^m$	-6340#	70#	330#	90#	1.0 s	0.2	(10 ⁻) 02 95Bi.A D $\alpha>86$; β^{+} ?
^{198}Rn	-1231	13			65 ms	3	0 ⁺ 02 $\alpha=?$; $\beta^{+}=1\#$
$^{198}\text{Rn}^m$			non existent	EU	50 ms	9	$\alpha=?$; $\beta^{+}=?$; IT=? *
* ^{198}Pt	T : lower limit is for $0\nu-2\beta^{-}$ decay **						
* $^{198}\text{Bi}^n$	E : 248.5(0.5) keV above $^{198}\text{Bi}^m$, from 92Hu04 **						
* $^{198}\text{Rn}^m$	I : α decay assigned to isomer by ENSDF'95, not accepted by NUBASE **						
^{199}Ir	-24400	40			20# s		3/2 ⁺ # 01 β^{-} ?
^{199}Pt	-27392	3			30.80 m	0.21	5/2 ⁻ 94 $\beta^{-}=100$
$^{199}\text{Pt}^m$	-26968	4	424	2	13.6 s	0.4	(13/2 ⁺) 94 IT=100
^{199}Au	-29095.0	0.6			3.139 d	0.007	3/2 ⁺ 94 $\beta^{-}=100$
$^{199}\text{Au}^m$	-28546.1	0.6	548.9368	0.0021	440 μs	30	(11/2 ⁻) 94 IT=100
^{199}Hg	-29547.1	0.4			STABLE		1/2 ⁻ 94 IS=16.87 22
$^{199}\text{Hg}^m$	-29014.6	0.4	532.48	0.10	42.66 m	0.08	13/2 ⁺ 94 01Li17 T IT=100 *

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...										
¹⁹⁹ Tl	-28059	28			7.42 h	0.08	1/2 ⁺	94	$\beta^+=100$	
¹⁹⁹ Tl ^m	-27309	28	749.7	0.3	28.4 ms	0.2	9/2 ⁻	94	IT=100	
¹⁹⁹ Pb	-25228	26			90 m	10	3/2 ⁻	01	$\beta^+=100$	
¹⁹⁹ Pb ^m	-24799	26	429.5	2.7	12.2 m	0.3	(13/2 ⁺)	01	IT=93; $\beta^+=7$	
¹⁹⁹ Pb ⁿ	-22664	26	2563.8	2.7	10.1 μ s	0.2	(29/2 ⁻)	01	IT=100	
¹⁹⁹ Bi	-20798	12			27 m	1	9/2 ⁻	94	$\beta^+=100$	
¹⁹⁹ Bi ^m	-20131	12	667	4	24.70 m	0.15	(1/2 ⁺)	94	$\beta^+=?$; IT<2; $\alpha \approx 0.01$	
¹⁹⁹ Po	-15215	23			5.48 m	0.16	(3/2 ⁻)	94	$\beta^+=92.5$ 3; $\alpha=7.5$ 3	
¹⁹⁹ Po ^m	-14903	23	312.0	2.8	AD	4.17 m	0.04	13/2 ⁺	94	$\beta^+=73.5$ 10; $\alpha=24$ 1; IT=2.5
¹⁹⁹ At	-8820	50			7.2 s	0.5	(9/2 ⁻)	94	$\alpha=89$ 6; $\beta^+?$	
¹⁹⁹ Rn	-1520	60			620 ms	30	3/2 ⁻ #	98	$\alpha=?$; $\beta^+=6$ #	
¹⁹⁹ Rn ^m	-1334	29	180	70	AD	320 ms	20	13/2 ⁺ #	98	$\alpha=?$; $\beta^+=3$ #
¹⁹⁹ Fr	6760	40			16 ms	7	1/2 ⁺ #	01	99Ta20 T $\alpha \approx 100$; $\beta^+?$	
* ¹⁹⁹ Hg ^m	T : average 01Li17=42.67(0.09) 69KI06=42.6(0.2)									**
* ¹⁹⁹ Pb ^m	E : 424.8 γ to level lower than 9.3 keV, from ENSDF									**
* ¹⁹⁹ Pb ⁿ	E : 2559.1 to level lower than 9.3 keV, from ENSDF									**
²⁰⁰ Pt	-26603	20			12.5 h	0.3	0 ⁺	95	$\beta^-=100$	
²⁰⁰ Au	-27270	50			48.4 m	0.3	1 ⁽⁻⁾	95	$\beta^-=100$	
²⁰⁰ Au ^m	-26300	50	970	70	BD	18.7 h	0.5	12 ⁻	95	$\beta^-=82$ 2; IT=18 2
²⁰⁰ Hg	-29504.1	0.4			STABLE		0 ⁺	95	IS=23.10 19	
²⁰⁰ Tl	-27048	6			26.1 h	0.1	2 ⁻	95	$\beta^+=100$	
²⁰⁰ Tl ^m	-26294	6	753.6	0.2	34.3 ms	1.0	7 ⁺	95	IT=100	
²⁰⁰ Pb	-26243	11			21.5 h	0.4	0 ⁺	95	$\varepsilon=100$	
²⁰⁰ Bi	-20370	24			* 36.4 m	0.5	7 ⁺	95	$\beta^+=100$	
²⁰⁰ Bi ^m	-20270#	70#	100#	70#	* 31 m	2	(2 ⁺)	95	$\beta^+>90$; IT<10	
²⁰⁰ Bi ⁿ	-19942	24	428.20	0.10	400 ms	50	(10 ⁻)	95	IT=100	
²⁰⁰ Po	-16954	14			11.5 m	0.1	0 ⁺	95	$\beta^+=88.9$ 3; $\alpha=11.1$ 3	
²⁰⁰ At	-8988	24			43.2 s	0.9	(3 ⁺)	95	96Ta18 T $\alpha=57$ 6; $\beta^+=43$ 6	
²⁰⁰ At ^m	-8875	25	112.7	3.0	AD	47 s	1	(7 ⁺)	95	$\alpha=43$ 7; $\beta^+=?$; IT?
²⁰⁰ At ⁿ	-8644	24	344	3	AD	3.5 s	0.2	(10 ⁻)	95	IT \approx 84; $\alpha \approx 10.5$; $\beta^+ \approx 4.5$
²⁰⁰ Rn	-4006	13			1.03 s	0.05	0 ⁺	98	96Ta18 T $\alpha=?$; $\beta^+=2$ #	
²⁰⁰ Fr	6120	80			* 24 ms	10	3 ⁺ #	97	96En01 TD $\alpha=100$	
²⁰⁰ Fr ^m	6180	70	60	110	AD *	650 ms	210	10 ⁻ #	97	95Mo14 TD $\alpha \approx 100$; IT?
* ²⁰⁰ At	T : average 96Ta18=44(2) 92Hu04=43(1)									**
* ²⁰⁰ At ⁿ	E : 230.9(0.2) keV above ²⁰⁰ At ^m , from ENSDF									**
* ²⁰⁰ Rn	T : average 96Ta18=0.96(0.03) 84Ca32=1.06(0.02)									**
²⁰¹ Pt	-23740	50			2.5 m	0.1	(5/2 ⁻)	94	$\beta^-=100$	
²⁰¹ Au	-26401	3			26 m	1	3/2 ⁺	94	$\beta^-=100$	
²⁰¹ Hg	-27663.3	0.6			STABLE		3/2 ⁻	94	IS=13.18 9	
²⁰¹ Hg ^m	-26897.1	0.6	766.23	0.15	94 μ s		13/2 ⁺			
²⁰¹ Tl	-27182	15			72.912 h	0.017	1/2 ⁺	94	$\varepsilon=100$	
²⁰¹ Tl ^m	-26263	15	919.50	0.09	2.035 ms	0.007	(9/2 ⁻)	94	IT=100	
²⁰¹ Pb	-25258	22			9.33 h	0.03	5/2 ⁻	94	$\beta^+=100$	
²⁰¹ Pb ^m	-24629	22	629.14	0.17	61 s	2	13/2 ⁺	94	IT>99; $\beta^+<1$	
²⁰¹ Bi	-21416	15			108 m	3	9/2 ⁻	94	$\beta^+=100$; $\alpha<1e-4$	
²⁰¹ Bi ^m	-20570	15	846.34	0.21	59.1 m	0.6	1/2 ⁺	94	$\beta^+=92.9$ #; IT<6.8; $\alpha=?$	
²⁰¹ Po	-16525	6			15.3 m	0.2	3/2 ⁻	94	$\beta^+=98.4$ 3; $\alpha=1.6$ 3	
²⁰¹ Po ^m	-16101	6	424.1	2.4	AD	8.9 m	0.2	13/2 ⁺	94	IT=56 14; $\beta^+=41$ 10; $\alpha \approx 2.9$
²⁰¹ At	-10789	8			85 s	3	(9/2 ⁻)	94	96Ta18 T $\alpha=71$ 7; $\beta^+=29$ 7	
²⁰¹ Rn	-4070	70			7.0 s	0.4	(3/2 ⁻)	94	96Ta18 T $\alpha=?$; $\beta^+=20$ #	
²⁰¹ Rn ^m	-3790#	90#	280#	90#	3.8 s	0.1	(13/2 ⁺)	94	96Ta18 T $\alpha=?$; $\beta^+=10$ #; IT=0.01#	
²⁰¹ Fr	3600	70			61 ms	12	(9/2 ⁻)	94	96En01 T $\alpha \approx 100$; $\beta^+<1$	
* ²⁰¹ Bi ^m	D : α decay is observed. Its branching ratio is estimated 0.3%# in ENSDF									**
* ²⁰¹ At	T : average 96Ta18=83(2) and two results in ENSDF=89(3)									**
* ²⁰¹ Rn	T : average 96Ta18=7.1(0.8) 71Ho01=7.0(0.4)									**
* ²⁰¹ Fr	T : average 96En01=69(+16-11) 80Ew03=48(15)									**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
²⁰² Pt	-22600#	300#			44 h	15	0 ⁺	97		β^- =100
²⁰² Au	-24400	170			28.8 s	1.9	(1 ⁻)	97		β^- =100
²⁰² Hg	-27345.9	0.6			STABLE		0 ⁺	97		IS=29.86 26 *
²⁰² Tl	-25983	15			12.23 d	0.02	2 ⁻	97		β^+ =100
²⁰² Tl ^m	-25033	15	950.19	0.10	572 μ s	7	7 ⁺	97		
²⁰² Pb	-25934	8			52.5 ky	2.8	0 ⁺	97		ϵ ≈100; α <1#
²⁰² Pb ^m	-23764	8	2169.83	0.07	3.53 h	0.01	9 ⁻	97		IT=90.5 5; β^+ =9.5 5
²⁰² Bi	-20733	20			1.72 h	0.05	5 ⁽⁺⁾	97		β^+ =100; α <1e-5 *
²⁰² Bi ^m	-20118	21	615	7	3.04 μ s	0.06	(10#) ⁻	97		
²⁰² Po	-17924	15			44.7 m	0.5	0 ⁺	97		β^+ =?; α =1.92 7
²⁰² Po ^m	-15297	15	2626.7	0.7	> 200 ns		11 ⁻	97		IT=100
²⁰² At	-10591	28			184 s	1	(2,3) ⁺	97		β^+ =?; α =18 3
²⁰² At ^m	-10401	28	190	40 MD	182 s	2	(7 ⁺)	97		IT ?; β^+ ?; α =8.7 15
²⁰² At ⁿ	-10010	28	580	40 MD	460 ms	50	(10 ⁻)	97	92Hu04 E	IT≈100; β^+ =0.25#; ... *
²⁰² Rn	-6275	18			9.94 s	0.18	0 ⁺	97	96Ta18 T	α =?; β^+ =14# *
²⁰² Fr	3140	50			290 ms	30	(3 ⁺)	97	96En01 T	α =?; β^+ =3# *
²⁰² Fr ^m	3470#	70#	330#	90#	340 ms	40	(10 ⁻)	97		α =?; β^+ =3#
²⁰² Ra	9210	60			2.6 ms	2.1	0 ⁺	98	96Le09 TD	α =100
* ²⁰² Hg	D : lower half-life limit for ²⁴ Ne decay $T>3.7$ Zy, from 90Bu28 **									
* ²⁰² Bi	J : re-evaluation to a possible 6 ⁺ is discussed in 96Ca02 **									
* ²⁰² At ⁿ	D : ... ; α =0.096 11 **									
* ²⁰² At ⁿ	E : 391.7(0.5) keV above ²⁰² At ^m **									
* ²⁰² Rn	T : average 96Ta18=10.3(0.4) 71Ho01=9.85(0.20) **									
* ²⁰² Fr	T : average 96En01=230(+80-40) 95Bi.A=300(40) **									
²⁰³ Au	-23143	3			53 s	2	3/2 ⁺	93		β^- =100
²⁰³ Hg	-25269.1	1.7			46.612 d	0.018	5/2 ⁻	93		β^- =100
²⁰³ Hg ^m	-24336.0	2.0	933.1	1.0	24 μ s		(13/2 ⁺)			
²⁰³ Tl	-25761.2	1.3			STABLE		1/2 ⁺	93		IS=29.524 14
²⁰³ Tl ^m	-22360	300	3400	300	7.7 μ s	0.5	(25/2 ⁺)	98Pf02 TJ		IT=100
²⁰³ Pb	-24787	7			51.873 h	0.009	5/2 ⁻	93		ϵ =100
²⁰³ Pb ^m	-23962	7	825.20	0.09	6.3 s	0.2	13/2 ⁺	93		IT=100
²⁰³ Pb ⁿ	-21838	7	2949.47	0.22	480 ms	20	29/2 ⁻	93		IT=100
²⁰³ Bi	-21540	22			11.76 h	0.05	9/2 ⁻	93		β^+ =100; α ≈1e-5
²⁰³ Bi ^m	-20442	22	1098.14	0.07	303 ms	5	1/2 ⁺	93		IT=100
²⁰³ Po	-17307	26			36.7 m	0.5	5/2 ⁻	93		β^+ ≈100; α =0.11 2
²⁰³ Po ^m	-16666	26	641.49	0.17	45 s	2	13/2 ⁺	93		IT≈100; α =0.04#
²⁰³ At	-12163	12			7.4 m	0.2	9/2 ⁻	93		β^+ =69 3; α =31 3
²⁰³ Rn	-6160	24			43.5 s	2.1	(3/2,5/2) ⁻	93	96Ta18 T	α =66 9; β^+ =34 9 *
²⁰³ Rn ^m	-5798	24	363	4 AD	26.7 s	0.5	13/2 ⁽⁺⁾	93	87Bo29 J	α =?; β^+ =20# *
²⁰³ Fr	861	16			550 ms	20	9/2 ⁻ #	98		α =?; β^+ =5#
²⁰³ Ra	8640	80			4 ms	3	(3/2 ⁻)	98	96Le09 TJD	α ≈100; β^+ ?
²⁰³ Ra ^m	8860	40	220	90 AD	41 ms	17	(13/2 ⁺)	98	96Le09 TJD	α ≈100; β^+ ?
* ²⁰³ Rn	T : average 96Ta18=42(3) 71Ho01=45(3) **									
* ²⁰³ Rn ^m	T : from 96Ta18 **									
²⁰⁴ Au	-20750#	200#			39.8 s	0.9	(2 ⁻)	94		β^- =100
²⁰⁴ Hg	-24690.2	0.3			STABLE		0 ⁺	94		IS=6.87 15; 2 β^- ?
²⁰⁴ Tl	-24346.0	1.3			3.78 y	0.02	2 ⁻	94		β^- =97.10 12; ϵ =2.90 12
²⁰⁴ Tl ^m	-23242.0	1.4	1104.0	0.4	63 μ s	2	(7 ⁺)	94		IT=100
²⁰⁴ Tl ⁿ	-21850	500	2500	500	2.6 μ s	0.2	(12 ⁻)	98Pf02 TJ		IT=100
²⁰⁴ Tl ^p	-20850	500	3500	500	1.6 μ s	0.2	(20 ⁺)	98Pf02 TJ		IT=100
²⁰⁴ Pb	-25109.7	1.2			STABLE	(>140 Py)	0 ⁺	94		IS=1.4 1; α ?
²⁰⁴ Pb ^m	-22923.9	1.2	2185.79	0.05	67.2 m	0.3	9 ⁻	94		IT=100
²⁰⁴ Bi	-20667	26			11.22 h	0.10	6 ⁺	94		β^+ =100
²⁰⁴ Bi ^m	-19862	26	805.5	0.3	13.0 ms	0.1	10 ⁻	94		IT=100
²⁰⁴ Bi ⁿ	-17834	26	2833.4	1.1	1.07 ms	0.03	(17 ⁺)	94		IT=100
²⁰⁴ Po	-18334	11			3.53 h	0.02	0 ⁺	94		β^+ =99.34 1; α =0.66 1

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...								
^{204}At	-11875	24		9.2 m	0.2	7 ⁺	94	$\beta^+=96.2$ 2; $\alpha=3.8$ 2
$^{204}\text{At}^m$	-11288	24	587.30	0.20				IT=100
^{204}Rn	-7984	15						$\alpha=73$ 1; β^+ ?
^{204}Fr	608	25						$\alpha=96$ 2; β^+ ?
$^{204}\text{Fr}^m$	658	25	50	4	AD		94 95Bi.A	D $\alpha=90$ 2; β^+ ?
$^{204}\text{Fr}^n$	934	25	326	4	AD		94 94Le05	T $\alpha=74$ 8; IT=26 8
^{204}Ra	6054	15						$\alpha\approx 100$; $\beta^+=0.3$ #
* $^{204}\text{Fr}^n$	E : 276.1 keV above $^{204}\text{Fr}^m$, from 95Bi.A				D : α intensity is from 95Bi.A			
* ^{204}Ra	T : average 95Le04=45(+55-21) 96Le09=59(+12-9)							
^{205}Au	-18750#	300#						$\beta^-=100$
^{205}Hg	-22287	4						$\beta^-=100$
$^{205}\text{Hg}^m$	-20730	4	1556.53	0.24				IT=100
^{205}Tl	-23820.6	1.3						IS=70.476 14
$^{205}\text{Tl}^m$	-20530.0	1.3	3290.63	0.17	STABLE			IT=100
^{205}Pb	-23770.1	1.2						$\epsilon=100$
$^{205}\text{Pb}^m$	-22756.3	1.2	1013.839	0.013				IT=100
$^{205}\text{Pb}^n$	-20574.5	1.4	3195.6	0.8				IT=100
^{205}Bi	-21062	7						$\beta^+=100$
^{205}Po	-17509	20						$\beta^+\approx 100$; $\alpha=0.04$ 1
$^{205}\text{Po}^m$	-16048	20	1461.20	0.21				IT=100
$^{205}\text{Po}^n$	-16629	20	880.30	0.04				
^{205}At	-12972	15						$\beta^+=90$ 2; $\alpha=10$ 2
$^{205}\text{At}^m$	-10909	15	2062.57	0.25				
$^{205}\text{At}^n$	-10632	15	2339.60	0.25				
^{205}Rn	-7710	50						$\beta^+=77$ 4; $\alpha=23$ 4
^{205}Fr	-1310	8						$\alpha\approx 100$; $\beta^+ < 1$
^{205}Ra	5840	90						$\alpha=?$; β^+ ?
$^{205}\text{Ra}^m$	6150#	100#	310#	110#				$\alpha=?$; IT ?
* ^{205}Ra	T : average 96Le09=210(+60-40) 87He10=220(60)							
^{206}Hg	-20946	20						$\beta^-=100$
^{206}Tl	-22253.1	1.4						$\beta^-=100$
$^{206}\text{Tl}^m$	-19610.0	1.4	2643.11	0.19				IT=100
^{206}Pb	-23785.4	1.2			STABLE			IS=24.1 1
$^{206}\text{Pb}^m$	-21585.3	1.2	2200.14	0.04				IT=100
$^{206}\text{Pb}^n$	-19758.1	1.4	4027.3	0.7				IT=100
^{206}Bi	-20028	8						$\beta^+=100$
$^{206}\text{Bi}^m$	-19968	8	59.897	0.017				IT=100
$^{206}\text{Bi}^n$	-18983	8	1044.8	0.5				IT=100
^{206}Po	-18182	8						$\beta^+=94.55$ 5; $\alpha=5.45$ 5
$^{206}\text{Po}^m$	-16596	8	1585.85	0.11				IT=100
$^{206}\text{Po}^n$	-15920	8	2262.22	0.14				IT=100
^{206}At	-12420	20						$\beta^+=99.11$ 8; $\alpha=0.89$ 8
$^{206}\text{At}^m$	-11613	20	807	3				IT=100
^{206}Rn	-9116	15						$\alpha=62$ 3; $\beta^+=38$ 3
^{206}Fr	-1243	28						$\beta^+=?$; $\alpha=42$ 24
$^{206}\text{Fr}^m$	-1048	28	190	40	MD			$\alpha=42$ 24; β^+ ?; IT ?
$^{206}\text{Fr}^n$	-517	28	730	40	MD			IT=?; $\alpha\approx 12$ #
^{206}Ra	3565	18						$\alpha=100$
^{206}Ac	13510	70			* &	25	ms	7
$^{206}\text{Ac}^m$	13590	90	80	50	* &	15	ms	6
$^{206}\text{Ac}^n$	13800#	80#	290#	110#	&	41	ms	16
* $^{206}\text{Po}^m$	E : less than 40 keV above 1573.4 level, from ENSDF							
* ^{206}Fr	D : $\alpha=84(2)$ % for mixture of ^{206}Fr and $^{206}\text{Fr}^m$, in 92Hu04. Value replaced by							
* ^{206}Fr	D : uniform distribution 0%-84% for each isomer							
* $^{206}\text{Fr}^n$	E : 531 keV above $^{206}\text{Fr}^m$, from ENSDF							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
^{207}Hg	-16220	150	2.9 m	0.2		(9/2 ⁺)	94 β^- =100
^{207}Tl	-21034	5	4.77 m	0.02		(1/2 ⁺)	94 β^- =100
$^{207}\text{Tl}^m$	-19686	5	1348.1	0.3		1.33 s	0.11 (11/2 ⁻) 94 IT≈100; $\beta^- < 0.1\%$
^{207}Pb	-22451.9	1.2				STABLE	94 IS=22.1 1
$^{207}\text{Pb}^m$	-20818.5	1.2	1633.368	0.005		806 ms	6 (13/2 ⁺) 94 IT=100
^{207}Bi	-20054.4	2.4				32.9 y	1.4 (9/2 ⁻) 94 β^+ =100
$^{207}\text{Bi}^m$	-17952.9	2.4	2101.49	0.16		182 μs	6 (21/2 ⁺) 94 IT=100
^{207}Po	-17146	7				5.80 h	0.02 (5/2 ⁻) 94 $\beta^+ \approx 100$; $\alpha=0.021$ 2
$^{207}\text{Po}^m$	-15763	7	1383.15	0.06		2.79 s	0.08 (19/2 ⁻) 94 IT=100
$^{207}\text{Po}^n$	-16031	7	1115.073	0.016		49 μs	(13/2 ⁺)
^{207}At	-13243	21				1.80 h	0.04 (9/2 ⁻) 94 $\beta^+=91.4$ 10; $\alpha=8.6$ 10
^{207}Rn	-8631	26				9.25 m	0.17 (5/2 ⁻) 94 $\beta^+=79$ 3; $\alpha=21$ 3
$^{207}\text{Rn}^m$	-7732	26	899.0	1.0		181 μs	18 (13/2 ⁺) 94 IT=100
^{207}Fr	-2840	50				14.8 s	0.1 (9/2 ⁻) 94 $\alpha=95$ 2; $\beta^+=5$ 2
^{207}Ra	3540	60				1.3 s	0.2 (5/2 ⁻ , 3/2 ⁻) 94 $\alpha \approx 90$; $\beta^+ \approx 10$
$^{207}\text{Ra}^m$	4095	25	560	50	AD	57 ms	8 (13/2 ⁺) 94 96Le09 T IT=85#; $\alpha=?$; ... *
^{207}Ac	11130	50				31 ms	8 (9/2 ⁻) 98 94Le05 TD $\alpha=100$ *
$^{207}\text{Ra}^m$	D : ... ; $\beta^+=0.55\%$ **						
$^{207}\text{Ra}^m$	T : average 96Le09=63(16) 87He10=55(10) **						
^{207}Ac	T : average 98Es02=27(+11-6) 94Le05=22(+40-9) **						
^{208}Hg	-13100#	300#				42 m	5 (0 ⁺) 98 98Zh22 T β^- =100 *
^{208}Tl	-16749.5	2.0				3.053 m	0.004 (5 ⁺) 98 β^- =100
^{208}Pb	-21748.5	1.2				STABLE	0 ⁺ 96 IS=52.4 1
$^{208}\text{Pb}^m$	-16853.5	2.3	4895	2		500 ns	10 (10 ⁺) 86 98Pf02 T IT=100
^{208}Bi	-18870.0	2.4				368 ky	4 (5 ⁺) 86 β^+ =100
$^{208}\text{Bi}^m$	-17298.9	2.4	1571.1	0.4		2.58 ms	0.04 (10 ⁻) 86 IT=100
^{208}Po	-17469.5	1.8				2.898 y	0.002 (0 ⁺) 86 $\alpha \approx 100$; $\beta^+=0.00223$ 23
^{208}At	-12491	26				1.63 h	0.03 (6 ⁺) 86 $\beta^+=99.45$ 6; $\alpha=0.55$ 6
^{208}Rn	-9648	11				24.35 m	0.14 (0 ⁺) 86 $\alpha=62$ 7; $\beta^+=38$ 7
^{208}Fr	-2670	50				59.1 s	0.3 (7 ⁺) 86 $\alpha=90$ 4; $\beta^+=10$ 4
^{208}Ra	1714	15				1.3 s	0.2 (0 ⁺) 86 $\alpha=?$; $\beta^+=5\%$
$^{208}\text{Ra}^m$	3510	200	1800	200		270 ns	(8 ⁺) 98Le.A ETJ
^{208}Ac	10760	60				97 ms	16 (3 ⁺) 96 96lk01 T $\alpha=?$; $\beta^+=1\%$ *
$^{208}\text{Ac}^m$	11258	28	500	50	AD	28 ms	7 (10 ⁻) 96 96lk01 T $\alpha=?$; IT<10#; $\beta^+=1\%$ *
^{208}Hg	T : 98Zh22=41(+5-4) supersedes 94Zh02=42(+23-12) of same group **						
^{208}Ac	T : average 96lk01=83(+34-19) 94Le05=95(+24-16) **						
$^{208}\text{Ac}^m$	E : if α decay goes to (7 ⁺) $^{204}\text{Fr}^m$, instead of (10 ⁻) as assumed in AME, then **						
$^{208}\text{Ac}^m$	E : E will become 234(22) keV **						
$^{208}\text{Ac}^m$	T : average 96lk01=21(+28-8) 94Le05=25(+9-5) **						
^{209}Hg	-8350#	200#				37 s	8 (9/2 ⁺)# 98Zh22 T β^- =100
^{209}Tl	-13638	8				2.161 m	0.007 (1/2 ⁺) 91 94Ar23 T β^- =100
^{209}Pb	-17614.4	1.8				3.253 h	0.014 (9/2 ⁺) 91 β^- =100
^{209}Bi	-18258.5	1.4				19 Ey	2 (9/2 ⁻) 91 03De11 TD IS=100.; $\alpha=100$
^{209}Po	-16365.9	1.8				102 y	5 (1/2 ⁻) 91 $\alpha \approx 100$; $\beta^+=0.48$ 4
^{209}At	-12880	7				5.41 h	0.05 (9/2 ⁻) 91 $\beta^+=95.9$ 5; $\alpha=4.1$ 5
^{209}Rn	-8929	20				28.5 m	1.0 (5/2 ⁻) 91 $\beta^+=83$ 2; $\alpha=17$ 2
$^{209}\text{Rn}^m$	-7755	20	1173.98	0.13		13.4 μs	(13/2 ⁺) 91
^{209}Fr	-3769	15				50.0 s	0.3 (9/2 ⁻) 91 $\alpha=89$ 3; $\beta^+=11$ 3
^{209}Ra	1850	50				4.6 s	0.2 (5/2 ⁻) 91 $\alpha \approx 90$; $\beta^+ \approx 10$
^{209}Ac	8840	50				92 ms	11 (9/2 ⁻) 91 00He17 T $\alpha=?$; $\beta^+=1\%$ *
^{209}Th	16500	100				7 ms	5 (5/2 ⁻)# 97 96lk01 TD $\alpha=?$; $\beta^+?$
^{209}Ac	T : average 00He17=98(+59-27) 96lk01=82(+18-13) 94Le05=91(+21-14) **						
^{209}Ac	T : and 68Va04=100(50) **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{210}Hg	-5110#	300#	10#	m (>300 ns)	0 ⁺	03 98Pf02 I	β^- ?	
^{210}Tl	-9246	12	1.30	m	0.03	5 ⁺ #	03 β^- =100; β^- n=0.009 6	
^{210}Pb	-14728.3	1.5	22.20	y	0.22	0 ⁺	03 β^- =100; α =1.9e-6 4	
$^{210}\text{Pb}^m$	-13450	5 1278	5	201	ns	17	8 ⁺ 03 IT=100	
^{210}Bi	-14791.8	1.4	5.012	d	0.005	1 ⁻	03 β^- =100; α =13.2e-5 10	
$^{210}\text{Bi}^m$	-14520.5	1.4 271.31	0.11	3.04	My	0.06	9 ⁻ 03 α =100	
$^{210}\text{Bi}^n$	-14358.3	1.4 433.49	0.10	57.5	ns	10	7 ⁻ 03 IT=100	
^{210}Po	-15953.1	1.2	138.376	d	0.002	0 ⁺	03 α =100	
$^{210}\text{Po}^m$	-14396.1	1.2 1556.96	0.03	98.9	ns	2.5	8 ⁺ 03 IT=100	
^{210}At	-11972	8	8.1	h	0.4	(5) ⁺	03 β^+ ≈100; α =0.175 20	
$^{210}\text{At}^m$	-9422	8 2549.6	0.2	482	μs	6	(15) ⁻ 03 IT=100	
$^{210}\text{At}^n$	-7944	8 4027.7	0.2	5.66	μs	0.07	(19) ⁺ 03 IT=100	
$^{210}\text{At}^p$	-5013	8 6959.3	0.6	98	ns	2	(26 ⁻) 03 IT=100	
^{210}Rn	-9598	9	2.4	h	0.1	0 ⁺	03 α =96 1; β^+ ?	
$^{210}\text{Rn}^m$	-7908	17 1690	15	644	ns	40	8 ⁺ # 03 IT ?	
$^{210}\text{Rn}^n$	-5761	17 3837	15	1.06	μs	0.05	(17) ⁻ 03 IT=100	
$^{210}\text{Rn}^p$	-3105	17 6493	15	1.04	μs	0.07	(22) ⁺ 03 IT=100	
^{210}Fr	-3346	22	3.18	m	0.06	6 ⁺	03 α =60 30; β^+ =40 30	
^{210}Ra	461	15	3.7	s	0.2	0 ⁺	03 α ?; β^+ =4#	
$^{210}\text{Ra}^m$	2260	200 1800	200	2.24	μs	(8 ⁺)	03 98Le.A EJ	
^{210}Ac	8790	60	350	ms	40	7 ⁺ #	03 00He17 T α ?; β^+ =9#	
^{210}Th	14043	25	17	ms	11	0 ⁺	03 α ?; β^+ =1#	
* $^{210}\text{Rn}^m$	E : ENSDF2003: less than 50 keV above 1664.6 level							**
* ^{210}Ac	T : average 00He17=335(+64-46) 68Va04=350(50)							**
^{211}Tl	-6080#	200#	1#	m (>300 ns)	1/2 ⁺ #	98Pf02 I	β^- ?	
^{211}Pb	-10491.4	2.7	36.1	m	0.2	9/2 ⁺	91 β^- =100	
^{211}Bi	-11858	6	2.14	m	0.02	9/2 ⁻	91 α ≈100; β^- =0.276 4	
$^{211}\text{Bi}^m$	-10631	6 1227.2	0.3	70	ns	5	(21/2 ⁻) 91 IT=100	
$^{211}\text{Bi}^n$	-10601	12 1257	10	1.4	μs	0.3	(25/2 ⁻) 91 98Pf02 T IT=100	
^{211}Po	-12432.5	1.3	516	ms	3	9/2 ⁺	91 α =100	
$^{211}\text{Po}^m$	-10970	5 1462	5	AD 25.2	s	0.6	(25/2 ⁺) 91 α ≈100; IT=0.016 4	
$^{211}\text{Po}^n$	-10298	5 2135	5	0.25	μs	0.07	(31/2 ⁻) 98Fo04 ETJ IT≈100; α ?	
$^{211}\text{Po}^p$	-7559	5 4874	5	2	μs	1	(43/2 ⁺) 98Fo04 ETJ IT≈100; α ?	
^{211}At	-11647.1	2.8	7.214	h	0.007	9/2 ⁻	96 ϵ =58.20 8; α =41.80 8	
^{211}Rn	-8756	7	14.6	h	0.2	1/2 ⁻	96 β^+ =72.6 17; α =27.4 17	
^{211}Fr	-4158	21	3.10	m	0.02	9/2 ⁻	91 α >80; β^+ <20	
^{211}Ra	836	26	13	s	2	5/2 ⁽⁻⁾	91 α >93; β^+ <7	
^{211}Ac	7200	70	213	ms	25	9/2 ⁻ #	91 00He17 T α ≈100; β^+ <0.2	
^{211}Th	13910	70	48	ms	20	5/2 ⁻ #	96 95Uu01 T α ?; β^+ =0.5#	
* ^{211}Ac	T : average 00He17=200(29) 68Va04=250(50)							**
^{212}Tl	-1650#	300#	30#	s (>300 ns)	5 ⁺ #	98Pf02 I	β^- ?	
^{212}Pb	-7547.4	2.2	10.64	h	0.01	0 ⁺	92 β^- =100	
$^{212}\text{Pb}^m$	-6212	10 1335	10	5	μs	1	(8 ⁺) 92 98Pf02 T IT=100	
^{212}Bi	-8117.3	2.0	60.55	m	0.06	1 ⁽⁻⁾	92 89Ha.A D β^- =64.06 6; α =35.94 6; ... *	
$^{212}\text{Bi}^m$	-7870	30 250	30	AD 25.0	m	0.2	(9 ⁻) 92 α =67 1; β^- =33 1; β^- α =30 1	
$^{212}\text{Bi}^n$	-5920#	200# 2200#	200#	7.0	m	0.3	> 15 92 β^- ≈100; IT ?	
^{212}Po	-10369.4	1.2	299	ns	2	0 ⁺	92 α =100	
$^{212}\text{Po}^m$	-7459	12 2911	12	AD 45.1	s	0.6	(18 ⁺) 92 α ≈100; IT=0.07 2	
^{212}At	-8621	7	314	ms	2	(1 ⁻)	92 α ≈100; β^+ <0.03; β^- <2e-6	
$^{212}\text{At}^m$	-8395	6 226	9	AD 119	ms	3	(9 ⁻) 92 α >99; IT<1	
$^{212}\text{At}^n$	-3849	8 4772	3	152	μs	5	(25 ⁻) 98By01 ETJ IT=100	
^{212}Rn	-8660	3	23.9	m	1.2	0 ⁺	92 α =100; 2 β^+ ?	
^{212}Fr	-3538	26	20.0	m	0.6	5 ⁺	92 β^+ =57 2; α =43 2	
^{212}Ra	-191	11	13.0	s	0.2	0 ⁺	92 α ?; β^+ =15#	
$^{212}\text{Ra}^m$	1767	11 1958.4	0.5	10.9	μs	0.4	(8 ⁺) 92 IT=100	

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
... A-group continued ...							
²¹² Ac	7280	70	920 ms	50	6 ⁺ #	92 00He17 T	$\alpha=?; \beta^+=3\#$
²¹² Th	12091	18	36 ms	15	0 ⁺	92	$\alpha\approx 100; \beta^+=0.3\#$
²¹² Pa	21610	70	8 ms	5	7 ⁺ #	97Mi03 TD	$\alpha=100$
* ²¹² Bi	D : ... ; $\beta^- \alpha=0.014$						**
* ²¹² Bi ⁿ	E : 1910 keV, if 100% β^- decay goes to 2922 level in ²¹² Po, and if $\log ft$ for						**
* ²¹² Bi ⁿ	E : this transition is 5.1 (see ENSDF), or higher						**
* ²¹² Ac	T : average 00He17=880(110) 68Va04=930(50)						**
* ²¹² Ac	J : ENSDF proposes to assign 7 ⁺ , if the observed α feeds the ²⁰⁸ Fr 7 ⁺ ground-state						**
²¹³ Pb	-3184	8	10.2 m	0.3	(9/2 ⁺)	92	$\beta^-=100$
²¹³ Bi	-5231	5	45.59 m	0.06	9/2 ⁻	92	$\beta^-=97.91\ 3; \alpha=2.09\ 3$
²¹³ Po	-6653	3	4.2 μ s	0.8	9/2 ⁺	92	$\alpha=100$
²¹³ At	-6579	5	125 ns	6	9/2 ⁻	92	$\alpha=100$
²¹³ Rn	-5698	6	19.5 ms	0.1	(9/2 ⁺)	92 00He17 T	$\alpha=100$
²¹³ Fr	-3550	8	34.6 s	0.3	9/2 ⁻	92	$\alpha=99.45\ 3; \beta^+=0.55\ 3$
²¹³ Ra	358	20	2.74 m	0.06	1/2 ⁻	92	$\alpha=80\ 5; \beta^+ ?$
²¹³ Ra ^m	2127	21	1769	6	AD	2.1 ms	0.1 17/2 ⁻ # 92 76Ra37 J
²¹³ Ac	6150	50	731 ms	17	9/2 ⁻ #	92 00He17 T	$\alpha=?; \beta^+ ?$
²¹³ Th	12120	70	140 ms	25	5/2 ⁻ #	92	$\alpha=?; \beta^+ ?$
²¹³ Pa	19660	70	7 ms	3	9/2 ⁻ #	97 95Ni05 TD	$\alpha=100$
* ²¹³ Rn	T : in same paper 18.0(0.4) 19.0(0.5), not used. Other 70Va13=25.0(0.2) at						**
* ²¹³ Rn	T : variance, not used						**
* ²¹³ Ra ^m	E : derived from difference in α decay energy in the AME evaluation.						**
* ²¹³ Ra ^m	E : ENSDF evaluation: less than 10 keV above 1769.7 level, thus 1775(3) keV						**
* ²¹³ Ra ^m	J : 17/2 ⁻ or 13/2 ⁺ as proposed by 76Ra37						**
²¹⁴ Pb	-181.3	2.4	26.8 m	0.9	0 ⁺	95	$\beta^-=100$
²¹⁴ Bi	-1200	11	19.9 m	0.4	1 ⁻	95 89Ha.A D	$\beta^-\approx 100; \alpha=0.021\ 1; \beta^- \alpha=0.003$
²¹⁴ Po	-4469.9	1.5	164.3 μ s	2.0	0 ⁺	95	$\alpha=100$
²¹⁴ At	-3380	4	558 ns	10	1 ⁻	95	$\alpha=100$
²¹⁴ At ^m	-3320	8	59	9	AD	268 ns	
²¹⁴ At ⁿ	-3146	5	234	6	AD	760 ns	
²¹⁴ Rn	-4320	9	270 ns	20	0 ⁺	95	$\alpha=100; 2\beta^+ ?$
²¹⁴ Rn ^m	-2695	9	1625.1	0.5		6.5 ns	3.0 8 ⁺
²¹⁴ Fr	-958	9	5.0 ms	0.2	(1 ⁻)	95	$\alpha=100$
²¹⁴ Fr ^m	-835	9	123	6	AD	3.35 ms	0.05 (8 ⁻) 95
²¹⁴ Ra	101	9	2.46 s	0.03	0 ⁺	95	$\alpha\approx 100; \beta^+=0.059\ 4$
²¹⁴ Ac	6429	22	8.2 s	0.2	5 ⁺ #	95	$\alpha\geq 89\ 3; \beta^+\leq 11\ 3$
²¹⁴ Th	10712	17	100 ms	25	0 ⁺	95	$\alpha\approx 100; \beta^+=0.1\ \#$
²¹⁴ Pa	19490	80	17 ms	3		95 95Ni05 D	$\alpha=100$
²¹⁵ Pb	4480#	410#	36 s	1	5/2 ⁺ #	96Ry.B T	$\beta^-=100$
²¹⁵ Bi	1649	15	7.6 m	0.2	(9/2 ⁻)	01	$\beta^-=100$
²¹⁵ Bi ^m	2997	15	1347.5	2.5		36.4 m	2.5 (25/2 ⁻) 01 02Fr.B D
²¹⁵ Po	-540.3	2.5	1.781 ms	0.004	9/2 ⁺	01	$\alpha=100; \beta^-=2.3e-4\ 2$
²¹⁵ At	-1255	7	100 μ s	20	9/2 ⁻	01	$\alpha=100$
²¹⁵ Rn	-1169	8	2.30 μ s	0.10	9/2 ⁺	01	$\alpha=100$
²¹⁵ Fr	318	7	86 ns	5	9/2 ⁻	01	$\alpha=100$
²¹⁵ Ra	2534	8	1.55 ms	0.07	9/2 ⁺ #	01	$\alpha=100$
²¹⁵ Ra ^m	4412	8	1877.8	0.5		7.1 μ s	0.2 (25/2 ⁺) 01
²¹⁵ Ra ⁿ	4781	8	2246.9	0.5		1.39 μ s	0.07 (29/2 ⁻) 01
²¹⁵ Ac	6012	21	170 ms	10	9/2 ⁻	01	$\alpha\approx 100; \beta^+=0.09\ 2$
²¹⁵ Th	10927	27	1.2 s	0.2	(1/2 ⁻)	01	$\alpha=100$
²¹⁵ Pa	17870	90	14 ms	2	9/2 ⁻ #	01	$\alpha=100$
* ²¹⁵ Pb	T : other preliminary result 02Fr.B=147(12) s						**
* ²¹⁵ Bi ^m	T : other preliminary result 02Fr.B=36.9(0.6) s						**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	J^π	Ens	Reference	Decay modes and intensities (%)			
^{216}Bi	5874	11		2.17 m	0.05	1 ⁻ #	97	96Ry.B T	$\beta^- = 100$	*	
^{216}Po	1783.8	2.2		145 ms	2	0 ⁺	97		$\alpha = 100; 2\beta^- ?$		
^{216}At	2257	4		300 μs	30	1 ⁽⁻⁾	97		$\alpha \approx 100; \beta^- < 0.006; \epsilon < 3e-7$		
$^{216}\text{At}^m$	2670	6	413	5	100#	μs	(9 ⁻)	97	$\alpha = 100$		
^{216}Rn	256	7		45 μs	5	0 ⁺	97		$\alpha = 100$		
^{216}Fr	2979	14		700 ns	20	(1 ⁻)	97		$\alpha = 100; \beta^+ < 2e-7\#$		
^{216}Ra	3291	9		182 ns	10	0 ⁺	97		$\alpha = 100; \epsilon < 1e-8$		
^{216}Ac	8123	27		440 μs	16	(1 ⁻)	97	00He17 T	$\alpha = 100; \beta^+ = 7e-5\#$		
$^{216}\text{Ac}^m$	8166	26	44	7	AD	443 μs	7	(9 ⁻)	97	00He17 T	$\alpha = 100; \beta^+ = 7e-5\#$
^{216}Th	10304	13		26.8 ms	0.3	0 ⁺	97	01Ha46 T	$\alpha \approx 100; \beta^+ = 0.006\#$	*	
$^{216}\text{Th}^m$	12346	16	2042	13	AD	137 μs	4	(8 ⁺)	97	01Ha46 TJD	IT=94.4; $\alpha = ?$
$^{216}\text{Th}^n$	12941	24	2637	20		615 ns	55	(11 ⁻)	97	01Ha46 TJ	IT=100
^{216}Pa	17800	70		105 ms	12		97	96An21 T	$\alpha = ?; \beta^+ = 2\#$	*	
* ^{216}Bi	T: also 90Ru02=3.6(0.4) outweighed, not used										**
* ^{216}Th	T: average 01Ha46=25.4(0.8) 00He17=27.0(0.3); other 68Va18=28(2) outweighed										**
* $^{216}\text{Th}^m$	T: average 01Ha46=128(8) 00He17=140(5)										**
* ^{216}Pa	T: not updated in 00He17: "could not be determined satisfactorily"										**
^{217}Bi	8820#	200#		97 s	3	9/2 ⁻ #		96Ry.B T	$\beta^- = 100$		
^{217}Po	5901	7		1.47 s	0.05	5/2 ⁺ #	91	96Ry.B T	$\alpha > 95; \beta^- < 5$		
^{217}At	4396	5		32.3 ms	0.4	9/2 ⁻	91	97Ch53 D	$\alpha \approx 100; \beta^- = 0.008$	*	
^{217}Rn	3659	4		540 μs	50	9/2 ⁺	91		$\alpha = 100$		
^{217}Fr	4315	7		16.8 μs	1.9	9/2 ⁻	94	90An19 T	$\alpha = 100$	*	
^{217}Ra	5887	9		1.63 μs	0.17	(9/2 ⁺)	91	90An19 T	$\alpha = 100$	*	
^{217}Ac	8707	13		69 ns	4	9/2 ⁻	91		$\alpha = ?; \beta^+ \leq 2$		
$^{217}\text{Ac}^m$	10719	19	2012	20	AD	740 ns	40	(29/2 ⁺)	91	IT=95.7 10; $\alpha = 4.3$	10
^{217}Th	12216	21		240 μs	5	(9/2 ⁺)	91	02He29 T	$\alpha = 100$	*	
^{217}Pa	17070	50		3.48 ms	0.09	9/2 ⁺ #	91	02He29 T	$\alpha = 100$	*	
$^{217}\text{Pa}^m$	18930	50	1860	7	AD	1.08 ms	0.03	29/2 ⁺ #	91	02He29 TD	$\alpha = 73$ 4; IT ?
^{217}U	22700	90		26 ms	14	1/2 ⁻ #		00Ma65 TD	$\alpha = ?$		
* ^{217}At	D: average β^- 97Ch53=0.0067(24) 69Le.A=0.012(4)										**
* ^{217}Fr	T: average 90An19=16(2) 70Bo13=22(5)										**
* ^{217}Ra	T: average 90An19=1.7(0.3) 70Bo13=1.6(0.2)										**
* ^{217}Th	T: average 02He29=237(2) 00He17=247(3) with Birge ratio $B=2.8$										**
* ^{217}Pa	T: average 02He29=3.8(0.2) 00He17=3.4(0.1)										**
^{218}Bi	13340#	360#		33 s	1	1 ⁻ #		02Fr.B TD	$\beta^- = 100$		
^{218}Po	8358.3	2.4		3.10 m	0.01	0 ⁺	96		$\alpha \approx 100; \beta^- = 0.020$	2	
^{218}At	8099	12		1.5 s	0.3	1 ⁻ #	96		$\alpha \approx 100; \beta^- = 0.1$		
^{218}Rn	5217.5	2.4		35 ms	5	0 ⁺	96		$\alpha = 100$		
^{218}Fr	7059	5		1.0 ms	0.6	1 ⁻	96		$\alpha = 100$		
$^{218}\text{Fr}^m$	7146	6	86	4	AD	22.0 ms	0.5		$\alpha \approx 100; \text{IT} ?$		
$^{218}\text{Fr}^p$	7260#	150#	200#	150#							
^{218}Ra	6651	11		25.6 μs	1.1	0 ⁺	96		$\alpha = 100; 2\beta^+ ?$		
^{218}Ac	10840	50		1.08 μs	0.09	1 ⁻ #	96		$\alpha = 100$		
$^{218}\text{Ac}^m$	10990#	70#	150#	50#		32 ns	9	(9 ⁻)	96	94De04 ET	*
$^{218}\text{Ac}^n$	11420#	70#	584#	50#		103 ns	11	(11 ⁺)	96		*
^{218}Th	12374	13		109 ns	13	0 ⁺	96		$\alpha = 100$		
^{218}Pa	18669	25		113 μs	10		96	00He17 T	$\alpha = 100$	*	
^{218}U	21920	30		6 ms	5	0 ⁺	96		$\alpha = 100$		
* $^{218}\text{Ac}^m$	E: at least 122.5 in 94De04										**
* $^{218}\text{Ac}^n$	E: 384.5(0.2) keV above $^{218}\text{Ac}^m$, from ENSDF										**
* ^{218}Pa	T: supersedes 96An21=110(20)										**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²¹⁹ Po	12800# 360#		2# m (>300 ns)	7/2 ⁺ #		98Pf02 I	$\beta^- ?; \alpha ?$
²¹⁹ At	10397 4		56 s 3	5/2 ⁻ #	01		$\alpha \approx 97; \beta^- \approx 3$
²¹⁹ Rn	8830.8 2.5		3.96 s 0.01	5/2 ⁺	01		$\alpha=100$
²¹⁹ Fr	8618 7		20 ms 2	9/2 ⁻	01		$\alpha=100$
²¹⁹ Ra	9394 8		10 ms 3	(7/2) ⁺	01		$\alpha=100$
²¹⁹ Ac	11570 50		11.8 μ s 1.5	9/2 ⁻	01		$\alpha=100; \beta^+=1e-6\#$
²¹⁹ Th	14470 50		1.05 μ s 0.03	9/2 ⁺ #	01		$\alpha=100; \beta^+=1e-7\#$
²¹⁹ Pa	18520 50		53 ns 10	9/2 ⁻	01		$\alpha=100; \beta^+=5e-9\#$
²¹⁹ U	23210 60		55 μ s 25	9/2 ⁺ #	01		$\alpha=100; \beta^+=1.4e-5\#$
²²⁰ Po	15470# 360#		40# s (>300 ns)	0 ⁺		98Pf02 I	$\beta^- ?$
²²⁰ At	14350 50		3.71 m 0.04	3(-#)	97		$\beta^-=92 2; \alpha=8 2$
²²⁰ Rn	10613.4 2.2		55.6 s 0.1	0 ⁺	97		$\alpha=100; 2\beta^- ?$
²²⁰ Fr	11483 4		27.4 s 0.3	1 ⁺	97		$\alpha \approx 100; \beta^- =0.35 5$
²²⁰ Ra	10273 9		17.9 ms 1.4	0 ⁺	97	00He17 T	$\alpha=100$ *
²²⁰ Ac	13752 15		26.36 ms 0.19	(3 ⁻)	97	90An19 T	$\alpha=100; \beta^+=5e-4\#$ *
²²⁰ Th	14669 22		9.7 μ s 0.6	0 ⁺	97		$\alpha=100; \epsilon=2e-7\#$
²²⁰ Pa	20380 60		780 ns 160	1 ⁻ #	97		$\alpha=100; \beta^+=3e-7\#$
²²⁰ U	23030# 200#		60# ns	0 ⁺			$\alpha ?; \beta^+ ?$
* ²²⁰ Ra	T : average 00He17=18(2) 90An19=17(2) 61Ru06=23(5)						**
* ²²⁰ Ac	T : average 90An19=26.4(0.2) 70Bo13=26.1(0.5)						**
²²¹ At	16810# 200#		2.3 m 0.2	3/2 ⁻ #	90		$\beta^-=100$
²²¹ Rn	14472 6		25 m 2	7/2(+)	90		$\beta^-=78 1; \alpha=22 1$
²²¹ Fr	13278 5		4.9 m 0.2	5/2 ⁻	90	97Ch53 D	$\alpha \approx 100; \beta^- =0.0048 15; \dots$ *
²²¹ Ra	12964 5		28 s 2	5/2 ⁺	90	94Bo28 D	$\alpha=100; {}^{14}\text{C}=1.2e-10 9$
²²¹ Ac	14520 50		52 ms 2	9/2 ⁻ #	90		$\alpha=100$
²²¹ Th	16938 9		1.68 ms 0.06	(7/2 ⁺)	90		$\alpha=100$ *
²²¹ Pa	20380 50		5.9 μ s 1.7	9/2 ⁻	90		$\alpha=100$
²²¹ U	24590# 100#		700# ns	9/2 ⁺ #			$\alpha ?; \beta^+ ?$
* ²²¹ Fr	D : ... ; ${}^{14}\text{C}=8.8e-11 11$						**
* ²²¹ Fr	D : β^- intensity is from 97Ch53; ${}^{14}\text{C}$ intensity is from 94Bo28						**
* ²²¹ Th	T : also 00He17=2.0(+0.3-0.2)						**
²²² At	20800# 300#		54 s 10		96		$\beta^-=100$
²²² Rn	16373.6 2.4		3.8235 d 0.0003	0 ⁺	96		$\alpha=100$
²²² Fr	16349 21		14.2 m 0.3	2 ⁻	96		$\beta^-=100$
²²² Ra	14321 5		38.0 s 0.5	0 ⁺	96		$\alpha=100; {}^{14}\text{C}=3.0e-8 10$
²²² Ac	16621 5		5.0 s 0.5	1 ⁻	96		$\alpha=99 1; \beta^+=1 1$
²²² Ac ^m	16820# 150# 200# 150# *		1.05 m 0.07	high	96		$\alpha=?; IT \leq 10; \beta^+=1.4 4$ *
²²² Th	17203 12		2.05 ms 0.07	0 ⁺	96	00He17 T	$\alpha=100; \epsilon < 1.3e-8\#$ *
²²² Pa	22120# 70#		3.2 ms 0.3		96	95Ni.A T	$\alpha=100$ *
²²² U	24300# 100#		1.4 μ s 0.7	0 ⁺	96		$\alpha=100; \beta^+ < 1e-6\#$
* ²²² Ac ^m	D : derived from $0.7\% < \beta^+ < 2\%$, in ENSDF						**
* ²²² Th	T : average 00He17=2.0(0.1) 99Gr28=2.1(0.1)						**
* ²²² Pa	T : average 95Ni.A=3.3(0.3) 79Sc09=2.9(+0.6-0.4)						**
* ²²² Pa	T : 70Bo13=5.7(0.5) at variance, not used						**
²²³ At	23460# 400#		50 s 7	3/2 ⁻ #	01		$\beta^- \approx 100; \alpha=0.008\#$
²²³ Rn	20300# 300#		24.3 m 0.4	7/2	01		$\beta^-=100; \alpha=0.0004\#$
²²³ Fr	18383.8 2.4		22.00 m 0.07	3/2(-)	01		$\beta^- \approx 100; \alpha=0.006$
²²³ Ra	17234.7 2.5		11.43 d 0.05	3/2 ⁺	01		$\alpha=100; {}^{14}\text{C}=8.9e-8 4$
²²³ Ac	17826 7		2.10 m 0.05	(5/2 ⁻)	01		$\alpha=99; \epsilon=1$
²²³ Th	19386 9		600 ms 20	(5/2) ⁺	01		$\alpha=100$
²²³ Pa	22320 70		5.1 ms 0.3	9/2 ⁻ #	01	99Ho28 T	$\alpha=100; \beta^+ < 0.001\#$ *
²²³ U	25840 70		21 μ s 8	7/2 ⁺ #	01		$\alpha \approx 100; \beta^+=0.2\#$
* ²²³ Pa	T : average 99Ho28=4.9(0.4) 95Ni.A=5.0(1.0) 70Bo13=6.5(1.0)						**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life		J^π	Ens	Reference	Decay modes and intensities (%)
²²⁴ Rn	22440# 300#		107	m	3	0 ⁺	97	$\beta^- = 100$
²²⁴ Fr	21660 50		3.33	m	0.10	1 ⁻	97	$\beta^- = 100$
²²⁴ Ra	18827.2 2.2		3.66	d	0.04	0 ⁺	97	$\alpha = 100$; ¹⁴ C=4.0e-9 12
²²⁴ Ac	20235 4		2.78	h	0.17	0 ⁻	97	$\beta^+ = 90.6$ 17; $\alpha = 9.4$ 17; $\beta^- < 1.6\#$
²²⁴ Th	19996 11		1.05	s	0.02	0 ⁺	97	$\alpha = 100$; $2\beta^+ ?$
²²⁴ Pa	23870 16		844	ms	19	5 ⁻ #	97	$\alpha \approx 100$; $\beta^+ = 0.1\#$
²²⁴ U	25714 25		940	μ s	270	0 ⁺	97	$\alpha = 100$; $\beta^+ < 1.2e-4\#$
* ²²⁴ Pa	T : average ⁹⁶ Li05=790(60) ⁹⁶ Wi.A=850(20)							
* ²²⁴ U	T : average ⁹² To02=1000(400) ⁹¹ An10=700(+500-200)							
²²⁵ Rn	26490# 300#		4.66	m	0.04	7/2 ⁻	90	$\beta^- = 100$
²²⁵ Fr	23810 30		4.0	m	0.2	3/2 ⁻	90	$\beta^- = 100$
²²⁵ Ra	21994.0 3.0		14.9	d	0.2	1/2 ⁺	90	$\beta^- = 100$
²²⁵ Ac	21638 5		10.0	d	0.1	(3/2 ⁻)	90	$\alpha = 100$; ¹⁴ C=6.0e-10 13
²²⁵ Th	22310 5		8.72	m	0.04	(3/2 ⁺)	90	$\alpha \approx 90$; $\epsilon \approx 10$
²²⁵ Pa	24340 70		1.7	s	0.2	5/2 ⁻ #	90	$\alpha = 100$
²²⁵ U	27377 12		61	ms	4	5/2 ⁺ #	90	$\alpha = 100$
²²⁵ Np	31590 70		3#	ms	(>2 μ s)	9/2 ⁻ #	97	$\alpha = 100$
* ²²⁵ U	T : ⁰⁰ He17=59(+5-2); others ⁹⁴ An02=68(+45-20) ⁹² To02=95(15) and							
* ²²⁵ U	T : ⁸⁹ He13=80(+40-10) outweighed, not used							
²²⁶ Rn	28770# 400#		7.4	m	0.1	0 ⁺	96	$\beta^- = 100$
²²⁶ Fr	27370 100		49	s	1	1 ⁻	96	$\beta^- = 100$
²²⁶ Ra	23669.1 2.3		1.600	ky	0.007	0 ⁺	96	$\alpha = 100$; ¹⁴ C=2.6e-9 6; $2\beta^- ?$
²²⁶ Ac	24310 3		29.37	h	0.12	(1) ^(-#)	96	$\beta^- = 83$ 3; $\epsilon = 17$ 3; $\alpha = 0.006$ 2
²²⁶ Th	23197 5		30.57	m	0.10	0 ⁺	96	$\alpha = 100$; ¹⁸ O<3.2e-12
²²⁶ Pa	26033 11		1.8	m	0.2		96	$\alpha = 74$ 5; $\beta^+ = 26$ 5
²²⁶ U	27329 13		269	ms	6	0 ⁺	96	$\alpha = 100$
²²⁶ Np	32740# 90#		35	ms	10		96	$\alpha = 100$; $\beta^+ = 0.003\#$
* ²²⁶ Ra	D : ¹⁴ C: average ⁹⁰ We01=2.3(0.8) ⁸⁶ Ba26=2.9(1.0) ⁸⁵ Ho21=3.2(1.6)							
* ²²⁶ U	T : average ⁰¹ Ca.B=258(13) ⁰⁰ He17=281(9) ⁹⁹ Gr28=260(10)							
²²⁷ Rn	32980# 420#		20.8	s	0.7	5/2 ^(+#)	01	$\beta^- = 100$
²²⁷ Fr	29650 100		2.47	m	0.03	1/2 ⁺	01	$\beta^- = 100$
²²⁷ Ra	27179.0 2.4		42.2	m	0.5	3/2 ⁺	01	$\beta^- = 100$
²²⁷ Ac	25850.9 2.4		21.772	y	0.003	3/2 ⁻	01	$\beta^- = 98.62$ 36; $\alpha = 1.38$ 36
²²⁷ Th	25806.2 2.5		18.68	d	0.09	1/2 ⁺	01	$\alpha = 100$
²²⁷ Pa	26832 7		38.3	m	0.3	(5/2 ⁻)	01	$\alpha = 85$ 2; $\epsilon = 15$ 2
²²⁷ U	29022 17		1.1	m	0.1	(3/2 ⁺)	01	$\alpha = 100$; $\beta^+ < 0.001\#$
²²⁷ Np	32560 70		510	ms	60	5/2 ⁻ #	01	$\alpha \approx 100$; $\beta^+ = 0.05\#$
²²⁸ Rn	35380# 410#		65	s	2	0 ⁺	97	$\beta^- = 100$
²²⁸ Fr	33280# 200#		38	s	1	2 ⁻	97	$\beta^- = 100$
²²⁸ Ra	28941.8 2.4		5.75	y	0.03	0 ⁺	97	$\beta^- = 100$
²²⁸ Ac	28896.0 2.5		6.15	h	0.02	3 ⁺	97	$\beta^- = 100$
²²⁸ Th	26772.2 2.2		1.9116	y	0.0016	0 ⁺	97	$\alpha = 100$; ²⁰ O=1.13e-11 22
²²⁸ Pa	28924 4		22	h	1	3 ⁺	97	$\beta^+ = 98.0$ 2; $\alpha = 2.0$ 2
²²⁸ U	29225 15		9.1	m	0.2	0 ⁺	97	$\alpha > 95$; $\epsilon < 5$
²²⁸ Np	33700# 200#		61.4	s	1.4		97	$\epsilon = 60$ 7; $\alpha = 40$ 7; $\beta^+ \text{SF} = 0.012$ 6
²²⁸ Pu	36090 30		10#	ms	(>2 μ s)	0 ⁺	97	$\alpha \approx 100$; $\beta^+ = 0.1\#$
* ²²⁸ Np	D : $\beta^+ \text{SF} = 0.020(9)\%$ defined by ⁹⁴ Kr13 relative to ϵ , thus 0.012(6)% of total							

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²²⁹ Fr	35820	40	50.2 s	0.4	1/2 ⁺ #	90	92Bo05 T β^- =100
²²⁹ Ra	32563	19	4.0 m	0.2	5/2 ⁽⁺⁾	90	β^- =100
²²⁹ Ac	30750	30	62.7 m	0.5	(3/2 ⁺)	90	β^- =100
²²⁹ Th	29586.5	2.8	7.34 ky	0.16	5/2 ⁺	90	α =100
²²⁹ Th ^m	29586.5	2.8	0.0035	0.0010	70 h	50	3/2 ⁺ 94He08 TEJ IT ? *
²²⁹ Pa	29898.0	2.7	1.50 d	0.05	(5/2 ⁺)	90	ϵ ≈100; α =0.48 5
²²⁹ Pa ^m	29909.6	2.7	11.6	0.3	420 ns	30	3/2 ⁻ 98Le15 EJD IT=100
²²⁹ U	31211	6	58 m	3	(3/2 ⁺)	90	β^+ ≈80; α ≈20
²²⁹ Np	33780	90	4.0 m	0.2	5/2 ⁺ #	90	α >50; β^+ <50
²²⁹ Np ^p	33850#	100#	70#	50#	5/2 ⁻ #		
²²⁹ Pu	37400	50	120 s	50	3/2 ⁺ #	97	01Ca.B TD α =100
* ²²⁹ Th ^m	D : ultraviolet γ -ray emission assigned by 97Ir02 and 98Ri03 to IT decay is **						
* ²²⁹ Th ^m	D : proved by 99Sh12 to be due to N ₂ discharge emission. 99Ut01 sees **						
* ²²⁹ Th ^m	D : no UV in vacuo. **						
²³⁰ Fr	39600#	450#	19.1 s	0.5		93	β^- =100
²³⁰ Ra	34518	12	93 m	2	0 ⁺	93	β^- =100
²³⁰ Ac	33810	300	122 s	3	(1 ⁺)	94	01Yu03 D β^- =100; β^- SF=1.19e-6 40
²³⁰ Th	30864.0	1.8	75.38 ky	0.30	0 ⁺	93	α =100; SF<5e-11; ... *
²³⁰ Pa	32175	3	17.4 d	0.5	(2 ⁻)	93	β^+ ≈91.6 13; β^- ≈8.4 13; ... *
²³⁰ U	31615	5	20.8 d		0 ⁺	93	01Bo11 D α =100; 22Ne=4.8e-12 20; ... *
²³⁰ Np	35240	50	4.6 m	0.3		93	β^+ <97; α ≥3
²³⁰ Np ^p	35540#	210#	300#	200#	am		
²³⁰ Pu	36934	15	1.70 m	0.17	0 ⁺	93	01Ca.B T α =?; β^+ ? *
* ²³⁰ Th	D : ... ; ²⁴ Ne=5.6e-11 10 **						
* ²³⁰ Pa	D : ... ; α =0.0032 1 **						
* ²³⁰ U	D : ... ; SF<1.4e-10#; 2 β^+ ? **						
* ²³⁰ Pu	T : also 90An22=154(66)s outweighed, not used **						
²³¹ Fr	42330#	470#	17.6 s	0.6	1/2 ⁺ #	01	β^- =100
²³¹ Ra	38400#	300#	103 s	3	(5/2 ⁺)	01	β^- =100
²³¹ Ra ^m	38470#	300#	66.21	0.09	53 μ s	(1/2 ⁺)	01 IT=100
²³¹ Ac	35920	100	7.5 m	0.1	(1/2 ⁺)	01	β^- =100
²³¹ Th	33817.3	1.8	25.52 h	0.01	5/2 ⁺	01	β^- =100; α =4e-11#
²³¹ Pa	33425.7	2.3	32.76 ky	0.11	3/2 ⁻	01	α =100; SF≤3e-10; ... *
²³¹ U	33807	3	4.2 d	0.1	(5/2 ⁽⁺⁾)	01	ϵ ≈100; α =0.004 1
²³¹ Np	35630	50	48.8 m	0.2	(5/2 ⁽⁺⁾)	01	β^+ ≈98 1; α =2 1
²³¹ Np ^p	35690#	60#	40#		5/2 ⁻ #		
²³¹ Pu	38285	26	8.6 m	0.5	3/2 ⁺ #	01	99La14 D β^+ ≈87 5; α =13 5
²³¹ Am	42440#	300#	30#	s			β^+ ?; α ?
* ²³¹ Pa	D : ... ; ²⁴ Ne=13.4e-10 17; ²³ F=9.9e-13 **						
²³² Fr	46360#	640#	5 s	1		97	90Me13 T β^- =100
²³² Ra	40650#	280#	250 s	50	0 ⁺	91	β^- =100
²³² Ac	39150	100	119 s	5	(1 ⁺)	91	β^- =100
²³² Th	35448.3	2.0	14.05 Gy	0.06	0 ⁺	91	95Bo18 D IS=100.; α =100; SF=11e-10 3; ... *
²³² Pa	35948	8	1.31 d	0.02	(2 ⁻)	91	β^- ≈100; ϵ =0.003 1
²³² U	34610.7	2.2	68.9 y	0.4	0 ⁺	91	90Bo16 D α =100; ²⁴ Ne=8.9e-10 7; ... *
²³² Np	37360#	100#	14.7 m	0.3	(4 ⁺)	91	β^+ ≈100; α ≈0.003
²³² Pu	38366	18	33.7 m	0.5	0 ⁺	91	ABBW D ϵ =?; α =11# *
²³² Am	43400#	300#	1.31 m	0.04		91	β^+ =?; α =2#; β^+ SF=0.069 10
* ²³² Th	D : ... ; ²⁴ Ne+ ²⁶ Ne<2.78e-10; 2 β^- ? **						
* ²³² U	D : ... ; ²⁸ Mg<5e-12; SF<1e-12 **						
* ²³² U	D : ²⁴ Ne: average, as adopted by 91Bo20, of 2 results from their group **						
* ²³² Pu	T : average 00La25=33.1(0.8) 73Ja06=34.1(0.7) **						
* ²³² Pu	D : derived from 1.6%# < α < 20%#, in ENSDF **						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²³³ Ra	44770#	470#	30 s	5	1/2 ⁺ #	97	β^- =100
²³³ Ac	41500#	300#	145 s	10	(1/2 ⁺)	90	β^- =100
²³³ Th	38733.2	2.0	22.3 m	0.1	1/2 ⁺	90	β^- =100
²³³ Pa	37490.1	2.2	26.967 d	0.002	3/2 ⁻	90	β^- =100
²³³ U	36920.0	2.7	159.2 ky	0.2	5/2 ⁺	96	α =100; SF<6e-9; ...
²³³ Np	37950	50	36.2 m	0.1	(5/2 ⁺)	90	β^+ ≈100; α <0.001
²³³ Np ^p	38000#	60#	50#	30#	(5/2 ⁻)	90	
²³³ Pu	40050	50	20.9 m	0.4	5/2 ⁺ #	90	β^+ ≈100; α =0.12 5
²³³ Am	43170#	100#	3.2 m	0.8		00Sa52	TD β^+ ?; α >3
²³³ Cm	47290	70	1#	m	3/2 ⁺ #	01Ca.B	D α =?; β^+ ?
* ²³³ U	D : . . . ; ²⁴ Ne=7.2e-11 9; ²⁸ Mg<1.3e-13						
²³⁴ Ra	47230#	490#	30 s	10	0 ⁺	94	β^- =100
²³⁴ Ac	45100#	400#	44 s	7		94	β^- =100
²³⁴ Th	40614	3	24.10 d	0.03	0 ⁺	94	β^- =100
²³⁴ Pa	40341	5	6.70 h	0.05	4 ⁺	94	78Ga07 D β^- =100; SF<3e-10
²³⁴ Pa ^m	40419	4	78	3	1.17 m	0.03	(0 ⁻) 94 78Ga07 D β^- ≈100; IT=0.16 4; SF<1e-10
²³⁴ U	38146.6	1.8	245.5 ky	0.6	0 ⁺	94	IS=0.0055 2; α =100; ...
²³⁴ U ^m	39567.9	1.8	1421.32	0.10	33.5 μ s	2.0	6 ⁻
²³⁴ Np	39956	9	4.4 d	0.1	(0 ⁺)	94	β^+ =100
²³⁴ Pu	40350	7	8.8 h	0.1	0 ⁺	94	ϵ ≈94; α ≈6
²³⁴ Am	44530#	210#	2.32 m	0.08		94	90Ha02 D β^+ ≈100; α =0.039 12; ...
²³⁴ Cm	46724	18	51 s	12	0 ⁺	01Ca.B	TD α =?; β^+ =47#; SF=3
* ²³⁴ U	D : . . . ; SF=1.73e-9 10; ²⁸ Mg=1.4e-11 3; ²⁴ Ne+ ²⁶ Ne=9e-12 7						
* ²³⁴ Am	D : . . . ; β^+ SF=0.0066 18						
²³⁵ Ac	47720#	360#	40#	s	1/2 ⁺ #		β^- ?
²³⁵ Th	44260	50	7.2 m	0.1	1/2 ⁺ #	03	β^- =100
²³⁵ Pa	42330	50	24.44 m	0.11	(3/2 ⁻)	03	β^- =100
²³⁵ U	40920.5	1.8	704 My	1	7/2 ⁻	03	IS=0.7200 51; α =100; ...
²³⁵ U ^m	40920.6	1.8	0.0765	0.0004	26 m		1/2 ⁺ 03 IT=100
²³⁵ Np	41044.7	2.0	396.1 d	1.2	5/2 ⁺	03	ϵ ≈100; α =0.00260 13
²³⁵ Pu	42184	21	25.3 m	0.5	(5/2 ⁺)	03	β^+ ≈100; α =0.0028 7
²³⁵ Am	44660#	120#	9.9 m	0.5	5/2 ⁻ #	03	β^+ ≈100; α =0.40 5
²³⁵ Cm	47910#	200#	5#	m	5/2 ⁺ #	03	β^+ ?; α ?
²³⁵ Cm ^p	47960#	210#	50#	50#			<i>am</i>
²³⁵ Bk	52700#	400#	20#	s			β^+ ?; α ?
* ²³⁵ U	D : . . . ; SF=7e-9 2; ²⁰ Ne=8e-10 4; ²⁵ Ne≈8e-10; ²⁸ Mg=8e-10						
²³⁶ Ac	51510#	500#	2#	m			β^- ?
²³⁶ Th	46450#	200#	37.5 m	0.2	0 ⁺	91	β^- =100
²³⁶ Pa	45350	200	9.1 m	0.1	1 ⁽⁻⁾	91	β^- =100; β^- SF=6e-8 4
²³⁶ U	42446.3	1.8	23.42 My	0.03	0 ⁺	91	α =100; SF=9.6e-8 6
²³⁶ U ^m	45196	10	2750	10	115 ns		0 ⁺
²³⁶ Np	43380	50	* 154 ky	6	(6 ⁻)	91	ϵ =87.3 5; β^- =12.5 5; α =0.16 4
²³⁶ Np ^m	43439	7	* 22.5 h	0.4	1	91	ϵ =52 1; β^- =48 1
²³⁶ Np ^p	43618	14	240	50			AD 3 ⁻
²³⁶ Pu	42902.7	2.2	2.858 y	0.008	0 ⁺	91	90Og01 D α =100; SF=1.36e-7 4; ...
²³⁶ Am	46180#	100#	30#	m		91	β^+ ?; α ?
²³⁶ Cm	47890#	200#	10#	m		91	β^+ ?; α ?
²³⁶ Bk	53400#	400#	1#	m			β^+ ?; α ?
* ²³⁶ Pa	D : β^- SF decay questioned by 90Ha02						
* ²³⁶ U	D : and Ne+Mg < 4e-10%, from 89Mi.A						
* ²³⁶ Pu	D : . . . ; ²⁸ Mg=2e-12; 2 β^+ ?						

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
²³⁷ Th	50200# 360#		4.8 m	0.5	5/2 ⁺ #	97 00Xu02 T	$\beta^- = 100$	*
²³⁷ Pa	47640 100		8.7 m	0.2	(1/2 ⁺)	95	$\beta^- = 100$	
²³⁷ U	45391.9 1.9		6.75 d	0.01	1/2 ⁺	95	$\beta^- = 100$	
²³⁷ Np	44873.3 1.8		2.144 My	0.007	5/2 ⁺	95 89Pr.A D	$\alpha = 100$; SF $\leq 2e-10$; ³⁰ Mg $< 4e-12$	*
²³⁷ Pu	45093.3 2.2		45.2 d	0.1	7/2 ⁻	95	$\epsilon \approx 100$; $\alpha = 0.0042$	
²³⁷ Pu ^m	45238.8 2.2	145.544 0.010	180 ms	20	1/2 ⁺	95	IT=100	
²³⁷ Am	46570# 60#		73.0 m	1.0	5/2 ⁽⁻⁾	95	$\beta^+ \approx 100$; $\alpha = 0.025$	3
²³⁷ Cm	49280# 210#		20# m		5/2 ⁺ #	95	$\beta^+ ?$; $\alpha ?$	
²³⁷ Cm ^p	49480# 260#	200# 150#			7/2 ⁻			
²³⁷ Bk	53100# 220#		1# m		7/2 ⁺ #		$\beta^+ ?$; $\alpha ?$	
²³⁷ Bk ^p	53170# 230#	70# 30#			(3/2 ⁻)			
²³⁷ Cf	57820# 500#		2.1 s	0.3	5/2 ⁺ #	98 95La09 TD	$\alpha ?$; SF ≈ 10 ; $\beta^+ ?$	
* ²³⁷ Th	T : average 00Xu02=4.69(0.60) 93Yu03=5.0(0.9)							**
* ²³⁷ Np	D : and cluster (Z=10-14) < 1.8e-12%, from 92Mo03							**
²³⁸ Th	52630# 280#		9.4 m	2.0	0 ⁺	02	$\beta^- = 100$	
²³⁸ Pa	50770 60		2.27 m	0.09	3 ⁻ #	02 85Ba57 D	$\beta^- = 100$; β^- SF $< 2.6e-6$	
²³⁸ U	47308.9 1.9		4.468 Gy	0.003	0 ⁺	02 91Tu02 D	IS=99.2745 106; $\alpha = 100$; ...	*
²³⁸ U ^m	49866.8 2.0	2557.9 0.5	280 ns	6	0 ⁺	02	IT=?; SF=2.6 4; $\alpha < 0.5$	
²³⁸ Np	47456.3 1.8		2.117 d	0.002	2 ⁺	02	$\beta^- = 100$	
²³⁸ Np ^m	49760# 200#	2300# 200#	112 ns	39		02	SF ≈ 100 ; IT ?	
²³⁸ Pu	46164.7 1.8		87.7 y	0.1	0 ⁺	02 89Wa10 D	$\alpha = 100$; SF=1.9e-7 1; ...	*
²³⁸ Am	48420 50		98 m	2	1 ⁺	02	$\beta^+ = 100$; $\alpha = 1.0e-4$	4
²³⁸ Am ^m	50920# 210#	2500# 200#	35 μ s	10		02	SF ≈ 100 ; IT ?	
²³⁸ Cm	49400 40		2.4 h	0.1	0 ⁺	02	$\epsilon ?$; $\alpha \leq 10$	
²³⁸ Bk	54290# 290#		2.40 m	0.08		02 94Kr03 D	$\beta^+ \approx 100$; $\alpha ?$; β^+ SF=0.048 2	
²³⁸ Bk ^p	54490# 330#	200# 150#			<i>am</i>			
²³⁸ Cf	57200# 400#		21.1 ms	1.3	0 ⁺	02 01Og08 TD	SF ≈ 100 ; $\alpha \approx 0.2$; $\beta^+ ?$	*
* ²³⁸ U	D : ... ; SF=5.45e-5 7; 2 $\beta^- = 2.2e-10$ 7							**
* ²³⁸ U	D : 2 $\beta^- = 2.2(7)e-10$ % derived from 2 β^- half-life T=2.0(0.6) Zy, in 91Tu02							**
* ²³⁸ Pu	D : ... ; ³² Si $\approx 1.4e-14$; ²⁸ Mg+ ³⁰ Mg $\approx 6e-15$							**
* ²³⁸ Cf	T : average 01Og08=21.1(+1.9-1.7) 95La09=21(2)							**
²³⁹ Pa	53340# 200#		1.8 h	0.5	(3/2 ⁻) ^(-#)	03	$\beta^- = 100$	
²³⁹ U	50573.9 1.9		23.45 m	0.02	5/2 ⁺	03	$\beta^- = 100$	
²³⁹ U ^m	50594# 20#	20# 20#	> 250 ns		(5/2 ⁺)	03	$\beta^- = 100$	
²³⁹ U ⁿ	50707.7 1.9	133.7990 0.0010	780 ns	40	1/2 ⁺	03	IT=100	
²³⁹ Np	49312.4 2.1		2.356 d	0.003	5/2 ⁺	03	$\beta^- = 100$; $\alpha = 5e-10$ #	
²³⁹ Pu	48589.9 1.8		24.11 ky	0.03	1/2 ⁺	03	$\alpha = 100$; SF=3.1e-10 6	
²³⁹ Pu ^m	48981.5 1.8	391.584 0.003	193 ns	4	7/2 ⁻	03	IT=100	
²³⁹ Am	49392.0 2.4		11.9 h	0.1	(5/2 ⁻)	03	$\epsilon \approx 100$; $\alpha = 0.010$ 1	
²³⁹ Am ^m	51890 200	2500 200	163 ns	12	(7/2 ⁺)	03	SF ≈ 100 ; IT ?	
²³⁹ Cm	51190# 100#		2.9 h		(7/2 ⁻)	03	$\beta^+ \approx 100$; $\alpha < 0.1$	
²³⁹ Cm ^p	51340# 140#	150# 100#			1/2 ⁺			
²³⁹ Bk	54290# 230#		3# m		7/2 ⁺ #	03	$\beta^+ ?$; $\alpha ?$	
²³⁹ Bk ^p	54330# 230#	41 11			(3/2 ⁻)			
²³⁹ Cf	58150# 210#		60 s	30	5/2 ⁺ #	03	$\alpha = ?$; $\beta^+ ?$	
²⁴⁰ Pa	56800# 300#		2# m				$\beta^- ?$	
²⁴⁰ U	52715 5		14.1 h	0.1	0 ⁺	96	$\beta^- = 100$; $\alpha < 1e-10$	
²⁴⁰ Np	52315 15		* 61.9 m	0.2	(5 ⁺)	96	$\beta^- = 100$	
²⁴⁰ Np ^m	52335 21	20 15	* 7.22 m	0.02	1 ⁽⁺⁾	96 81Hs02 E	$\beta^- \approx 100$; IT=0.11 3	
²⁴⁰ Pu	50127.0 1.8		6.564 ky	0.011	0 ⁺	01 89Pr.A D	$\alpha = 100$; SF=5.7e-6 2; ³⁴ Si $< 1.3e-13$	
²⁴⁰ Am	51512 14		50.8 h	0.3	(3 ⁻)	96	$\beta^+ = 100$; $\alpha \approx 1.9e-4$	
²⁴⁰ Cm	51725.4 2.3		27 d	1	0 ⁺	96	$\alpha \approx 100$; $\epsilon < 0.5$; SF=3.9e-6 8	
²⁴⁰ Bk	55670# 150#		4.8 m	0.8		96	$\beta^+ ?$; $\alpha = 10$ #; β^+ SF=0.0020 13	
²⁴⁰ Bk ^p	55910# 180#	240# 100#			<i>am</i>			
²⁴⁰ Cf	58030# 200#		1.06 m	0.15	0 ⁺	96 95La09 D	$\alpha \approx 98$; SF ≈ 2 ; $\beta^+ ?$	
²⁴⁰ Es	64200# 400#		1# s				$\alpha ?$; $\beta^+ ?$	

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²⁴¹ U	56200#	300#			5# m	7/2 ⁺ #			β^- ?
²⁴¹ Np	54260	70			13.9 m	0.2 (5/2 ⁺)	94		β^- =100
²⁴¹ Pu	52956.8	1.8			14.35 y	0.10 5/2 ⁺	96		$\beta^- \approx 100$; $\alpha=0.00245$ 2; ... *
²⁴¹ Pu ^m	53118.4	1.8	161.60	0.10	880 ns	1/2 ⁺			
²⁴¹ Pu ⁿ	55160	200	2200	200	21 μ s	3			
²⁴¹ Am	52936.0	1.8			432.2 y	0.7 5/2 ⁻	94		$\alpha=100$; SF=4.3e-10 18; ... *
²⁴¹ Am ^m	55140	100	2200	100	1.5 μ s				
²⁴¹ Cm	53703.4	2.2			32.8 d	0.2 1/2 ⁺	94		$\epsilon=99.0$ 1; $\alpha=1.0$ 1
²⁴¹ Bk	56100#	200#			4.6 m	0.4 (7/2 ⁺)	94	03As01 T	α ?; β^+ ?
²⁴¹ Bk ^p	56150#	200#	51	3	AD	3/2 ⁻			
²⁴¹ Cf	59360#	260#			3.8 m	0.7 7/2 ⁻ #	94		$\beta^+ \approx 75$; $\alpha \approx 25$
²⁴¹ Cf ^p	59510#	270#	150#	100#	Nm	(1/2 ⁺)			
²⁴¹ Es	63840#	230#			10 s	5 (3/2 ⁻)	97	96Ni09 TJD	α =?; β^+ ?
²⁴¹ Es ^p	64240#	300#	400#	200#		(7/2 ⁺)			
* ²⁴¹ Pu	D : ... ; SF<2.4e-14 **								
* ²⁴¹ Am	D : ... ; ²⁴¹ Si<7.4e-14 **								
²⁴² U	58620#	200#			16.8 m	0.5 0 ⁺	02		β^- =100
²⁴² Np	57420	200			2.2 m	0.2 (1 ⁺)	02		β^- =100
²⁴² Np ^m	57420#	210#	0#	50#	* 5.5 m	0.1 6 ⁺ #	02		β^- =100
²⁴² Pu	54718.4	1.9			375 ky	2 0 ⁺	02		$\alpha=100$; SF=5.50e-4 6
²⁴² Am	55469.7	1.8			16.02 h	0.02 1 ⁻	02		β^- =82.7 3; $\epsilon=17.3$ 3
²⁴² Am ^m	55518.3	1.8	48.60	0.05	141 y	2 5 ⁻	02		IT \approx 100; $\alpha=0.45$ 2; SF<4.7e-9
²⁴² Am ⁿ	57670	80	2200	80	14.0 ms	1.0 (2 ⁺ , 3 ⁻)	02		SF \approx 100; IT=?; α ?
²⁴² Cm	54805.2	1.8			162.8 d	0.2 0 ⁺	02		$\alpha=100$; SF=6.2e-6 3; ... *
²⁴² Bk	57740#	200#			7.0 m	1.3 2 ⁻ #	02	80Ga07 D	$\beta^+ \approx 100$; β^+ SF<3e-5; α ?
²⁴² Bk ^m	57940#	280#	200#	200#	600 ns	100	02		SF \approx 100; IT ?
²⁴² Bk ^p	57990#	220#	250#	100#		4 ⁻			
²⁴² Cf	59340	40			3.49 m	0.15 0 ⁺	02	70Si19 T	$\alpha=80$ 20; β^+ ?; SF<0.014 *
²⁴² Es	64970#	330#			13.5 s	2.5 0 ⁺	02	94Ke.B D	α =?; β^+ =?; β^+ SF=0.6 *
²⁴² Fm	68400#	400#			800 μ s	200 0 ⁺	02		SF=?; α ?
* ²⁴² Cm	D : ... ; ²⁴² Si=1.1e-14 4; 2 β^+ ? **								
* ²⁴² Cf	T : average 70Si19=3.68(0.44) 67Si07=3.4(0.2) 67Fi04=3.2(0.5) 67Hl01=3.7(0.3) **								
* ²⁴² Es	D : β^+ SF=0.6% assuming α and β^+ are equal **								
²⁴³ Np	59880#	30#			1.85 m	0.15 (5/2 ⁻)	93		β^- =100
²⁴³ Np ^p	59925	11	50#	30#	Nm	(5/2 ⁻)			
²⁴³ Pu	57756	3			4.956 h	0.003 7/2 ⁺	93		β^- =100
²⁴³ Pu ^m	58140	3	383.6	0.4	330 ns	30 (1/2 ⁺)	93		IT=100
²⁴³ Am	57176.1	2.3			7.37 ky	0.04 5/2 ⁻	93		$\alpha=100$; SF=3.7e-9 2
²⁴³ Cm	57183.6	2.1			29.1 y	0.1 5/2 ⁺	93		$\alpha \approx 100$; $\epsilon=0.29$ 3; SF=5.3e-9 9
²⁴³ Cm ^p	57312	10	129	9	AD	7/2 ⁺			
²⁴³ Bk	58691	5			4.5 h	0.2 (3/2 ⁻)	93		$\beta^+ \approx 100$; $\alpha \approx 0.15$
²⁴³ Bk ^p	58740#	30#	50#	30#		(7/2 ⁻)			
²⁴³ Cf	60950#	140#			10.7 m	0.5 (1/2 ⁺)	93		$\beta^+ \approx 86$; $\alpha \approx 14$
²⁴³ Es	64780#	230#			21 s	2 3/2 ⁻ #	93		$\beta^+ \leq 70$; $\alpha \geq 30$
²⁴³ Es ^p	65180#	310#	400#	200#	am				
²⁴³ Fm	69260#	220#			210 ms	60 7/2 ⁻ #	93	ABBW D	$\alpha=60$ 40; β^+ ?; SF=0.57# *
* ²⁴³ Fm	D : $\alpha=40(20)$ % if α branching of ²³⁹ Cf is 100%, see ENSDF **								
²⁴⁴ Np	63200#	300#			2.29 m	0.16 (7 ⁻)	03		β^- =100
²⁴⁴ Pu	59806	5			80.0 My	0.9 0 ⁺	03	92Mo25 D	$\alpha \approx 100$; SF=0.121 4; ... *
²⁴⁴ Am	59881.0	2.1			10.1 h	0.1 6 ⁻ #	03		β^- =100
²⁴⁴ Am ^m	59969.5	2.3	88.6	1.7	RQ	26 m	1 1 ⁺	03	$\beta^- \approx 100$; $\epsilon=0.0361$ 13
²⁴⁴ Cm	58453.7	1.8			18.10 y	0.02 0 ⁺	03		$\alpha=100$; SF=1.37e-4 3
²⁴⁴ Cm ^m	59493.9	1.8	1040.188	0.012	34 ms	2 6 ⁺	03		IT=100

... A-group is continued on next page ...

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
... A-group continued ...								
²⁴⁴ Bk	60716	14	4.35 h	0.15	4 ⁻ #	03	$\beta^+ ?; \alpha=0.006\ 3$	
²⁴⁴ Bk ^p	60860#	50#			am			
²⁴⁴ Cf	61479.2	2.9	19.4 m	0.6	0 ⁺	03	$\alpha \approx 100; \epsilon ?$	
²⁴⁴ Es	66030#	180#	37 s	4		03	$\beta^+ = ?; \alpha = 5\ 3; \beta^+ \text{SF} = 0.01$	
²⁴⁴ Es ^p	66230#	240#			am			
²⁴⁴ Fm	69010#	280#	3.3 ms	0.5	0 ⁺	03	SF $\approx 100; \alpha = 0.4\#$	
* ²⁴⁴ Pu	D : ... ; $2\beta^- < 7.3e-9$							**
* ²⁴⁴ Pu	T : and $T(2\beta^-) > 1.1\ \text{Ey}$, from ⁹² Mo25; thus $2\beta^- < 7.3\ e-9\%$							**
²⁴⁵ Pu	63106	14	10.5 h	0.1	(9/2 ⁻)	93	$\beta^- = 100$	
²⁴⁵ Am	61900	3	2.05 h	0.01	(5/2 ⁺)	93	$\beta^- = 100$	
²⁴⁵ Cm	61004.7	2.1	8.5 ky	0.1	7/2 ⁺	93	$\alpha = 100; \text{SF} = 6.1e-7\ 9$	
²⁴⁵ Cm ^m	61360.6	2.1	290 ns	20	1/2 ⁺	93	IT=100	
²⁴⁵ Bk	61815.4	2.3	4.94 d	0.03	3/2 ⁻	93	$\epsilon \approx 100; \alpha = 0.12\ 1$	
²⁴⁵ Bk ^p	61870#	30#			(7/2 ⁻)			
²⁴⁵ Cf	63386.9	2.9	45.0 m	1.5	(5/2 ⁺)	93	$\beta^+ = 64\ 3; \alpha = 36\ 3$	
²⁴⁵ Cf ^p	63540#	100#			7/2 ⁺			
²⁴⁵ Es	66440#	200#	1.1 m	0.1	(3/2 ⁻)	93	$\beta^+ = 60\ 10; \alpha = 40\ 10$	
²⁴⁵ Es ^p	66740#	220#			am			
²⁴⁵ Es ^q	66790#	250#			am			
²⁴⁵ Fm	70220#	280#	4.2 s	1.3	1/2 ⁺ #	93	$\alpha = ?; \beta^+ = 4.2\#; \text{SF} = 0.13\#$	
²⁴⁵ Md	75290#	320#	* 900 μs	250	1/2 ⁻ #	97	96Ni09 TJD SF = ?; $\alpha ?$	
²⁴⁵ Md ^m	75490#	310#	* 400 ms	200	(7/2 ⁺)	97	96Ni09 TJD $\alpha = ?; \beta^+ ?$	
²⁴⁶ Pu	65395	15	10.84 d	0.02	0 ⁺	98	$\beta^- = 100$	
²⁴⁶ Am	64995	18	39 m	3	(7 ⁻)	98	$\beta^- = 100$	
²⁴⁶ Am ^m	65025	15	25.0 m	0.2	2 ⁽⁻⁾	98	$\beta^- \approx 100; \text{IT} < 0.02$	
²⁴⁶ Cm	62618.4	2.1	4.76 ky	0.04	0 ⁺	98	$\alpha \approx 100; \text{SF} = 0.02615\ 7$	
²⁴⁶ Bk	63970	60	1.80 d	0.02	2 ⁽⁻⁾	98	$\beta^+ \approx 100; \alpha = 0.1\#$	
²⁴⁶ Cf	64091.7	2.1	35.7 h	0.5	0 ⁺	98	$\alpha = 100; \text{SF} = 2.5e-4\ 2; \epsilon < 4e-3$	
²⁴⁶ Es	67900#	220#	7.7 m	0.5	4 ⁻ #	98	$\beta^+ = 90.1\ 18; \alpha = 9.9\ 18; \dots$	
²⁴⁶ Es ^p	68250#	300#			am			
²⁴⁶ Fm	70140	40	1.1 s	0.2	0 ⁺	98	96Ni09 D $\alpha = ?; \beta^+ > 10; \text{SF} = 4.5\ 13; \dots$	
²⁴⁶ Md	76280#	330#	1.0 s	0.4		98	$\alpha = ?; \beta^+ ?; \text{SF} ?$	
²⁴⁶ Md ^m	76490#	340#	1.0 s	0.4		96Ni09	TD $\alpha = ?; \beta^+ ?$	
* ²⁴⁶ Es	D : ... ; $\beta^+ \text{SF} \approx 0.003$							**
* ²⁴⁶ Fm	D : ... ; $\beta^+ \text{SF} = 10\ 5$							**
* ²⁴⁶ Md ^m	I : no longer considered to exist, see ENSDF'98							**
²⁴⁷ Pu	69000#	300#	2.27 d	0.23	1/2 ⁺ #	93	$\beta^- = 100$	
²⁴⁷ Am	67150#	100#	23.0 m	1.3	5/2#	93	$\beta^- = 100$	
²⁴⁷ Cm	65534	4	15.6 My	0.5	9/2 ⁻	93	$\alpha = 100$	
²⁴⁷ Bk	65491	6	1.38 ky	0.25	(3/2 ⁻)	93	$\alpha \approx 100; \text{SF} ?$	
²⁴⁷ Cf	66137	8	3.11 h	0.03	7/2 ⁺ #	93	$\epsilon \approx 100; \alpha = 0.035\ 5$	
²⁴⁷ Es	68610#	30#	4.6 m	0.3	7/2 ⁺ #	93	$\beta^+ \approx 93; \alpha \approx 7; \text{SF} \approx 9e-5\#$	
²⁴⁷ Es ^p	68930#	200#			am			
²⁴⁷ Fm	71580#	140#	35 s	4	5/2 ⁺ #	93	$\alpha \geq 50; \beta^+ \leq 50$	
²⁴⁷ Fm ^m		non existent	9.2 s	2.3		93	67F115 I $\alpha \approx 100; \text{IT} ?$	
²⁴⁷ Fm ^p	71730#	170#			(7/2 ⁺)			
²⁴⁷ Fm ^q	71980#	210#						
²⁴⁷ Md	76040#	320#	* 270 ms	160	1/2 ⁻ #	93	93Ho.A TD SF = ?; $\alpha ?$	
²⁴⁷ Md ^m	76170#	310#	Nm * 1.12 s	0.22	(7/2 ⁺)	93	93Ho.A TD $\alpha = 100; \text{SF} = 0.0001\#$	
* ²⁴⁷ Fm ^m	I : existence of this isomer is discussed in ENSDF							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²⁵² Cm	79060# 300#		< 1 d	0 ⁺	99		β^- ?
²⁵² Bk	78530# 200#		1.8 m 0.5		99	92Kr.A TD	β^- ?; α ?
²⁵² Cf	76034 5		2.645 y 0.008	0 ⁺	99		α =96,908 8; SF=3.092 8
²⁵² Es	77290 50		471.7 d 1.9	(5 ⁻)	99		α =78 2; ϵ =22 2
²⁵² Fm	76817 6		25.39 h 0.04	0 ⁺	99		α \approx 100; SF=0.0023 2; 2 β^+ ?
²⁵² Md	80630# 200#		2.3 m 0.8		99		β^+ >50; α <50
²⁵² Md ^p	80670# 220#	40# 100#		<i>am</i>			
²⁵² No	82881 13		2.44 s 0.04	0 ⁺	99	01Og08 TD	α \approx 67; SF=32.2 5; β^+ ?
²⁵² Lr	88840# 250#		390 ms 90		99	01He35 TD	β^+ =71#; α =?; SF<1
²⁵² Lr ^p	89140# 290#	300# 150#					
* ²⁵² No	T : other 03Be18=2.38(+0.26-0.22)		D : SF from 01Og08; α estimated by NUBASE				**
²⁵³ Bk	80930# 360#		10# m			91Kr.A I	β^- ?
²⁵³ Cf	79301 6		17.81 d 0.08	(7/2 ⁺)	99		β^- \approx 100; α =0.31 4
²⁵³ Es	79013.7 2.6		20.47 d 0.03	7/2 ⁺	99		α =100; SF=8.7e-6 3
²⁵³ Fm	79350 4		3.00 d 0.12	(1/2) ⁺	99		ϵ =88 1; α =12 1
²⁵³ Md	81300# 210#		12 m 8	7/2 ⁻ #	99		β^+ \approx 100; α =0.6#
²⁵³ Md ^p	81300# 210#	0# 30#		<i>am</i>			
²⁵³ No	84470# 100#		1.62 m 0.15	9/2 ⁻ #	99		α =?; β^+ =20#; SF=0.001#
²⁵³ No ^m	84590# 100#	129 19 AD	31 μ s	5/2 ⁺ #			α =?
²⁵³ Lr	88690# 220#		* & 580 ms 70	(7/2 ⁻)	99	01He35 TJD	α =90 10; SF=2.6 21; β^+ =1#
²⁵³ Lr ^m	88710# 250#	30# 100#	* & 1.5 s 0.3	(1/2 ⁻)	99	01He35 TJD	α =90 10; SF=8 5; β^+ =1#
²⁵³ Rf	93790# 450#		* 13 ms 5	(7/2) ^(+#)		95Ho.B TJ	SF \approx 50; α \approx 50
²⁵³ Rf ^m	93990# 470#	200# 150#	* 52 μ s 14	(1/2) ^(-#)	99	97He29 J	SF=?; α =5#
* ²⁵³ Bk	I : possible identification, in 91Kr.A. Needs confirmation						**
* ²⁵³ Rf	I : the state with \approx 1.8 s reported in ENSDF is not confirmed						**
²⁵⁴ Bk	84390# 300#		1# m				β^- ?
²⁵⁴ Cf	81341 12		60.5 d 0.2	0 ⁺	01		SF \approx 100; α =0.31 2; 2 β^- ?
²⁵⁴ Es	81992 4		275.7 d 0.5	(7 ⁺)	01		α \approx 100; ϵ =0.03#; ...
²⁵⁴ Es ^m	82076 3	84.2 2.5 AD	39.3 h 0.2	2 ⁺	01		β^- =98 2; IT<3; α =0.32 1; ...
²⁵⁴ Fm	80904.2 2.8		3.240 h 0.002	0 ⁺	01		α \approx 100; SF=0.0592 3
²⁵⁴ Md	83510# 100#		* 10 m 3	(0 ⁻)	01		β^+ \approx 100; α ?
²⁵⁴ Md ^m	83560# 140#	50# 100#	* 28 m 8	(3 ⁻)	01		β^+ \approx 100; α ?
²⁵⁴ No	84724 18		51 s 10	0 ⁺	01		α =90 4; β^+ =10 4; SF=0.17 5
²⁵⁴ No ^m	85220# 100#	500# 100#	280 ms 40		01		IT>80; α ?
²⁵⁴ Lr	89850# 340#		13 s 3		01		α =76 11; β^+ =24 11; SF ?
²⁵⁴ Lr ^p	89880# 340#	30# 70#					
²⁵⁴ Rf	93320# 290#		23 μ s 3	0 ⁺	01	97He29 TD	SF=?; α <1.5
* ²⁵⁴ Es	D : ... ; β^- =1.74e-4 8; SF<3e-6						**
* ²⁵⁴ Es ^m	D : ... ; ϵ =0.076 7; SF<0.045						**
* ²⁵⁴ Lr	T : also 01Ga20=13.4(4.2)						**
²⁵⁵ Cf	84810# 200#		85 m 18	(7/2 ⁺)	99		β^- =100; SF<0.001#; α =2e-7#
²⁵⁵ Es	84089 11		39.8 d 1.2	(7/2 ⁺)	99		β^- =92.0 4; α =8.0 4; SF=0.0041 2
²⁵⁵ Fm	83799 5		20.07 h 0.07	7/2 ⁺	99		α =100; SF=2.4e-5 10
²⁵⁵ Fm ^p	84050# 100#	250# 100# Nm		(9/2 ⁺)			
²⁵⁵ Md	84843 7		27 m 2	(7/2 ⁻)	99		β^+ =92 2; α =8 2; SF<0.15
²⁵⁵ Md ^p	84850# 70#	10# 70#		<i>am</i>			
²⁵⁵ No	86854 10		3.1 m 0.2	(1/2 ⁺)	99		α =61 3; β^+ =39 3
²⁵⁵ No ^p	86950# 70#	100# 70# Nm		(7/2 ⁺)			
²⁵⁵ Lr	90060# 210#		22 s 4	7/2 ⁻ #	99		α =?; β^+ <30#; SF<1#
²⁵⁵ Rf	94400# 180#		* 1.64 s 0.11	9/2 ⁻ #	99	01He35 TD	α =?; SF=52 6
²⁵⁵ Rf ^m	94320# 210#	-80# 180#	* 1.0 s 0.4	5/2 ⁺ #	99	97He29 D	α =100
²⁵⁵ Db	100040# 420#		1.7 s 0.5		99		α ?; SF \approx 20
* ²⁵⁵ Lr	T : also 01Ga20=21(8)						**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
²⁵⁶ Cf	87040#	300#	12.3 m	1.2	0 ⁺	99	SF=100; $\alpha=6.2e-7\#$; $2\beta^-$?	
²⁵⁶ Es	87190#	100#	25.4 m	2.4	(1 ⁺ , 0 ⁻)	99	β^- =100	
²⁵⁶ Es ^m	87190#	140#	7.6 h		(8 ⁺)	99	$\beta^- \approx 100$; β^- -SF=0.002	
²⁵⁶ Fm	85486	7	157.6 m	1.3	0 ⁺	99	SF=91.9 3; $\alpha=8.1$ 3	
²⁵⁶ Md	87620	50	77 m	2	(1 ⁻)	99	$\beta^+=?$; $\alpha=9.2$ 7; SF<3	
²⁵⁶ Md ^p	87700#	110#	80#	100#			am	
²⁵⁶ No	87824	8	2.91 s	0.05	0 ⁺	99	$\alpha \approx 100$; SF=0.53 6; $\epsilon < 0.01\#$	
²⁵⁶ Lr	91870#	220#	27 s	3		99	$\alpha=85$ 10; $\beta^+=15$ 10; SF<0.03	
²⁵⁶ Lr ^p	91970#	230#	100	70	XL			
²⁵⁶ Rf	94236	24	6.45 ms	0.14	0 ⁺	99	97He29 TD SF=?; $\alpha=0.32$ 17	*
²⁵⁶ Db	100720#	290#	1.9 s	0.4		99	01He35 TD $\alpha=?$; $\beta^+=36$ 12; SF=?	*
* ²⁵⁶ Rf	T : average 97He29=6.2(0.2) 84Og02=6.7(0.2)							**
* ²⁵⁶ Db	T : average 01He35=1.6(+0.5-0.3) 83Og.A=2.6(+1.4-0.8)							**
²⁵⁷ Es	89400#	410#	7.7 d	0.2	7/2 ⁺ #	99	β^- =100; $\alpha=4e-4\#$	
²⁵⁷ Fm	88589	6	100.5 d	0.2	(9/2 ⁺)	99	$\alpha \approx 100$; SF=0.210 4	
²⁵⁷ Md	88996.2	2.8	5.52 h	0.05	(7/2 ⁻)	99	$\epsilon=85$ 3; $\alpha=15$ 3; SF<4	
²⁵⁷ No	90241	22	25 s	2	(7/2 ⁺)	99	02Ho11 D $\alpha=?$; $\beta^+=15$ 8	
²⁵⁷ No ^p	90550#	110#	310#	100#			am	
²⁵⁷ Lr	92740#	210#	646 ms	25	9/2 ⁺ #	99	$\alpha \approx 100$; $\beta^+=0.01\#$; SF=0.001#	
²⁵⁷ Lr ^p	92890#	230#	150#	100#			am	
²⁵⁷ Rf	95930#	100#	4.7 s	0.3	(1/2 ⁺)	99	97He29 JD $\alpha=?$; $\beta^+=11$ 1; SF<1.4	
²⁵⁷ Rf ^m	96050#	100#	114	17	AD	99	97He29 EJ $\alpha \approx 100$; SF=0.7#; β^+ ?	*
²⁵⁷ Rf ^p	96030#	120#	100#	70#			(7/2 ⁺)	
²⁵⁷ Db	100340#	230#	* & 1.53 s	0.17	(9/2 ⁺)	99	01He35 TJD $\alpha > 94$; SF<6; $\beta^+=1\#$	
²⁵⁷ Db ^m	100450#	250#	* & 790 ms	130	(1/2 ⁻)	99	01He35 TJD $\alpha > 87$; SF<13; $\beta^+=1\#$	
* ²⁵⁷ Rf ^m	E : 97He29=118(4) keV form direct comparison of two alpha lines							**
²⁵⁸ Es	92700#	300#	3#	m			β^- ?; α ?	
²⁵⁸ Fm	90430#	200#	370 μ s	14	0 ⁺	01	86Hu05 T SF \approx 100; α ?	*
²⁵⁸ Md	91688	5	51.5 d	0.3	8 ⁻ #	01	93Mo18 D $\alpha \approx 100$; $\beta^+ < 0.0015$; $\beta^- < 0.0015$	*
²⁵⁸ Md ^m	91690#	200#	* 57.0 m	0.9	1 ⁻ #	01	93Mo18 D $\epsilon=?$; SF<20; $\beta^- < 10\#$; $\alpha < 1.2$	*
²⁵⁸ No	91480#	200#	1.2 ms	0.2	0 ⁺	01	SF \approx 100; $\alpha=0.001\#$; $2\beta^+$?	
²⁵⁸ Lr	94840#	100#	4.1 s	0.3		01	$\alpha > 95$; $\beta^+ < 5$	
²⁵⁸ Lr ^p	95040#	180#	200#	150#			am	
²⁵⁸ Rf	96400#	200#	12 ms	2	0 ⁺	01	SF=87 2; $\alpha=13$ 2	
²⁵⁸ Db	101750#	340#	* 4.5 s	0.6		01	$\alpha=64$ 7; $\beta^+=36$ 7; SF<1#	
²⁵⁸ Db ^m	101810#	350#	* 20 s	10		01	$\beta^+ \approx 100$; IT ?	
²⁵⁸ Sg	105420#	410#	3.3 ms	1.0	0 ⁺	01	SF=?; $\alpha < 20$	
* ²⁵⁸ Fm	T : average 86Hu05=360(20) 71Hu03=380(20) (all 1 σ) ENSDF gives 3 σ							**
* ²⁵⁸ Md	D : derived from: "the sum of SF, ϵ and β^- decay branches < 0.003%" in							**
* ²⁵⁸ Md	D : 93Mo18 and T(SF)>150000 y, from 86Lo16, thus SF<1e-4#							**
* ²⁵⁸ Md ^m	D : SF<20% derived from 93Mo18 "the sum of SF and β^- decay branches < 30%"							**
²⁵⁹ Fm	93700#	280#	1.5 s	0.3	3/2 ⁺ #	99	SF=100	
²⁵⁹ Md	93620#	200#	1.60 h	0.06	7/2 ⁻ #	99	93Mo18 T SF=?; $\alpha < 1.3$	
²⁵⁹ No	94110#	100#	58 m	5	9/2 ⁺ #	99	$\alpha=75$ 4; $\epsilon=25$ 4; SF<10	
²⁵⁹ No ^p	94390#	180#	280#	150#				
²⁵⁹ Lr	95850#	70#	6.2 s	0.3	9/2 ⁺ #	99	$\alpha=78$ 2; SF=22 2; $\beta^+=0.6\#$	
²⁵⁹ Lr ^p	96200#	170#	350#	150#				
²⁵⁹ Rf	98400#	70#	2.8 s	0.4	7/2 ⁺ #	99	94Gr08 T $\alpha=92$ 2; SF=8 2; $\beta^+=0.3\#$	*
²⁵⁹ Rf ^p	98500#	100#	100#	70#	Nm		(3/2 ⁺)	
²⁵⁹ Rf ^l	98610#	130#	210#	110#	Nm		(9/2 ⁺)	
²⁵⁹ Db	102100#	210#	510 ms	160		99	01Ga20 TD $\alpha=100$	
²⁵⁹ Sg	106660#	180#	580 ms	210	1/2 ⁺ #	99	$\alpha=90$ 10; SF<20	
* ²⁵⁹ Rf	T : average 94Gr08=1.7(+0.8-0.5) 85So03=3.4(1.7) 81Be03=3.0(1.3)							**
* ²⁵⁹ Rf	T : 73Dr10=3.2(0.8) and 69Gh01=3.2(0.8)							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{260}Fm	95640# 500#	EU	1# m	0^+			SF ?	*
^{260}Md	96550# 320#		27.8 d	0.8	99	92Lo.B TD	SF=?; $\alpha < 5$; $\epsilon < 5$; $\beta^- < 3.5$	*
^{260}No	95610# 200#		106 ms	8	99		SF=100	
^{260}Lr	98280# 120#		3.0 m	0.5	99		$\alpha=80\ 20$; $\beta^+=20\ 20$	
^{260}Rf	99150# 200#		21 ms	1	99		SF=?; $\alpha=2\#$; $\epsilon=0.01\#$	
^{260}Db	103680# 230#		1.52 s	0.13	99		$\alpha \geq 90.4\ 6$; SF $\leq 9.6\ 6$; $\beta^+ < 2.5$	
$^{260}\text{Db}^p$	103880# 280# 200# 150#							
^{260}Sg	106580 40		3.8 ms	0.8	99		SF=60 30; $\alpha=40\ 30$	
^{260}Bh	113610# 580#		300# μs		99		$\alpha=100$	
* ^{260}Fm	I: half-life ≈ 4 ms and SF=100 mode were reported in the 92Lo.B internal							**
* ^{260}Fm	I: report. Not confirmed in subsequent experiment by same group (97Lo.A)							**
* ^{260}Fm	I: Discovery of this nuclide is considered unproven							**
* ^{260}Md	T: supersedes 86Hu01=31.8(0.5) of same group							**
^{261}Md	98480# 650#		40# m	$7/2^- \#$			$\alpha ?$	
^{261}No	98500# 300#		3# h	$3/2^+ \#$			$\alpha ?$	
^{261}Lr	99560# 200#		39 m	12	99		SF=?; $\alpha ?$	
^{261}Rf	101315 29		* & 5.5 s	2.5	99	02Ho11 T	$\alpha=?$; SF=40	
$^{261}\text{Rf}^m$	101390# 100# 70# 100#		* & 81 s	9	99	02Ho11 TD	$\alpha=?$; $\beta^+ < 15$; SF<10	
$^{261}\text{Rf}^p$	101420 70 100 60 AD						$3/2^+ \#$	
^{261}Db	104380# 230#		1.8 s	0.4	99		$\alpha > 82$; SF<18	
^{261}Sg	108160# 130#		230 ms	60	99		$\alpha \approx 100$; SF<1	
$^{261}\text{Sg}^p$	108290# 140# 130 50 AD						$(9/2^+)$	
$^{261}\text{Sg}^q$	108320# 140# 160 50 AD						$(3/2^+)$	
^{261}Bh	113330# 230#		13 ms	4	99		$\alpha=95\ 5$; SF<10	
^{262}Md	101410# 580#		3# m				SF ?; $\alpha ?$	
^{262}No	99950# 450#		5 ms	0^+	01		SF ≈ 100 ; $\alpha ?$	
^{262}Lr	102120# 200#		4 h		01		$\beta^+ = ?$; SF<10; $\alpha ?$	
^{262}Rf	102390# 280#		* 2.3 s	0.4	01		SF ≈ 100 ; $\alpha < 0.8$	
$^{262}\text{Rf}^m$	102990# 490# 600# 400#		* 47 ms	5	high	96La11 I	SF=100	*
^{262}Db	106270# 180#		35 s	5	01		$\alpha \approx 67$; SF ≈ 30 ; $\beta^+ = 3\#$	
$^{262}\text{Db}^p$	106390# 200# 120# 70#						$\alpha ?$	
^{262}Sg	108420# 280#		8 ms	3	0^+	01 01Ho06 TD	SF=?; $\alpha < 22$	
^{262}Bh	114470# 350#		290 ms	160	01	97Ho14 T	$\alpha=?$; SF<20	*
$^{262}\text{Bh}^m$	114780# 350# 300 60 AD		14 ms	4	01	97Ho14 T	$\alpha=?$; SF<10	*
* $^{262}\text{Rf}^m$	I: assigned by 96La11 to K-isomeric state							**
* ^{262}Bh	T: 3 events at 225, 255 and 278 ms yielding 175(+240–64), see 84Sc13							**
* $^{262}\text{Bh}^m$	T: 11 events yielding 12.2(+5.5–2.8)							**
^{263}No	102980# 490#		20# m				$\alpha ?$; SF ?	
^{263}Lr	103670# 360#		5# h				$\alpha ?$	
^{263}Rf	104840# 180#		11 m	3	$3/2^+ \#$	99 93Gr.C TD	SF=?; $\alpha=30$	*
^{263}Db	107110# 170#		29 s	9	99	92Kr01 D	SF=56 14; $\alpha=?$; $\beta^+ = 6.9\ 16$	*
$^{263}\text{Db}^p$	107510# 260# 400# 200#							
^{263}Sg	110220# 120#		1.0 s	0.2	$9/2^+ \#$	99	$\alpha > 70$; SF ?	
$^{263}\text{Sg}^m$	110320# 100# 100# 70# Nm *		120 ms		$3/2^+ \#$	99	$\alpha=?$; IT ?	
^{263}Bh	114610# 370#		200# ms			99	$\alpha ?$	
^{263}Hs	119750# 350#		1# ms		$7/2^+ \#$	99	$\alpha=100$	
$^{263}\text{Hs}^p$	120250# 360# 500# 100#				am		$\alpha ?$; SF ?	
* ^{263}Rf	T: average 03Kr.1=24(+19–7) m 93Gr.C=500(+300–200) s 92Cz.A=600(+300–200) s							**
* ^{263}Db	D: SF from 92Kr01=57(+13–15); β^+ average 03Kr.1=3(+4–1) 93Gr.C=8(2)							**
* ^{263}Db	T: Possibly a candidate for the 54(+98–21) s SF decay observed by 98Ik02							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
^{264}No	104650# 640#		1# m	0^+			α ?; SF ?	
^{264}Lr	106230# 440#		10# h				α ?; SF ?	
^{264}Rf	106180# 450#		1# h	0^+			α ?	
^{264}Db	109360# 230#		3# m				α ?	
^{264}Sg	110780# 280#		400# ms	0^+	99		α ?	
^{264}Bh	116070# 280#		1.3 s	0.5	99	02Ho11 T	α =?; β^+ ?	
$^{264}\text{Bh}^p$	116370# 310#	300# 150#					*	
^{264}Hs	119600 40		540 μs	300	99	95Ho.B T	α ≈50; SF≈50	
* ^{264}Bh	T: mean lifetime of 6 events 1.5 s							**
* ^{264}Hs	T: 95Ho.B (2 events 76 μs and 825 μs) 87Mu15 (1 event 80 μs). Average of							**
* ^{264}Hs	T: the 3 events: 327(+448–120) μs , see 84Sc13							**
^{265}Lr	107900# 710#		10# h				α ?; SF ?	
^{265}Rf	108710# 420#		13 h	$3/2^+$ #	00	99Og.A TD	α ?	
^{265}Db	110480# 280#		15# m				α ?	
^{265}Sg	112820 60		8 s	3	99		α >50; SF ?	
$^{265}\text{Sg}^p$	113120# 120#	300# 100#					$11/2^-$ #	
^{265}Bh	116570# 380#		500# ms				α ?	
^{265}Hs	121170# 140#		2.1 ms	0.3	99		α ≈100; SF<1	
$^{265}\text{Hs}^m$	121480# 140#	300 70 AD	780 μs	150	99		α ≈100; IT ?	
^{265}Mt	126820# 460#		2# ms				α ?	
* ^{265}Rf	T: one case only after a 1.3 h measurement							**
^{266}Lr	111130# 660#		1# h				α ?; SF ?	
^{266}Rf	109880# 540#		10# h	0^+			α ?; SF ?	
^{266}Db	112740# 360#		20# m				α ?; SF ?	
^{266}Sg	113700# 290#		21 s	6	01	98Tu01 T	α =34 9; SF=66 9	
^{266}Bh	118250# 200#		5 s	3	01		α ≈100; β^+ ?; SF ?	
^{266}Hs	121190# 280#		2.7 ms	1.0	01	01Ho06 TD	α =?; SF≈1.4#	
^{266}Mt	127890# 350#		1.2 ms	0.4	01	84Og03 D	α =?; SF<5.5	
$^{266}\text{Mt}^m$	129120# 350#	1230 80 AD	6 ms	3	01	97Ho14 TD	α =100	
* ^{266}Sg	T: average 98Tu01=21(+20–12) 94La22=10–30 D: from 18%< α <50% 50%<SF<82%							**
* ^{266}Bh	T: from $T=1-10$; estimated 1#s from systematics							**
* ^{266}Mt	T: 10 events yielding 1.01(+0.47–0.24)							**
* $^{266}\text{Mt}^m$	T: 3 events at 7.8, 2.0 and 5.0 yield 3.4(+4.7–1.3)							**
^{267}Rf	113200# 580#		5# h				α ?; SF ?	
^{267}Db	113990# 470#		2# h				α ?; SF ?	
^{267}Sg	115900# 270#		19 ms			99Og.B T	α =100	
^{267}Bh	118910# 260#		22 s	10		00Wi15 TD	α =100	
^{267}Hs	122760# 100#		32 ms	15	00		α =100	
$^{267}\text{Hs}^m$		non existent EU	200 ms			95Ho.A TDI	α =?; IT ?	
^{267}Mt	127900# 540#		10# ms				α ?	
^{267}Ea	134450# 370#		10 μs	8	00	95Gh04 T	α =100	
* $^{267}\text{Hs}^m$	I: tentative only							**
* ^{267}Ea	T: one single event, lifetime 4 μs , thus $T=2.8(+13.0-1.3)$, see 84Sc13							**
^{268}Rf	115170# 710#		1# h	0^+			α ?; SF ?	
^{268}Db	116850# 530#		6# h				α ?; SF ?	
^{268}Sg	117000# 540#		30# s	0^+			α ?; SF ?	
^{268}Bh	120870# 380#		25# s				α ?; SF ?	
^{268}Hs	123110# 410#		2# s	0^+			α ?	
^{268}Mt	129220# 320#		53 ms	21	00	02Ho11 T	α =100	
$^{268}\text{Mt}^p$	129470# 330#	250# 100#					α ?; SF ?	
^{268}Ea	133940# 500#		100# μs	0^+			α ?	
* ^{268}Mt	T: mean lifetime of 6 events 60 ms							**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)	
²⁶⁹ Db	118730# 770#		3# h				α ?; SF ?	
²⁶⁹ Sg	119930# 660#		35 s	23	00		$\alpha < 100$; SF ?	
²⁶⁹ Bh	121740# 410#		25# s				α ?	
²⁶⁹ Hs	124870# 120#		27 s	17	00	02Ho11 T	$\alpha = 100$ *	
²⁶⁹ Mt	129530# 550#		200# ms				α ?	
²⁶⁹ Ea	135180# 140#		230 μ s	110	3/2 ⁺ #	00 95Ho03 T	$\alpha = 100$	
* ²⁶⁹ Hs	T : 2 events at 19.7 and 22.0 s yield 14(+26–6)							**
²⁷⁰ Db	121760# 720#		1# h				α ?; SF ?	
²⁷⁰ Sg	121400# 620#		10# m	0 ⁺			α ?; SF ?	
²⁷⁰ Bh	124460# 470#		30# s				α ?; SF ?	
²⁷⁰ Hs	125430# 290#		30# s	0 ⁺		01Tu.B D	$\alpha = 100$	
²⁷⁰ Mt	131020# 540#		2# s				α ?	
²⁷⁰ Ea	134810# 290#		160 μ s	100	0 ⁺	01Ho06 TD	$\alpha \approx 100$; SF ≈ 0.2	
²⁷⁰ Ea ^m	135940# 290#	1140 70	10 ms	6	(10) ^(-#)	01Ho06 ETJ	$\alpha = ?$; IT ?	
²⁷¹ Sg	124330# 650#		2# h				α ?; SF ?	
²⁷¹ Bh	125920# 560#		40# s				α ?; SF ?	
²⁷¹ Hs	128230# 340#		40# s				α ?; SF ?	
²⁷¹ Mt	131470# 570#		5# s				α ?	
²⁷¹ Ea	136060# 110#		210 ms	170	11/2 ⁻ #	00	$\alpha = 100$	
²⁷¹ Ea ^m	136090# 110#	29 29 AD *	1.3 ms	0.5	9/2 ⁺ #	00	$\alpha = 100$	
²⁷² Sg	125900# 770#		1# h	0 ⁺			α ?; SF ?	
²⁷² Bh	128580# 610#		2# m				α ?; SF ?	
²⁷² Hs	129530# 580#		40# s	0 ⁺			α ?; SF ?	
²⁷² Mt	133890# 480#		10# s				α ?; SF ?	
²⁷² Ea	136290# 650#		1# s	0 ⁺			SF ?	
²⁷² Eb	143090# 330#		2.0 ms	0.8	5 ⁺ #, 6 ⁺ #	00 02Ho11 T	$\alpha = 100$ *	
* ²⁷² Eb	T : mean lifetime of 6 events 2.3 ms							**
²⁷³ Sg	128750# 660#		1# m				SF ?	
²⁷³ Bh	130050# 830#		90# m				α ?; SF ?	
²⁷³ Hs	132260# 830#	RN	50# s	3/2 ⁺ #	00	02Ni10 I	α ? *	
²⁷³ Mt	134990# 510#		20# s				α ?; SF ?	
²⁷³ Ea	138670# 130#		360 μ s	280	13/2 ⁻ #	00	$\alpha = 100$	
²⁷³ Ea ^m	138870# 130#	198 20 EU	120 ms		3/2 ⁺ #	00	$\alpha = 100$	
²⁷³ Ea ^p	138950# 130#	290 40 AD					α ?; SF ?	
²⁷³ Eb	143150# 610#		5# ms				α ?	
* ²⁷³ Hs	T : 99Ni03=1.2(+1.7–0.6) alpha decay retracted by authors in 02Ni10							**
²⁷⁴ Bh	132680# 780#		90# m				α ?; SF ?	
²⁷⁴ Hs	133330# 650#		1# m	0 ⁺			α ?; SF ?	
²⁷⁴ Mt	137390# 560#		20# s				α ?; SF ?	
²⁷⁴ Ea	139250# 490#		2# s	0 ⁺			α ?; SF ?	
²⁷⁴ Eb	145050# 620#		5# ms				α ?	
²⁷⁵ Bh	134370# 650#		40# m				SF ?	
²⁷⁵ Hs	135950# 710#		30# m				α ?; SF ?	
²⁷⁵ Mt	138460# 590#		30# s				α ?; SF ?	
²⁷⁵ Ea	141750# 450#		2# s				α ?; SF ?	
²⁷⁵ Eb	145450# 690#		10# ms				α ?	

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)				
²⁷⁶ Hs	137120#	820#	1#	h			α ?; SF ?				
²⁷⁶ Mt	140800#	680#	40#	s			α ?; SF ?				
²⁷⁶ Ea	142550#	610#	5#	s			α ?; SF ?				
²⁷⁶ Eb	147640#	630#	100#	ms			α ?; SF ?				
²⁷⁷ Hs	139580#	730#	40	m	30	3/2 ⁺ #	00	99Og10	TD	SF=100	*
²⁷⁷ Mt	141980#	880#	1#	m						α ?; SF ?	
²⁷⁷ Ea	144980#	960#	5#	s		11/2 ⁺ #	00	02Ni10	I	α ?	*
²⁷⁷ Eb	148590#	620#	1#	s						α ?; SF ?	
²⁷⁷ Ec	152710#	130#	1.1	ms	0.7	3/2 ⁺ #	00	02Ho11	T	α =100	*
* ²⁷⁷ Hs	T : one single event 16.5 m yields 11(+55–5)										**
* ²⁷⁷ Ea	T : 99Ni03=3.0(+4.7–1.5) alpha decay retracted by authors in 02Ni10										**
* ²⁷⁷ Ec	T : two events at 0.280 ms and 1.406 ms										**
²⁷⁸ Mt	144210#	840#	30#	m						α ?; SF ?	
²⁷⁸ Ea	145750#	680#	10#	s		0 ⁺				α ?; SF ?	
²⁷⁸ Eb	150530#	630#	1#	s						α ?; SF ?	
²⁷⁸ Ec	153060#	530#	10#	ms		0 ⁺				α ?; SF ?	
²⁷⁹ Mt	145490#	720#	6#	m						α ?; SF ?	
²⁷⁹ Ea	147980#	740#	10#	s						α ?; SF ?	
²⁷⁹ Eb	151340#	660#	3#	s						α ?; SF ?	
²⁷⁹ Ec	155140#	490#	100#	ms						α ?; SF ?	
²⁸⁰ Ea	148850#	850#	11	s	6	0 ⁺		01Og01	TD	SF=100	*
²⁸⁰ Eb	153210#	740#	10#	s						α ?; SF ?	
²⁸⁰ Ec	155600#	640#	1#	s		0 ⁺				α ?; SF ?	
* ²⁸⁰ Ea	T : 3 events at 6.93, 14.3 and 7.4 yield 6.6(+9–2.4)										**
²⁸¹ Ea	150960#	730#	4	m	3	3/2 ⁺ #	00	99Og10	TD	α =100	*
²⁸¹ Eb	154040#	930#	1#	m						α ?; SF ?	
²⁸¹ Ec	157690#	990#	10#	s		3/2 ⁺ #	00	02Ni10	I	α ?	*
* ²⁸¹ Ea	T : one single event 1.6 m yields 1.1(+5.3–0.5), see 84Sc13										**
* ²⁸¹ Ec	T : 99Ni03=0.89(+1.30–0.45) alpha decay retracted by authors in 02Ni10										**
²⁸² Eb	156010#	890#	4#	m						α ?; SF ?	
²⁸² Ec	158140#	710#	30#	s		0 ⁺				α ?; SF ?	
²⁸³ Eb	156880#	780#	10#	m						α ?; SF ?	
²⁸³ Ec	160020#	770#	4.2	m	2.1			99Og05	TD	SF=100	*
²⁸³ Ed	164360#	730#	10#	s						α ?; SF ?	
* ²⁸³ Ec	T : 4 events at 99Og07=9.3 m, 3.8 m, 99Og05=3.0 m and 0.9 m yield 3(+3–1) m										**
²⁸⁴ Ec	160570#	850#	31	s	18	0 ⁺		01Og01	TD	α =100	
²⁸⁴ Ed	165880#	800#	1#	m						α ?; SF ?	
²⁸⁵ Ec	162180#	730#	40	m	30	5/2 ⁺ #	00	99Og10	TD	α =100	*
²⁸⁵ Ed	166490#	980#	2#	m						α ?; SF ?	
²⁸⁵ Ee	171110#	1030#	5#	s		3/2 ⁺ #	00	02Ni10	I	α ?	*
* ²⁸⁵ Ec	T : one single event 15.4 s yields 11(+51–5), see 84Sc13										**
* ²⁸⁵ Ee	T : 99Ni03=580(+870–290) alpha decay retracted by authors in 02Ni10										**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	J^π	Ens	Reference	Decay modes and intensities (%)
²⁸⁶ Ed	168120#	940#	5#	m			$\alpha ?$; SF ?
²⁸⁶ Ee	171260#	770#	5#	s			$\alpha ?$; SF ?
²⁸⁷ Ed	168640#	830#	20#	m			$\alpha ?$; SF ?
²⁸⁷ Ee	172880#	770#	10	s	7	99Og07 T	$\alpha=100$ *
²⁸⁷ Ef	178090#	790#	500#	ms			$\alpha ?$; SF ?
* ²⁸⁷ Ee	T : 2 events at 1.32 s and 14.4 s yield 5.5(+10–2)						**
²⁸⁸ Ee	172970#	850#	2.8	s	1.4	01Og01 TD	$\alpha=100$
²⁸⁸ Ef	179310#	850#	1#	s			$\alpha ?$; SF ?
²⁸⁹ Ee	174450#	730#	80	s	60	5/2+# 00 99Og10 TD	$\alpha=100$ *
²⁸⁹ Ef	179510#	1020#	10#	s			$\alpha ?$; SF ?
²⁸⁹ Eg	185240#	1090#	10#	ms		5/2+# 00 02Ni10 I	$\alpha ?$ *
* ²⁸⁹ Ee	T : one single event at 30.4 s yields 21(+101–10)						**
* ²⁸⁹ Eg	T : 99Ni03=600(+860–300) alpha decay retracted by authors in 02Ni10						**
²⁹⁰ Ef	180840#	980#	10#	s			$\alpha ?$; SF ?
²⁹⁰ Eg	184990#	840#	50#	ms			$\alpha ?$; SF ?
²⁹¹ Ef	181070#	890#	1#	m			$\alpha ?$; SF ?
²⁹¹ Eg	186310#	850#	100#	ms			$\alpha ?$; SF ?
²⁹¹ Eh	192410#	880#	10#	ms			$\alpha ?$; SF ?
²⁹² Eg	186100#	850#	120	ms	100	01Og01 TD	$\alpha=100$ *
²⁹² Eh	193330#	940#	50#	ms			$\alpha ?$; SF ?
* ²⁹² Eg	T : one single event at 46.9 ms yields 33(+155–15)						**
²⁹³ Ei	199960#	1200#	RN	5#	ms	1/2+# 00 02Ni10 I	$\alpha ?$ *
* ²⁹³ Ei	T : 99Ni03=120(+180–60) alpha decay retracted by authors in 02Ni10						**